

## Study on Enterotoxigenic *Escherichia coli* Producing Extended Spectrum Beta Lactamase (ESBL) from Chicken Meat and its Products

Mohamed A. Hassan<sup>1</sup>, Gaber E. Batiha<sup>2</sup>, Saber A. Saad<sup>1</sup> and Eman Mahrous<sup>3\*</sup>

<sup>1</sup>Damanhur Branch, Animal Health Research Institute, Agriculture Research Center (ARC), Egypt

<sup>2</sup>Department of Pharmacology and Therapeutics, Faculty of Veterinary Medicine, Damanhour University, Damanhour, 22511, Egypt

<sup>3</sup>Bacteriology Department, Animal Health Research Institute, Agriculture Research Center (ARC), Egypt

\*Corresponding author: [eman\\_mahrous12@yahoo.com](mailto:eman_mahrous12@yahoo.com)

Article History: 22-726

Received: 18-Oct-22

Revised: 07-Dec-22

Accepted: 11-Dec-22

### ABSTRACT

This study aimed to identify the prevalence of enterotoxigenic *E. coli* producing ESBL in some chicken meat and chicken meat products. A total of 300 samples, including 120 chicken meat samples, 180 nuggets, frankfurter and luncheon (60 each) were taken from different supermarkets of different sanitation levels. These samples were subjected to bacteriological examination. The results showed that the prevalence of *E. coli* in chicken meat, nuggets, frankfurter and luncheon was 33.33, 13.3, 16.66 and 23.33%, respectively. In addition, the mean values of Aerobic plate count in chicken meat, nuggets, frankfurter and luncheon were  $2.05 \times 10^5 \pm 0.43 \times 10^4$ ,  $3.4 \times 10^3 \pm 0.22 \times 10^3$ ,  $4.7 \times 10^3 \pm 0.11 \times 10^3$  and  $6.5 \times 10^3 \pm 0.15 \times 10^3$  cfu/g; and the mean values of Coliform count were  $2.4 \times 10^3 \pm 0.11 \times 10^3$ ,  $6.3 \times 10^2 \pm 0.36 \times 10^2$ ,  $8.24 \times 10^2 \pm 0.38 \times 10^2$  and  $8.5 \times 10^2 \pm 0.34 \times 10^2$  cfu/g, respectively. *E. coli* isolates were serogrouped into O<sub>86</sub>:k<sub>61</sub>, O<sub>119</sub>:k<sub>69</sub>, O<sub>55</sub>:k<sub>59</sub>, O<sub>125</sub>:k<sub>70</sub>, O<sub>25</sub>:k<sub>11</sub>, O<sub>128</sub>:k<sub>67</sub>, O<sub>78</sub>:k<sub>80</sub> and untyped serogroups. Antimicrobial susceptibility testing of most isolates revealed the presence of multidrug-resistant strains. In addition, virulence genes (*Stx*, *Lt* and *Stx2* genes) were detected in 2, 1 and 4 isolates; and antibiotic resistance genes (*bla* TEM, *Bla* CTX-M and *AmpC* genes) were detected in 100% of isolates indicating high resistance to beta-lactamase antibiotics.

**Key words:** *E. coli*, ESBL, Chicken Meat, Chicken meat Products, Virulence Genes, Antibiotic resistance.

### INTRODUCTION

People of all ages get nutritional benefits from chicken and chicken products because it contains all the essential amino acids, high level of unsaturated fatty acids and low cholesterol level. Furthermore, vitamins, including niacin, riboflavin, thiamine, and ascorbic acid are abundant in chicken meat (Cartoni et al. 2022). On the other hand, chicken meat is a common source of many food-borne bacteria such as *Escherichia coli*, *Salmonellae* and coagulase-positive *Staph aureus* (Gonçalves-Tenório et al. 2018). The carcass could be contaminated at several points throughout the processing operation; during scalding, de-feathering, evisceration, processing equipment, improper cooking, refrigeration or storage (Gaafar et al. 2019).

*Escherichia coli* is regularly recovered from the digestive tracts of domestic and wild animals as well as humans. It is one of the most important foods borne

infections of public health significance incriminated in poultry meat and meat products worldwide (Samy et al. 2022). Enterotoxigenic *E. coli* infections cause abdominal cramps, low-grade fever, watery diarrhea, and nausea (Yang et al. 2017). It produces two types of enterotoxins: heat-stable (ST) and heat-labile (LT) each has a different function in pathogenesis.

Antibiotic usage in food-producing animals is a public health concern because antibiotic-resistant zoonotic infections could be transmitted to humans through the consumption and/or handling of animal-derived food (Adzitey et al. 2021).

Misuse of antimicrobial medications and poor infection control methods hastened the spread of antimicrobial resistance (El Jalil et al. 2020; WHO 2021). Infections with extended-spectrum Beta-lactamase (ESBL)-producing *Enterobacteriaceae* become a significant problem Worldwide Badr et al. (2022). The existence of ESBL genes in food-producing animals

**Cite This Article as:** Hassan MA, Batiha GE, Saad SA and Mahrous E, xxxx. Study on enterotoxigenic *Escherichia coli* producing extended spectrum beta lactamase (ESBL) from chicken meat and its products. International Journal of Veterinary Science x(x): xxxx. <https://doi.org/10.47278/journal.ijvs/2022.217>

indicates a potential hazard for animals (Moawad et al. 2018) and humans (Rahman et al. 2020) which highlights the importance of continuous monitoring.

This study was planned to investigate the prevalence of enterotoxigenic *E. coli* producing ESBL in chicken meat and its products. Moreover, monitoring the sensitivity to various antimicrobial agents. Also, the presence of virulence and antibiotic resistance genes were determined using PCR.

## MATERIALS AND METHODS

### Ethical approval

The study was conducted at Bacteriology Department, Animal Health Research Institute, Dokki, Giza, Egypt. The experimental protocol was approved by the Institutional Animal Care and Use Committee, Agriculture Research Center, Giza, Egypt.

### Sampling

A total of Three hundred samples including 120 chicken meat and 180 chicken nuggets, frankfurter and luncheon samples (60 each) were randomly collected from different supermarkets and retailers of different sanitation levels in different cities in El-Behera governorate, Egypt. Samples were collected aseptically, packed, identified and transferred in cooling iceboxes to the laboratory for bacteriological examination.

### Preparation of the samples according to (APHA 2001):

Twenty-five grams of each examined samples were transferred to a septic blender jar, and 225 ml of 0.1% sterile peptone water was aseptically added to the jar's contents. Each sample was then homogenized in the stomacher at 2000rpm for 1-2min to provide a homogenate from which ten-fold serial dilutions were prepared.

### Determination of Aerobic plate count according to APHA (2001):

It was conducted using standard plate count agar media.

**Determination of Coliform count according to ISO 4832 (2006):** Violet red bile agar medium has been used.

### Isolation and identification of *E. coli* according to Quinn et al. (2011):

Samples were inoculated in buffered peptone water and incubated at 37°C for 18 to 24h under aerobic conditions. A loopful from the enriched broth was streaked onto MacConkey's agar and eosin methylene blue agar (EMB) and incubated at 37°C for 24h. The metallic green colonies were picked up and identified biochemically and serologically. Media and biochemical test reagents were purchased from Hi-media®.

### Serogrouping of *E. coli* isolate according to Quinn et al. (2011):

*E. coli* isolates were serogrouped by slide agglutination test using polyvalent and monovalent antisera (Denka Seiken Co. LTD, Tokyo, Japan for antisera).

### Antibiotic sensitivity of *E. coli* isolates according to CLSI (2022):

Antibiogram of *E. coli* isolates was performed by disc diffusion method using available antibiotic discs purchased from MAST® Diagnostics. As the antibiotic discs used in the study were Levofloxacin (5µ), Amikacin (30µg), Gentamycin (5µg), Amoxicillin (25µg), Oxytetracycline (30µg), Vancomycin (30µg), Cefotaxime (30µg), Ampicillin (30µg), Ciprofloxacin (10µg) and Penicillin G (10µg). Antibiotic sensitivity in relation to zone of inhibition was interpreted by the Manufacturing Company.

### Molecular characterization of *E. coli* isolates

**DNA extraction:** The QIAamp DNA Mini kit (Qiagen, Germany, GmbH) was used to extract DNA from samples with slight modifications to the manufacturer's recommendations. Briefly, 200µL of the sample suspension was incubated with 10µL of proteinase K and 200µL of lysis buffer at 56°C for 10min. After incubation, 200µL of 100% ethanol was added to the lysate. The sample was then washed and centrifuged following the manufacturer's recommendations. Nucleic acid was eluted with 100µL of elution buffer provided in the kit.

**Table 1:** Primers sequences, target genes, amplicon sizes and cycling conditions for conventional PCR

Target Gene	Primers sequences	Product size(bp)	Amplification (35 cycles)				Final extension	References
			Primary Denaturation	Secondary denaturation	Annealing	Extension		
<i>phoA</i>	CGATTCTGGAAATGGCAAAG CGTGATCAGCGGTGACTATGAC	720bp	94°C 5min	94°C 30S	55°C 40S	72°C 45S	72°C 10min	Hu et al. (2011)
<i>blaTEM</i>	ATCAGCAATAAACCAGC CCCCGAAGAACGTTTTTC	516bp	94°C 5min.	94°C 30S	54°C 40S	72°C 45S	72°C 10min	Colom et al. (2003)
<i>ampC</i>	TTCTATCAAMACTGGCARCC CCYTTTTATGTACCCAYGA	550bp	94°C 5S	94°C 30S	50°C 40S	72°C 45S	72°C 10min.	Srinivasan et al. (2005)
<i>blaCTX-M</i>	ATG TGC AGY ACC AGT AAR GTK ATG GC TGG GTR AAR TAR GTS ACC AGA AYC AGC GG	593bp	94°C 5min	94°C 30S	54°C 40S	72°C 45S	72°C. 10min	Archambault et al. (2006)
<i>Stx2</i>	CCATGACAACGGACAGCAGTT CCTGTCAACTGAGCAGCACTT TG	779bp	94°C 5min	94°C 30S	58°C 40S	72°C 45S	72°C 10min	Dipineto et al. (2006)
<i>Stx/St1</i>	GAAACAACATGACGGGAGGT GCACAGGCAGGATTACAACA	229bp	94°C 5min	94°C 30S	57°C 30S	72°C 30S	72°C 7min	Lee et al. (2008)
<i>Lt</i>	GGTTTCTGCGTTAGGTGGAA GGGACTTCGACCTGAAATGT	606bp	94°C 5min	94°C 30S	57°C 40S	72°C 45S	72°C 10min	

*phoA*=Alkaline Phosphatase (Fig.1), *Stx2*=Shiga toxins 2 (Fig. 2), *LT*=heat-labile toxin (Fig. 3), *Stx/St1*=heat-stable toxin I (Fig.4), *blaCTX-M*=Beta-LactamaseCTX-M (Fig. 5), *blaTEM*=Beta-LactamaseTEM (Fig. 6), and *ampC*=AmpC Beta-Lactamase (Fig. 7).


**Oligonucleotide Primer:** The primers used were provided by Metabion (Germany), and they are listed in Table 1.

### Statistical Analysis

All the obtained results were statistically evaluated according to Feldman (2003).

## RESULTS AND DISCUSSION

Contamination of poultry meat with food-borne pathogens is still a significant public health concern. Chicken meat may contain many microorganisms that cause food poisoning. In the last two decades *E. coli*-producing ESBLs have been widely reported in animals and humans posing a severe public health threat (Koju et al. 2022). Moreover, the presence of *E. coli* in chicken meat implies fecal pollution which could be caused by poor slaughtering practices, contaminated surfaces and/or meat handling (Samy et al. 2022).



**Fig. 1:** Agarose gel electrophoresis of PCR products showing amplification of *E. coli phoA* gene products at 720bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative) + six isolates of *E. coli* as isolates No. (2, 4 and 6) chicken meat and isolate No. (1) nuggets, No. (3) frankfurter and isolate No. (5) from luncheon. All isolates were positive *E. coli* gene.

Results relating to the prevalence of *E. coli* in chicken meat and chicken meat products shown in Table 2 appear to be similar to Eman and Sherifa (2012) as 25% in luncheon. Lower results (6.66%) in nuggets were detected by Gaafar et al. (2019). Abdel Tawab et al. (2015) and Al-Jobori et al. (2016) recorded higher results in chicken meat as 40 and 45%, respectively and results from nuggets as 33.3 and 30%, respectively. However, El-Kewaiey (2012) didn't detect *E. coli* in nuggets.

Results regarding aerobic plate count (APC) shown in Table 3 indicated the highest level of contamination in chicken luncheon which could be due to insufficient cooking, post-processing contamination, cross-contamination via slicing machines or cutting knives used in food service facilities, raw materials and spices introduced during manufacturing. In contrast, chicken nuggets were the lowest product, which may be due to the freezing effect that minimizes the total count of *E. coli* (Abdel Fattah 2014). Findings were like the findings reported by Ibrahim et al. (2018) and Khalafalla et al. (2019) in chicken meat and nuggets as  $3.04 \times 10^5 \pm 0.39 \times 10^5$  and  $3 \times 10^3 \pm 5 \times 10^1$ , respectively. In addition, Eman and


Sherifa (2012) recorded  $8.5 \times 10^3$  in chicken luncheon. Higher results have also been reported by Hassanin et al. (2016) and Ibrahim et al. (2018) in chicken meat and nuggets as  $3.78 \times 10^6$  cfu/g and  $4.78 \times 10^5 \pm 0.51 \times 10^5$ , respectively. Furthermore, Nabil et al. (2014) recorded the mean APC in luncheon and frankfurter as  $7.4 \times 10^6 \pm 3.1 \times 10^6$  and  $5.8 \times 10^7 \pm 2.5 \times 10^7$  cfu/g, respectively. Lower results were obtained by Daoud (2012) and Khalafalla et al. (2019) in chicken meat and frankfurter as  $2.1 \times 10^3$  and  $2 \times 10^3 \pm 3 \times 10^3$  cfu/g, respectively.

**Table 2:** Prevalence of *E. coli* in examined chicken meat and chicken meat products samples

Sample Type	Samples examined	<i>E. coli</i> positive samples	
		No.	%
Chicken meat	120	40	33.33
Nuggets	60	8	13.3
Frankfurter	60	10	16.66
Luncheon	60	14	23.33
Total	300	72	24


**Table 3:** Total aerobic plate count (APC) of examined chicken meat and chicken meat samples.

Samples	Minimum	Maximum	Mean±SEM
Chicken meat	$13.3 \times 10^4$	$2.7 \times 10^5$	$2.05 \times 10^5 \pm 0.43 \times 10^4$
Nuggets	$1.2 \times 10^3$	$6.3 \times 10^3$	$3.4 \times 10^3 \pm 0.22 \times 10^3$
Frankfurter	$3.6 \times 10^3$	$5.8 \times 10^3$	$4.7 \times 10^3 \pm 0.11 \times 10^3$
Luncheon	$5.8 \times 10^3$	$8.4 \times 10^3$	$6.5 \times 10^3 \pm 0.15 \times 10^3$



**Fig. 2.** Agarose gel electrophoresis of PCR products showing amplification of *Stx2* gene products at 779bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative) + six isolates of *E. coli*. The isolates (2,4,5 and 6) were positive, while isolates (1,3) were negative.


Results regarding coliform count (Table 4) are similar to  $2.07 \times 10^3$  reported by Hassanin et al. (2016). The source of coliform contamination could be due to chopping, dressing of carcasses, unclean hands, shopping blocks, knives used for handling and cutting as well as contaminated water (Elsaid et al. 2019). Hamada et al. (2008) and Javadi and Safarmashaei (2011) recorded higher values of coliform count in luncheon and chicken meat as  $5.46 \times 10^3$  and  $1.4 \times 10^4$ , respectively. In addition, higher values in luncheon and frankfurter were reported by Bkheet et al. (2007) as  $2.3 \times 10^4$  and  $6.4 \times 10^3$  cfu/g, respectively. Lower counts were reported by Daoud (2012) and Hassan (2015) as  $5.1 \times 10^3$  and  $1.7 \times 10^3$ , respectively. Lower counts in luncheon and frankfurter were reported by Nabil et al. (2014) as  $3.0 \times 10^2 \pm 0.9 \times 10^2$  and  $5.3 \times 10^2 \pm 5.0 \times 10^2$ , respectively.




**Fig. 3.** Agarose gel electrophoresis of PCR products showing amplification of *Lt* gene products at 606bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative) + six isolates of *E. coli*. Isolate No (2) was positive, while other isolates (1,3,4,5, 6) were negative.

**Table 4:** Total coliform count of examined chicken meat and chicken meat samples

Samples	Minimum	Maximum	Mean±SEM
Chicken meat	$1.0 \times 10^3$	$4.2 \times 10^3$	$2.4 \times 10^3 \pm 0.11 \times 10^3$
Nuggets	$4.0 \times 10^2$	$1.2 \times 10^3$	$6.3 \times 10^2 \pm 0.36 \times 10^2$
Frankfurter	$6.0 \times 10^2$	$1.3 \times 10^3$	$8.24 \times 10^2 \pm 0.38 \times 10^2$
Luncheon	$3.0 \times 10^2$	$1.1 \times 10^3$	$8.5 \times 10^2 \pm 0.34 \times 10^2$




**Fig. 4.** Agarose gel electrophoresis of PCR products showing amplification of *Sta/StI* gene products at 229bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative) + six isolates of *E. coli*. The isolates No (2 and 6) were positive while isolates (1,3,4,5) were negative.




**Fig. 5.** Agarose gel electrophoresis of PCR products showing amplification of *blaCTX-M* gene products at 593bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative), all six isolates were positive for *blaCTX-M* gene.

Similar to the results of the present study (Table 5), *E. coli* serogroups O<sub>55</sub>: K<sub>59</sub> and O<sub>119</sub>: K<sub>69</sub> were detected in poultry cuts (thigh and breast) by Saad et al. (2011). In addition, Samah and Erfan (2013) detected O<sub>125</sub>:K<sub>70</sub>,



**Fig. 6.** Agarose gel electrophoresis of PCR products showing amplification of *blaTEM* gene products at 516bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (positive, negative), all six isolates were positive for *blaTEM* gene.



**Fig. 7.** Agarose gel electrophoresis of PCR products showing amplification of *ampC* gene products at 516bp. MWM-molecular weight marker (100-1000bp DNA ladder), + control (Positive, negative), all six isolates were positive for *ampC* gene.

O<sub>55</sub>:K<sub>59</sub>, O<sub>111</sub>:K<sub>58</sub> and O<sub>25</sub>:K<sub>11</sub> serogroups from broiler flocks showed high mortality rates in Sharkia Governorate, Egypt. However, Afify et al. (2020) detected *E. coli* O<sub>157</sub>:H<sub>7</sub>, O<sub>114</sub>:H<sub>21</sub> and O<sub>12</sub> and O<sub>26</sub> from raw chicken meat. In addition, Abdel Tawab et al. (2015) detected different serotypes of *E. coli* O<sub>55</sub>:H<sub>7</sub>, O<sub>78</sub>, O<sub>119</sub>:H<sub>4</sub>, O<sub>125</sub>:H<sub>18</sub> and 10 untyped strains from chicken meat and meat products.

Results regarding to susceptibility of *E. coli* shown in Table 6 are similar to those of Badau (2021) and Badr et al. (2022), who reported a significant antibiotic resistance among *E. coli* strains, that could be due to the unregulated use of antibiotics in chickens, resulting in antimicrobial-resistant *E. coli* strains in chicken meat and its products.

Results of detection of some virulence genes and antibiotic resistance genes shown in Table 7 are in accordance with those of Pehlivanlar et al. (2015), who detected *blaTEM*, *blaCTX-M*, *blaSHV*, *blaOXA* and *blaAmpC* genes from 95.4% of *E. coli* isolates. In addition, Beutin et al. (2007) detected *Stx2* gene in 67.44% of *E. coli* isolates. On the contrary, Kagambèga et al. (2012) detected *Stx2* genes in only 8% of isolates. In addition, Zende et al. (2013) detected *Sta* gene in 18.18% of the *E. coli* isolates and *Lt* gene was not detected in any isolates. Lower detection rate for the *ampC* gene obtained by Rasmussen et al. (2015) was 9%.

**Table 5:** Serogrouping of *E. coli* isolates from examined samples (n=20)

<i>E. coli</i> serogroups	Chicken meat		Nuggets		Frankfurter		Luncheon		Characterization
	No	%	No	%	No	%	No	%	
O <sub>86</sub> :k <sub>61</sub>	1	5	-	-	-	-	1	5	Enteropathogenic
O <sub>55</sub> :k <sub>59</sub>	2	10	-	-	-	-	1	5	Enteropathogenic
O <sub>128</sub> :k <sub>67</sub>	1	5	1	5	-	-	1	5	Enterotoxigenic
O <sub>119</sub> :k <sub>69</sub>	1	5	-	-	-	-	-	-	Enteropathogenic
O <sub>125</sub> :k <sub>70</sub>	2	10	-	-	-	-	1	5	Enterotoxigenic
O <sub>78</sub> :k <sub>80</sub>	1	5	-	-	1	5	-	-	Enteropathogenic
O <sub>25</sub> :k <sub>11</sub>	2	10	-	-	-	-	-	-	Enterotoxigenic
Untyped serogroups	2	10	1	5	1	5	-	-	-
Total	12	60	2	10	2	10	4	20	20 serotypes

**Table 6:** Antimicrobial susceptibility of some isolated *E. coli*.

Antimicrobial Discs	N=10	
	Resistant	Sensitive
Amoxicillin	9	1
Amikacin	4	6
Ampicillin	9	1
Ciprofloxacin	2	8
Cefotaxime	7	3
Gentamycin	4	6
Levofloxacin	1	9
Oxytetracyclin	9	1
Penicillin G	10	0
Vancomycin	8	2

**Table 7:** Detection of some virulence genes and antibiotic resistant genes in some selected *E. coli* isolates

<i>E. coli</i> Sample	phoA	blaTEM	blaCTX-M	ampC	Stx1	Stx2	Lt	Stx2
1	+	+	+	+	-	-	-	-
2	+	+	+	+	+	+	+	+
3	+	+	+	+	-	-	-	-
4	+	+	+	+	-	-	-	+
5	+	+	+	+	-	-	-	+
6	+	+	+	+	+	-	-	+

phoA=Alkaline Phosphatase, blaTEM=Beta-LactamaseTEM, blaCTX-M=Beta-LactamaseCTX-M, ampC=AmpC Beta-Lactamase, Stx1=heat-stable toxin 1, LT=heat-labile toxin, and Stx2=Shiga toxins 2.

## Conclusion

ESBL-producing *E. coli* was detected in most of examined chicken meat and chicken meat products. Antimicrobial susceptibility testing revealed the presence of Multidrug resistant strains. PCR assay was used for confirmation and molecular characterization of some virulence genes and antibiotic resistance genes in *E. coli* isolates.

## Author Contributions

MAH, SAS and EM jointly developed the hypothesis and concept of the study and contributed to the manuscript's chemistry, materials, and technique. For this research and scientific paper, MAH, SAS, GEB and EM are all involved in the experimental procedures and analyses for this study and scientific paper. All authors participated in the experimental analysis and helped rewrite the manuscript. The final manuscript was read and approved by all authors.

## REFERENCES

Abdel El-Tawab AA, Maarouf AA, El-Hofy FI and El-Said AA, 2015. Bacteriological studies on some foodborne bacteria

- isolated from Chicken meat and meat products in Kaliobia Governorate. Benha Veterinary Medicine Journal 29 (2): 47-59. <https://dx.doi.org/10.21608/bvmj.2015.31545>
- Abdel Fattah MG, 2014. Incidence of salmonellae and *Escherichia coli* in ready to eat foods. Thesis, Master of Veterinary Medicine, Benha University, Egypt. <https://dx.doi.org/10.21608/bvmj.2017.31107>
- Adzitey, F., Huda, N. and Shariff, A.H.M, 2021. Phenotypic antimicrobial susceptibility of *E. coli* from raw meats, ready-to-eat meats, and their related samples in one health context. Microorganisms 9(2): 1-11. <https://doi.org/10.3390/microorganisms9020326>
- Afify EA, Shaltout FA and Zakaria IM, 2020. Aerobic plate count of contaminants and molecular characterization of eschereichia coli in raw chicken meat in Ismailia, Egypt. Journal of Veterinary Healthcare 2(2): 23-30. <https://doi.org/10.14302/issn.2575-1212.jvhc-20-3477>
- AL-Jobori KM, Mreh LM and Nader MI, 2016. Detection of *E. coli*, *Salmonella* spp., and *Listeria Monocytogenes* in retail chicken meat and chicken giblets samples using multiplex PCR in Baghdad City. International Journal of Current Microbiology and Applied Sciences 5(9): 290-301. <http://dx.doi.org/10.20546/ijemas.2016.509.033>
- APHA "American Public Health Association" (2001). Compendium of methods for microbiological examination of foods, 4th Ed, 1st, NW Washington DC 2000. 1-3710. USA, pp: 365-366 and 800.
- Archambault M, Petrov P, Hendriksen RS, Asseva G, Bangtrakulnonth A, Hasman H and Aarestrup FM, 2006. Molecular characterization and occurrence of extended-spectrum beta-lactamase resistance genes among *Salmonella enterica* serovar Corvallis from Thailand, Bulgaria, and Denmark. Microbial Drug Resistance 12(3): 192-198. <https://doi.org/10.1089/mdr.2006.12.192>
- Badau EA, 2021. One health perspective on the issue of the antibiotic resistance. Parasite 15(3): 743-749. <https://doi.org/10.14202%2Fvetworld.2022.743-749>
- Badr H, Reda RM, Hagag NM, Kamel E, Elnomrosy SM, Mansour AI, Shahein MA, Ali SF and Ali HR, 2022. Multidrug-resistant and genetic characterization of extended-spectrum beta-lactamase-producing *E. coli* recovered from chickens and humans in Egypt. Animals 12(3): 346. <https://doi.org/10.3390/ani12030346>
- Beutin L, Steinruck H, Krause G, Steege K, Haby S, Hultsch G, and Appel B, 2007. Comparative evaluation of the Rida screen Verotoxin enzyme immunoassay for detection of Shiga-toxin producing strains of *Escherichia coli* (STEC) from food and other sources. Journal of Applied Microbiology 102: 630-639. <https://doi.org/10.1111/j.1365-2672.2006.03139.x>
- Bkheet AA, Rezk MS and Mousa MM, 2007. Study on the microbiological content of local manufactured poultry meat products in El-Bohiera governorate. Assuit Veterinary Medicine Journal 53(112): 1-11. <https://dx.doi.org/10.21608/avmj.2007.175950>
- Cartoni MA, Mattioli S, Twining C, Dal Bosco A, Donoghue

- AM, Arsi K, Angelucci E, Chiattelli D and Castellini C, 2022. Poultry Meat and Eggs as an Alternative Source of n-3 Long-Chain Polyunsaturated Fatty Acids for Human Nutrition. *Nutrients* 14: 1969. <https://doi.org/10.3390/20nu14091969>
- CLSI, 2022. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; M100, 32nd Ed. Clinical and Laboratory Standards Institute, Wayne, PA, USA.
- Colom K, Pérez J, Alonso R, Fernández-Aranguiz A, Lariño E and Cisterna R, 2003. Simple and reliable multiplex PCR assay for detection of *bla*<sub>TEM</sub>, *bla*<sub>SHV</sub> and *bla*<sub>OXA-1</sub> genes in Enterobacteriaceae. *Federation of European Microbiological Societies, Microbiology Letters* 223: 147-151. [https://doi.org/10.1016/S0378-1097%20\(03\)00306-9](https://doi.org/10.1016/S0378-1097%20(03)00306-9)
- Daoud JR, 2012. Microbial quality of frozen chicken meat at grocery stores in Qena city, International Conference and Exhibition on Food Processing & Technology, 2012.
- Dipineto L, Santaniello A, Fontanella M, Lagos K, Fioretti A and Menna LF, 2006. Presence of Shiga toxin-producing *Escherichia coli* O157:H7 in living layer hens. *Letters in Applied Microbiology* 43: 293-295. <https://doi.org/10.1111/j.1472-765x.2006.01954.x>
- El-Kewaiey IA, 2012. Quality assessment of some ready-to-eat and locally produced chicken meat products. *Assiut Veterinary Medicine Journal* 58: 132-151. <https://doi.org/10.21608/avmj.2012.171845>
- El Jalil MH, Khamar M, Maaninou S, Dahha M, Zinedine A and Ameer N, 2020. Antibiotic resistance of *Escherichia coli* strains isolated from broiler meat in Morocco. *International Journal of Veterinary Science* 9: 305-308.
- Elsaid S, Reham A, Amin and Eeilwa-Nesreein Z, 2019. Assessment of bacterial contamination in cattle carcasses at Gharbia Abattoirs. *Benha Veterinary Medical Journal* 36(2): 247-251. <https://doi.org/10.21608/bvmj.2019.13952.1028>
- Eman MS and Sherifa M, 2012. Microbiological loads for some types of cooked chicken meat products at Al-Taif Governorate, KSA. *World Applied Science Journal* 17(5): 593-597.
- Feldman D, Ganon J, Haffman R and Simpson J, 2003. The solution for data analysis and presentation graphics. 2<sup>nd</sup> Ed., Abacus Lancripts, Inc., Berkeley, USA.
- Gaafar R, Hasanine F, Shaltout F, and Zaghoul M, 2019. Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. *Benha Veterinary Medicine Journal* 37(1): 16-21. <https://doi.org/10.21608/bvmj.2019.15642.1064>
- Gonçalves-Tenório A, Silva BN, Rodrigues V, Cadavez V and Gonzales-Barron U, 2018. Prevalence of pathogens in poultry meat: A meta-analysis of European published surveys. *Foods* 7(5): 69. <https://doi.org/10.3390/foods7050069>
- Hamada M, El-Bagory A and Edris A, 2008. Microbial evaluation of poultry meat and its products. Ph.D. Thesis, Dept. Food Hygiene, Faculty of Veterinary Medicine Menoufia University.
- Hassan O, 2015. Microbiological status of poultry carcasses from retailed outlets in Alexandria province. M.V.Sc., Thesis, Faculty of Veterinary Medicine, Alexandria University, Egypt.
- Hassanin M, El-Sabagh RA, Marionet ZN and Mohammed SR, 2016. Bacterial and chemical quality of frozen chicken meat received at governmental hospital modern. *Benha Veterinary Medical Journal* 30(1): 109-117. <https://dx.doi.org/10.21608/bvmj.2016.31355>
- Hu Q, Tu J, Han X, Zhu Y, Ding C and Yu S, 2011. Development of multiplex PCR assay for rapid detection of *Riemerella anatipestifer*, *Escherichia coli*, and *Salmonella enterica* simultaneously from ducks. *Journal of Microbiological Methods* 87: 64-69. <https://doi.org/10.1016/j.mimet.2011.07.007>
- Ibrahim HM, Hassan MA, Amin RA, Shawqy NA and Elkoly RL, 2018. The bacteriological quality of some chicken meat products. *Benha Veterinary Medical Journal* 35(1): 50-57. <http://dx.doi.org/10.21608/bvmj.2018.37967>
- ISO 4832:2006. Microbiology of food and animal feeding stuffs, Horizontal method for enumeration of coliforms, Colony count tech. 3rd Ed. <https://www.iso.org/standard/38282.html>
- Javadi A and Safarmashaei S, 2011. Microbial profile of marketed broiler meat. *Middle East Journal of Sciences Reserches* 9 (5): 652-656.
- Kagambèga A, Martikainen O, Lienemann T, Siitonen A, Traoré AS, Barro N and Haukka K, 2012. Diarrheagenic *Escherichia coli* detected by 16-plex PCR in raw meat and beef intestines sold at local markets in Ouagadougou, Burkina Faso. *International Journal of Food Microbiology* 153: 154-158. <https://doi.org/10.1016/j.ijfoodmicro.2011.10.032>
- Khalafalla FA, Fatma HMA and El-Fouley A, 2019. Microbiological evaluation of chicken meat products. *Journal of Veterinary Medical Research* 26(2): 151-163. <https://dx.doi.org/10.21608/jvmr.2019.66094>
- Koju P, Shrestha R, Shrestha A, Tamrakar S, Rai A, Shrestha P, Madhup SK, Katuwal N and Shrestha A, Shrestha S, Sandip KC, Karki P, Tamang P, Thekkur P and Shrestha SS, 2022. Antimicrobial resistance in *E. coli* isolated from chicken cecum samples and factors contributing to antimicrobial resistance in Nepal. *Tropical Medicine and Infectious Disease* 7: 249. <https://doi.org/10.3390/tropicalmed7090249>
- Lee SI, Kang SG, Kang ML and Yoo HS, 2008. Development of multiplex polymerase chain reaction assays for detecting enterotoxigenic *Escherichia coli* and their application to field isolates from piglets with diarrhea. *Journal of Veterinary Diagnostic Investigation* 20: 492-496. <https://doi.org/10.1177/104063870802000413>
- Moawad AA, Hotzel H, Neubauer H, Ehrlich R, Monecke S, Tomaso H, Hafez HM, Roesler U and El-Adawy H, 2018. Antimicrobial resistance in Enterobacteriaceae from healthy broilers in Egypt: Emergence of colistin-resistant and extended spectrum  $\beta$ -lactamase-producing *Escherichia coli*. *Gut Pathogens* 10: 39. <https://doi.org/10.1186/s13099-018-0266-5>
- Nabil ME, Edris Am, Shaltout FA and Zakaria IM, 2014. Effect of some preservatives on Bacterial Load of some poultry Meat products. *Benha Veterinary medical Journal* 26(1): 94-103.
- Pehlivanlar SO, Ozkan A, Ebru S and Cemil K, 2015. Prevalence of  $\beta$ -Lactamase Producing *Escherichia coli* from Retail Meat in Turkey. *Journal of Food Sciences* 80(9): 2023-2029. <https://doi.org/10.1111/1750-3841.12984>
- Quinn PJ, Markey BK, Leonard FC, Hartigan P, Fanning S and Fitz Patrick ES, 2011. *Veterinary Microbiology and Microbial Disease*. 2nd Ed, Wiley-Blackwell, Chichester.
- Rahman MM, Husna A, Elshabrawy HA, AlamJ, Runa NY, Badruzzaman ATM, Banu NA, Al Mamun M, Paul B, Das S, Rahman M, E-Elahi M, Khairalla AS and Ashou HM, 2020. Isolation and molecular characterization of multidrug-resistant *Escherichia coli* from chicken meat. *Scientific Reports* 10: 21999. <https://doi.org/10.1038/s41598-020-78367-2>
- Rasmussen MM, Opintan JA, Frimodt-Møller N and Styrihshave B, 2015. Beta-lactamase producing *Escherichia coli* isolates in imported and locally produced chicken meat from Ghana. *PLoS ONE* 10(10): e0139706. <https://doi.org/10.1371/journal.pone.0139706>
- Saad MS, Edris AM, Shaltout FA and Shimaa N, 2011. Isolation and identification of salmonellae and *E. coli* from meat and

- poultry cuts by using multiplex PCR. Benha Veterinary Medicine Journal 22 (2): 152-160.
- Samah Eid and Erfan AM, 2013. Characterization of *E. coli* associated with high mortality in poultry flocks. Assiut Veterinary Medicine Journal 59(139): 51-61. <https://dx.doi.org/10.21608/avmj.2013.171926>
- Samy AA, Mansour AS, Khalaf DD and Khairy EA, 2022. Development of multidrug-resistant *Escherichia coli* in some Egyptian veterinary farms, Veterinary World 15(2): 488-495. <https://doi.org/10.14202/vetworld.2022.488-495>
- Srinivasan K, Viswanad B, Asrat L, Kaul CL, Ramarao P. Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. Pharmacological Research 52(4): 313-320. <https://doi.org/10.1016/j.phrs.2005.05.004>
- World Health Organization, 2021. Manual for the Laboratory Identification and Antimicrobial Susceptibility Testing of Bacterial Pathogens of Public Health Importance in the Developing World: *Haemophilus influenzae*, *Neisseria meningitidis*, *Streptococcus pneumoniae*, *Neisseria gonorrhoea*, *Salmonella* serotype Typhi, *Shigella*, and *Vibrio cholera*. Geneva, Switzerland.
- Yang S, Lin CH, Aljuffali IA and Fang JY, 2017. Current pathogenic *Escherichia coli* foodborne outbreak cases and therapy development. Archive of Microbiology 199: 811–825. <https://doi.org/10.1007/s00203-017-1393-y>
- Zende RJ, havhan DM, Suryawanshi PR, Rai AK and Vaidya VM, 2013. PCR detection and serotyping of enterotoxigenic and shigatoxigenic *Escherichia coli* isolates obtained from chicken meat in Mumbai, India. Veterinary World 6(10): 770-773. <http://dx.doi.org/10.14202/vetworld.2013.770-773>