



Microsatellite Variation in Jeju and American horses and their Phylogenetic Relationship

Cheul Jang¹, Baatarsogt Oyungerel² and Gil Jae Cho^{3*}

¹Korean Farrier Association, Gyeonggi 14942, Korea

²School of Animal Science and Biotechnology, Mongolian University of Life Sciences, Ulaanbaatar 17024, Mongolia

³College of Veterinary Medicine, Kyungpook National University, Daegu 41566, Korea

*Corresponding author: chogj@knu.ac.kr

Article History: 23-131

Received: 06-Feb-23

Revised: 22-Feb-23

Accepted: 23-Feb-23

ABSTRACT

The aim of this study was to evaluate the microsatellite (Ms) variation in Jeju and American horses and their phylogenetic relationship, and to establish a lineage-based breeding plan for domestic American miniature horse (AMH) and Quarter horse (QH) in Korea. A total of 83 horse samples (27 AMHs, 32 QHs, and 24 Jeju horses [JHs]) were genotyped with 17 Ms markers. The observed number of alleles per locus ranged from 4 (HTG7) to 12 (ASB17), with an average value of 7.71, 7.41, and 6.11 in AMH, QH, and JH, respectively. Of the 17 Ms markers, HTG4, HTG6, HTG7, HMS1 markers had relatively low PIC values (<0.6) in AMH and HMS1 in QH, and ASB2, HMS7, HTG4, ASB23, HMS1, HTG10 in JH. An average level of genetic variation was as follows: AMH, $H_E = 0.7300$ and $H_O = 0.7317$; QH, $H_E = 0.7505$ and $H_O = 0.7011$; and JH, $H_E = 0.6917$ and $H_O = 0.7011$. QH and JH had a genetic distance of 0.3831, and JH and AMH had a genetic distance of 0.2883. Of the three breeds, QH and JH formed clearly near groups and the AMH formed clearly different groups. This study is judged to be suitable for individual identification and parentage testing for registration of the lineage of AMH and QH. Therefore, the results of this study are an effective tool for genetic research and preservation of these horse breeds.

Key words: American Miniature Horse, Microsatellite Marker, Parentage Verification, Quarter Horse.

INTRODUCTION

The domestication of horses is believed to have been domesticated by humans for meat or milk between 4,000 and 3,000 BC in northern Ukraine or Kazakhstan. Therefore, it is estimated that humans started raising horses from the Neolithic Age to the Bronze Age. The origin of the Quarter horse (QH) is believed to be a cross between a native American horse originating in Spain in the 1600s and an English horse imported to Virginia from west of the Mississippi (Tryon et al. 2007).

The QH also has more than 2 million horses registered by the American Quarter Horse Association, one of the most popular horses in the world (AQHA 2019). QH has a thick neck, long, sloping shoulders, thick chest, short back, and long hips. These features are impeccably straight, coupled with structurally correct legs and feet, and are equally well-developed and well-balanced. These characteristics, based on agility and speed (Avila et al. 2018), demonstrate the best abilities seen in professional rodeos and Wrangler National Finals rodeos

in rodeo time events team roping, barrel racing, tie-down roping and steer wrestling, and are the only western additions to the Fédération Équestre Internationale (FEI) World Equestrian Games. It also has an edge in racing, a sport. In addition to racing, ranching and western events, quarter horses are also used for a variety of events from jumping to dressage. Its best ability is by far the sprint compared to other breeds (Roth et al. 2021). Some horses have been measured at speeds of 88.5 km/h. In addition, the amazing character (gentleness, calmness) and trustworthiness of the QH are used in treatment and rehabilitation riding classes by horse auxiliary activities (Watson et al. 2020). In Korea, it is often used for horseback riding for adults who are experiencing horseback riding for the first time. Since 2011, Gyeongsang buk-do has imported 50 horses to the United States every year for 5 years for the purpose of increasing farm household income and distributing riding horses and supplying them to farm households.

American miniature horses (AMH) are 'miniature' versions of well-balanced horses that are no larger than

Cite This Article as: Jang C, Oyungerel B and Cho GJ, 2023. Microsatellite variation in Jeju and American horses and their phylogenetic relationship. International Journal of Veterinary Science x(x): xxxx. <https://doi.org/10.47278/journal.ijvs/2023.022>

larger dogs and have the morphological characteristics found in most horse breeds (Brooks et al. 2010). AMH come in a variety of colors and types, and their gentle nature makes it easy for children and toddlers who are new to horses to become familiar with horses without hesitation. Despite their small size, they are very versatile, so they excel in various fields such as driving, halter, jumping, and obstacles. With a height of less than 34 inches, it is mainly used for light driving rather than horseback riding, but it is widely used for children's horseback riding in Korea.

Currently, about 27,000 animals are raised in Korea, including 300 QHs, 100 AMHs, and 7,000 Jeju horses (JHs), a Korean native horse (Shin et al. 2020). Each breed has its own characteristics and strengths, and despite the need to be managed and maintained, systematic personal and lineage management has not been carried out yet.

Genetic markers have been used to determine the ancestry of animals for decades. There are various methods for identification and paternity in horses. Currently, parentage verification of horses is performed using a short tandem repeater (STR), and microsatellite (Ms) refers to a class of codominant DNA markers inherited in Mendelian fashion. In horses, Ms DNA marker was first reported by Ellegren et al. (1992) and Marklund et al. (1994). At present, Ms is analyzing using the StockMarks® for Horse Equine 17-plex genotyping kit (Applied Biosystems, Foster City, CA, USA) and the Equine genotype TM Panel 1.1 (Finnzymes Diagnostics, Espoo, Finland) in horse. Based on these results, it is applied to the parentage testing. And to get more accurate results, each laboratory is using TKY additional markers. But, recently, single nucleotide polymorphisms (SNPs) have been trying to introduce a horse's parentage verification. The SNPs are biallelic sequences that are abundantly distributed across most eukaryotic genomes (Salisbury et al. 2003; McKay et al. 2008). SNPs have a very low mutation rate (10^{-8} vs 10^{-3}) compared to STRs, and are short in length, making them ideal for analysis using automated high-throughput techniques, and can be analyzed with multiple techniques, such as denaturing high-performance liquid chromatography (DHPLC) (Liu et al. 1998), invader assays (Mein et al. 2000), and TaqMan® assays (Hui et al. 2008). Recently, interest in SNPs research has been increasing for forensic science (Lee et al. 2005; Lessig et al. 2005) and parentage testing of domestic animals (Werner et al. 2004; Rohrer et al. 2007; Hirota et al. 2010; Lopes et al. 2018; Browning et al. 2018; Pook 2019; Randhawa et al. 2020; Pook et al. 2020; Přibáňová et al. 2020; Long 2021; Hu et al. 2021; Gebrehiwot et al. 2021; Nolte et al. 2022).

The aim of this study was to evaluate the Ms variation in Jeju and American horses and their phylogenetic relationship, and to secure basic data for accurate pedigree management of QH and AMH in Korea.

MATERIALS AND METHODS

Genomic DNA extraction and Microsatellite loci analysis

Genomic DNA from 83 horses (27 AMHs, 32 QHs, and 24 JHs) hair roots was extracted using MFX-2000

(Toyobo, Osaka, Japan) according to the manufacturer's protocols (Tozaki et al. 2001). A total of 17 Ms loci were used for analysis in this study (Table 1). Analysis of this study was performed according to the manufacturer's protocols (Stockmarks®, Applied Biosystems, USA). Each DNA marker was tested using an ABI 3130 xl Genetic Analyzer and GeneMapper Software ver. 4.0 (Applied Biosystems, USA), the size of the alleles (base pairs) for each marker was determined according to the International Society for Animal Genetics (ISAG) standard.

Statistical Analysis

Statistical analyses of 17 Ms loci were performed using R software (Jombart and Ahmed 2011) to analyze basic population-level statistics and genetic diversity. The total number of alleles, polymorphism information content (PIC), observed heterozygosity (H_O), expected heterozygosity (H_E) were calculated. And the gene distance between breeds was calculated based on the standard gene distance value of Nei obtained using the R package (Kamvar et al. 2014).

RESULTS

Analysis of Genetic Diversity

In this study, 17 Ms loci were used to identify genetic diversity and relationships among horse breeds. The observed number of alleles per locus ranged from 4 (HTG7) to 12 (ASB17), with an average value of 7.71, 7.41, and 6.11 in AMH, QH, and JH, respectively. Of the 17 Ms loci, HTG4, HTG6, HTG7, HMS1 loci had relatively low PIC values (<0.6) in AMH and HMS1 in QH, and ASB2, HMS7, HTG4, ASB23, HMS1, HTG10 in JH (Table 2). An average level of genetic variation was as follows: AMH, $H_E=0.7300$ and $H_O=0.7317$; QH, $H_E=0.7505$ and $H_O=0.7011$; and JH, $H_E=0.6917$ and $H_O=0.7011$ (Table 2).

Parentage Testing

The results of parentage testing in 2 foals are shown in Table 3. Foal 1 did not inherit alleles from sire or dam, and excluded by the incompatibility of 11 markers (AHT5, HMS3, HMS6, HMS7, HTG10, ASB17, ASB23, LEX3, CA425, HMS2, and HTG6). But foal 2 was identified as paternity in accordance with Mendel's genetic law.

Genetic Distance and Genetic Relationships of Horse Populations

The results of analyzing the genetic relationships of the 3 horse breeds are as shown in Table 4 and Fig. 1. QH and JH had a genetic distance of 0.3831, and JH and AMH had a genetic distance of 0.2883. Of the three breeds, QH and JH formed clearly near groups and the AMH formed clearly different groups.

DISCUSSION

Ms are beneficial because they have a high rate of polymorphism and are useful for parentage testing in animals (Bowling et al. 1997; Cho et al. 2002; Kim and

Table 1: The primer sequences of the Ms loci used in this study

Locus	Primer sequence (5'→3')	Allele size (bp)	References
AHT4	(FAM)-AACCGCCTGAGCAAGGAAGT, GCTCCCAGAGATTTACCTT	138-170	Binns et al. (1995)
AHT5	(VIC)-ACGGACACATCCCTGCCTGC, GCAGGCTAAGGGGGCTCAGC	128-152	Binns et al. (1995)
ASB2	(VIC)-CCACTAAGTGTCTTCAGAAGG, CACAACTGAGTTCTCTGATAGG	222-266	Breen et al. (1997)
ASB17	(PET)-GAGGGCGGTACCTTTGTACC, ACCAGTCAGGATCTCCACCG	89-131	Breen et al. (1997)
ASB23	(VIC)-GCAAGGATGAAGAGGGCAGC, CTGGTGGGTTAGATGAGAAGTC	176-212	Irvin et al. (1998)
CA425	(PET)-AGCTGCCTCGTTAATTCA, CTCATGTCCGCTTGTCTC	230-250	Eggleston-Stott et al. (1997)
HMS1	(PET)-CATCACTCTTCATGTCTGCTTGG, TTGACATAAATGCTTATCCTATGGC	166-178	Guerin et al. (1994)
HMS2	(NED)-CTTGCAGTCGAATGTGTATTAAT, ACGGTGGCAACTGCCAAGGAAG	218-238	Guerin et al. (1994)
HMS3	(NED)-CCAACCTTTGTACATAACAAGA, CCATCCTCACTTTTCACTTTTGT	150-174	Guerin et al. (1994)
HMS6	(VIC)-GAAGCTGCCAGTATCAACCATTG, CTCCATCTTGTGAAGTGTAACCTCA	153-171	Guerin et al. (1994)
HMS7	(FAM)-CAGGAACTCATGTTGATAACCATC, TGTTGTTGAAACATAACCTTGACTGT	167-189	Guerin et al. (1994)
HTG4	(FAM)-CTATCTCAGTCTTGATTGCAGGAC, CTCCCTCCCTCCCTCTGTTCTC	127-141	Ellegren et al. (1992)
HTG6	(VIC)-GAAGCTGCCAGTATCAACCATTG, CTCCATCTTGTGAAGTGTAACCTCA	153-171	Ellegren et al. (1992)
HTG7	(NED)-CCTGAAGCAGAACATCCCTCCTTG, ATAAAGTGTCTGGGCAGAGCTGCT	118-130	Marklund et al. (1994)
HTG10	(NED)-CAATTCCC GCCC ACCCCCGGCA, TTTTTATCTGATCTGTCACATTT	89-171	Marklund et al. (1994)
LEX3	(PET)-ACACTCTAACCCAGTGTGAGACT, GAAGGAAAAAAGGAGGAAGAC	137-160	Coogle et al. (1996)
VHL20	(FAM)-CAAGTCCTCTTACTTGAAGACTAG, AACTCAGGGAGAATCTTCCCTGAG	89-107	Van Haeringen et al. (1994)

Table 2: Number of alleles, heterozygosity, and PIC of the 17 Ms markers in 3 horse breeds

Marker	No. of allele			H _o			H _e			PIC		
	AMH	QH	JH	AMH	QH	JH	AMH	QH	JH	AMH	QH	JH
AHT4	7	6	9	0.7800	0.5900	0.8800	0.7900	0.6900	0.8300	0.7596	0.6559	0.8116
AHT5	7	7	7	0.7800	0.7800	0.7900	0.7600	0.7500	0.7400	0.7333	0.7131	0.7058
ASB2	6	9	6	0.7400	0.8400	0.6700	0.8100	0.8200	0.6400	0.7779	0.7942	0.5982
HMS3	7	7	6	0.6700	0.9400	0.7500	0.7400	0.8100	0.7800	0.7040	0.7864	0.7469
HMS6	6	7	5	0.8900	0.7200	0.6700	0.8000	0.6800	0.7100	0.7667	0.6282	0.6583
HMS7	7	7	6	0.8100	0.7800	0.5000	0.8100	0.8100	0.5200	0.7833	0.7852	0.4878
HTG4	4	7	6	0.3000	0.5900	0.7100	0.3200	0.6900	0.6500	0.3088	0.6503	0.5899
HTG10	9	10	8	0.8100	0.9100	0.9200	0.8000	0.8400	0.7800	0.7772	0.8202	0.7481
VHL20	9	8	5	0.8900	0.8800	0.9200	0.8300	0.8400	0.7200	0.8107	0.8151	0.6753
ASB17	12	10	8	0.8500	0.7500	0.8300	0.8200	0.8200	0.7800	0.7970	0.7995	0.7560
ASB23	7	6	6	0.8100	0.8100	0.6200	0.8100	0.8300	0.6100	0.7827	0.8023	0.5825
HMS1	7	5	4	0.6300	0.5900	0.5800	0.6400	0.5600	0.5800	0.5944	0.5166	0.5098
LEX3	9	10	7	0.7800	0.7800	0.4600	0.8300	0.8600	0.7800	0.8144	0.8450	0.7472
CA425	9	7	7	0.8900	0.8400	0.8300	0.7700	0.6800	0.7600	0.7439	0.6416	0.7217
HMS2	6	8	6	0.7000	0.7800	0.5800	0.7900	0.7300	0.7100	0.7547	0.7025	0.6636
HTG6	6	7	4	0.4800	0.6600	0.3800	0.4500	0.6800	0.4400	0.4254	0.6327	0.3884
HTG7	4	5	4	0.6300	0.6900	0.8300	0.6400	0.6700	0.7300	0.5911	0.6109	0.6820
Mean	7.71	7.41	6.11	0.7317	0.7605	0.7011	0.7300	0.7505	0.6917	0.7014	0.7176	0.6513

*AMH: American miniature horse, QH: Quarter horse, JH: Jeju horse, H_o: Observed heterozygosity, H_e: Expected heterozygosity, PIC: Polymorphism Information Content.

Choi 2002; Cho and Cho 2004; Li et al. 2004; Sun et al. 2004; Sugiura et al. 2020); they have been used extensively to examine the structure of closely related populations and breed allocation of animals (Kim and Choi 2002; Li et al. 2004; Sun et al. 2004; Yoon et al.

2005; Cho 2005). In the case of cows, pigs, horses, donkeys, and dogs, pedigree control has been routinely carried out by paternity in most countries, which conforms to the standards of the International Society for Animal Genetics (ISAG).




Fig. 1: The phylogenetic tree analysis in 3 horse populations. *AM: American miniature horse, Quarter: Quarter horse, Jeju: Jeju horse.

Table 3: One case of parentage testing by 17 Ms markers in Quarter horse

Horse	AHT4	AHT5	ASB2	HMS3	HMS6	HMS7	HTG4	HTG10	VHL20	ASB17	ASB23	HMS1	LEX3	CA 425	HMS2	HTG6	HTG7
Sire	J/O	K/N	N/O	I/R	P/P	N/O	K/M	L/Q	I/M	N/O	J/K	J/M	S/S	M/N	L/L	G/J	N/O
Dam	H/O	N/N	K/K	I/I	L/O	O/O	K/P	R/S	I/Q	N/Q	L/U	J/M	L/O	J/N	J/O	K/O	K/O
Foal 1	O/O	J/N	K/N	M/R	P/P	J/J	K/M	L/O	M/Q	N/R	S/U	J/M	K/L	N/O	L/L	K/O	K/O
Foal 2	O/O	N/N	K/N	I/I	L/P	O/O	K/K	L/R	I/M	N/N	J/L	J/J	L/S	N/N	L/O	K/J	K/O

*Alphabetical allele codes for all loci are identical to the assignments from the 2000 ISAG Horse Comparison Test.

Table 4: Matrix of Da genetic distances observed among the three horse breeds

Breeds	AMH	JH	QH
AMH	0		
JH	0.2883	0	
QH	0.2105	0.3831	0

*AMH: American miniature horse, QH: Quarter horse, JH: Jeju horse.

To meet the demand of the domestic QH market and secure better QH quality in Korea, it is necessary to select QHs with excellent pedigree to start and strengthen QH breeding. Currently, however, there is insufficient research on the reproduction and inheritance of QH compared to other breeds such as JH and other horses in Korea. Studies have shown that the validity and reliability of markers are estimated based on the polymorphism information content (PIC) values of each marker. Therefore, it is judged that markers with relatively high PIC values (PIC >0.5) will be useful for paternity verification of QH and AMH.

An indicator of diversity in a single gene pool is heterozygosity. In association analysis or connection imbalance analysis, high heterogeneous bonding is more desirable (Kang et al. 2009; Goor et al. 2010). Although heterozygosity increases when multiple groups are mixed, heterozygosity is generally associated with mutations in the population without interbreeding of groups (Oberbauer et al. 2010). In the case of genetic characteristic analysis using a Ms marker, heterogeneity may be determined from the degree of mixing of the target group and the other group. If pure lineage is preserved through strong selection without species mixing, the value of heterozygosity is low. If other varieties are mixed, heterozygosity is high. However, heterozygosity is higher when more individuals are used in the study, so it can be difficult to judge the hybridization of species by heterozygosity alone.

It was found that the average number of alleles was 7.71, 7.41, and 6.11 in AMH, QH, and JH, respectively,

and an average genetic variation levels in AMH ($H_E = 0.7300$ and $H_O = 0.7317$) and QH ($H_E = 0.7505$ and $H_O = 0.7011$), and JH ($H_E = 0.6917$ and $H_O = 0.7011$). These results suggest that the genetic diversity of QH or AMH is more abundant than that of JH. Also, it is considered useful for individual identification or paternity test of these horse breeds, except for markers that show relatively low PIC values (<0.6) for HTG4, HTG6, HTG7, HMS1 markers in AMH and HMS1 marker in QH. As a result of examining the relationships of the AMH and the QH through the genetic distance, it was found to be close to QH and JH. It was confirmed that this supports rumors that QH has been introduced to Jeju Island and raised.

While discussions are currently underway on whether to use SNPs instead of Ms DNA markers for parentage testing and individual identification of horses, the findings suggest that Ms is also effective as a tool for genetic research and preservation of QH and AMH.

Conclusion

This study is judged to be suitable for individual identification and parentage testing for registration of the lineage of AMH and QH. Therefore, the results of this study are an effective tool for genetic research and preservation of these horse breeds.

Author’s Contribution

All research protocols in this study were designed, and conducted by C Jang, who also contributed to data acquisition. GJ Cho and O Baatartsogt contributed to the interpretation of the experimental results and the writing the manuscript.

Funding

This research was supported by the National Research Foundation of Korea, funded by the Ministry of Education, Science and Technology (NRF-2020R111A3067905).

Conflicts of Interest

The authors declare no conflict of interest.

REFERENCES

- AQHA, 2019. Annual Report. American Quarter Horse Association (AQHA), Amarillo, Texas. <https://www.aqha.com/documents/82601/1589238/2019+AQHA+Annual+Report.pdf/f16217ed-1057-37d1-5138-7927af693d62> (accessed on 9 March 2019).
- Avila F, Mickelson JR, Schaefer RJ and McCue M, 2018. Genome-wide signatures of selection reveal genes associated with performance in American Quarter Horse Subpopulations. *Frontiers in Genetics* 9: 249. <https://doi.org/10.3389/fgene.2018.00249>
- Binns MM, Uolmes NG and Holliman AM, 1995. The identification of polymorphic microsatellite loci in the horse and their use in Thoroughbred parentage testing. *British Veterinary Journal* 151: 9-15. [https://doi.org/10.1016/s0007-1935\(05\)80057-0](https://doi.org/10.1016/s0007-1935(05)80057-0)
- Brooks SA, Makvandi-Nejad S, Chu E, Allen JJ, Streeter C, Gu E, McCleery B, Murohy BA, Bellone R and Sutter NB, 2010. Morphological variation in the horse: defining complex traits of body size and shape. *Animal Genetics* 41:159-165. <https://doi.org/10.1111/j.1365-2052.2010.02127.x>
- Bowling AT, Eggleston SML, Byrns G, Clark RS, Dileanis D and Wictum E, 1997. Validation of microsatellite markers for routine horse parentage testing. *Animal Genetics* 28: 247-252. <https://doi.org/10.1111/j.1365-2052.1997.00123.x>
- Breen M, Lindgren G, Binns MM, Norman J, Irvin Z, Bell K, Sandberg K and Ellegren H, 1997. Genetical and physical assignments of equine microsatellite-first integration of anchored markers in horse genome mapping. *Mammalian Genome* 8: 267-273. <https://doi.org/10.1007/s003359900407>
- Browning BL, Zhou Y and Browning SR, 2018. A one-penny imputed genome from next-generation reference panels. *American Journal of Human Genetics* 103: 338-348. <https://doi.org/10.1016/j.ajhg.2018.07.015>
- Cho GJ, 2005. Microsatellite polymorphism and genetic relationship in dog breeds in Korea. *Asian-Australian Journal of Animal Science* 18: 1071-1074.
- Cho GJ and Cho BW, 2004. Microsatellite DNA typing using 16 markers for parentage verification of the Korean native horse. *Asian-Australian Journal of Animal Science* 17: 750-754.
- Cho GJ, Yang YJ, Kang HS and Cho BW, 2002. Genetic diversity and validation of microsatellite markers for Jeju native horse parentage testing. *Korean Journal of Genetics* 24: 359-365.
- Coogle L, Bailet E, Reid R and Russ M, 1996. Equine dinucleotide repeat polymorphisms at loci LEX 002, -003, -004, -005, -007, -008, -009, -010, -011, -013, and -014. *Animal Genetics* 27: 126-127.
- Eggleston-Stott ML, DelValle A, Bautista M, Dileanis S, Wictum E and Bowling AT, 1997. Nine equine dinucleotide repeats at microsatellite loci UCDEQ136, UCDEQ412, UCDEQ425, UCDEQ437, UCDEQ467, UCDEQ487, UCDEQ502 and UCDEQ505, *Animal Genetics* 28: 370-371.
- Ellegren H, Jihansson M, Sandberg K and Andersson L, 1992. Cloning of highly polymorphic microsatellites in the horse. *Animal Genetics* 23: 133-142. <https://doi.org/10.1111/j.1365-2052.1992.tb00032.x>
- Gebrehiwot NZ, Strucken EM, Marshall K, Aliloo H and Gibson JP, 2021. SNP panels for the estimation of dairy breed proportion and parentage assignment in African crossbred dairy cattle. *Genetics Selection Evolution* 53: 21. <https://doi.org/10.1186/s12711-021-00615-4>
- Goor VLH, Panneman H and Haeringen VWA, 2010. Proposal for standardization in forensic equine DNA typing: allele nomenclature for 17 equine-specific STR loci. *Animal Genetics* 41: 122-127. <https://doi.org/10.1111/j.1365-2052.2009.01975.x>
- Guerin G, Bertaud M and Amigues Y, 1994. Characterization of seven new horse microsatellites: HMS1, HMS2, HMS3, HMS5, HMS6, HMS7 and HMS8. *Animal Genetics* 25: 62.
- Hirota KI, Kakoi H, Gawahara H, Hasegawa and Tozaki T, 2010. Construction and validation of parentage testing for thoroughbred horses by 53 single nucleotide polymorphisms *Journal of Veterinary Medical Science* 72: 719-726. <https://doi.org/10.1292/jvms.09-0486>
- Hu LR, Li D, Chu Q, Wang YC, Zhou L, Yu Y, Zhang Y, Zhang SL, Usman T, Xie ZQ, Hou SY, Liu L and Shi WH, 2021. Selection and implementation of single nucleotide polymorphism markers for parentage analysis in crossbred cattle population. *Animal*. Jan; 15(1): 100066. <https://doi.org/10.1016/j.animal.2020.100066>
- Hui L, DelMonte T and Ranade K, 2008. Genotyping using the TaqMan assay. *Current Protocols in Human Genetics*. Chapter 2: Unit 2.
- Irvin Z, Giffard J, Brandon R, Breen M and Bell K, 1998. Equine dinucleotide repeat polymorphisms at loci ASB21, 23, 25 and 37-43. *Animal Genetics* 29: 67.
- Jombart T and Ahmed I, 2011. new tools for the analysis of genome-wide SNP data. *Bioinformatics*. 27: 3070-2071. <https://doi.org/10.1093/bioinformatics/btr521>
- Kamvar Z, Tabima J and Grünwald N, 2014. Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *Peer Journal* 2:e281. <https://doi.org/10.7717/peerj.281>
- Kang BT, Kim KS, Min MS, Chae YJ, Kang JW, Yoon J, Choi J, Seong JK, Park HC, An J, Lee MH, Park HM and Lee H, 2009. Microsatellite loci analysis for the genetic variability and the parentage test of five dog breeds in South Korea. *Genes & Genetics Systems* 84: 245-251. <https://doi.org/10.1266/ggs.84.245>
- Kim KS, Choi CB, 2002. Genetic structure of Korean native pig using microsatellite markers. *Korean Journal of Genetics* 24: 1-7.
- Lee HY, Park MJ, Yoo JE, Chung U, Han GR and Shin KJ, 2005. Selection of twenty-four highly informative SNP markers for human identification and paternity analysis in Koreans. *Forensic Science International* 148: 107-112. <https://doi.org/10.1016/j.forsciint.2004.04.073>
- Lessig R, Zoledziewska M, Fahr K, Edelmann J, Kostrzewa M, Dobosz T and Kleemann WJ, 2005. Y-SNP-genotyping a new approach in forensic analysis. *Forensic Science International* 154: 128-136. <https://doi.org/10.1016/j.forsciint.2004.09.129>
- Li C, Wang Z, Liu B, Yang S, Zhu Z, Fan B, Yu M, Zhao S and Li SK, 2004. Evaluation of the genetic relationship among ten Chinese indigenous pig breeds with twenty-six microsatellite markers. *Asian-Australian Journal of Animal Science* 17: 441-444.
- Liu W, Smith DI, Rechtzigel KJ, Thibodeau SN and James CD, 1998. Denaturing high performance liquid chromatography (DHPLC) used in the detection of germline and somatic mutations. *Nucleic Acids Res.* 26: 1396-1400. <https://doi.org/10.1093/nar/26.6.1396>
- Long J, 2021. Parentage analysis using genome-wide high-density SNP microarray. *Gene* 785: 145605. <https://doi.org/10.1016/j.gene.2021.145605>
- Lopes FB, Wu XL, Li H, Xu J, Perkins T, Genho J, Ferretti R, Tait RG Jr, Bauck S and Rosa GJM, 2018. Improving accuracy of genomic prediction in Brangus cattle by adding

- animals with imputed low-density SNP genotypes. *Journal of Animal Breed Genetics* 135: 14-27. <https://doi.org/10.1111/jbg.12312>
- Marklund S, Ellegren H, Eriksson S, Sandberg K and Andersson L, 1994. Parentage testing and linkage analysis in the horse using a set of highly polymorphic horse microsatellites. *Animal Genetics* 25: 19-23.
- Mckay SD, Schnabel RD, Murdoch BM, Matukumalli LK, Aerts J, Coppeters W, Crews D, Dias NE, Gill CA, Gao C, Mannen H, Wang Z, Tassell VCP, Williams JL, Taylor JF and Moore SS, 2008. An assessment of population structure in eight breeds of cattle using a whole genome SNP panel. *BMC Genetics* 9: 37. <https://doi.org/10.1186/1471-2156-9-37>
- Mein CA, Barratt BJ, Dunn MG, Siegmund T, Smith AN, Esposito L, Nutland S, Stevens HE, Wilson AJ, Phillips MS, Jarvis N, Law S, Arruda DM and Todd JA, 2000. Evaluation of single nucleotide polymorphism typing with invader on PCR amplicons and its automation. *Genome Research* 10: 330-343. <https://doi.org/10.1101/gr.10.3.330>
- Nolte W, Alkhoder H, Wobbe M, Stock KF, Kalm E, Vosgerau S, Krattenmacher N, Thaller G, Tetens J and Kühn C, 2022. Replacement of microsatellite markers by imputed medium-density SNP arrays for parentage control in German warmblood horses. *Journal of Applied Genetics*. 63: 783-792. <https://doi.org/10.1007/s13353-022-00725-9>
- Oberbauer AM, Belanger JM, Grossman DI, Regan KR and Famula TR, 2010. Genome wide linkage scan for loci associated with epilepsy in Belgian shepherd dogs. *BMC Genetics* 4: 11-35. <https://doi.org/10.1186/1471-2156-11-35>
- Pook T, 2019. Methods and software to enhance statistical analysis in large scale problems in breeding and quantitative genetics. Thesis University of Göttingen, Göttingen (Germany).
- Pook T, Mayer M, Geibel J, Weigend S, Cavero D, Schoen CC and Simianer H, 2020. Improving imputation quality in BEAGLE for crop and livestock data. *G3 (Bethesda)* 10: 177-188. <https://doi.org/10.1534/g3.119.400798>
- Přibáňová M, Schroffelová D, Lipovský D, Kučera J, Šteiger V, Hromádková J and Němcová L, 2020. Use of SNPs from Illumina BovineSNP50K BeadChip v3 for imputation of microsatellite alleles for parentage verification and QTL reporting. *Czech Journal of Animal Science* 65: 482-490. <https://doi.org/10.17221/208/2020-CJAS>
- Randhawa IAS, Burns BM, McGowan MR, Porto-Neto LR, Hayes BJ, Ferretti R, Schutt KM and Lyons RE, 2020. Optimized Genetic Testing for Polledness in Multiple Breeds of Cattle. *G3 (Bethesda)* 10: 539-544. <https://doi.org/10.1534/g3.119.400866>
- Rohrer GA, Freking BA and Nonneman D, 2007. Single nucleotide polymorphisms for pig identification and parentage exclusion. *Animal Genetics* 38: 253-258. <https://doi.org/10.1111/j.1365-2052.2007.01593.x>
- Roth IT, Schielke B, Rensing M and Bernau M, 2021. Comparison of American quarter horses competing in western pleasure, hunter under saddle, and reining using linear traits. *Animals* 11: 2861. <https://doi.org/10.3390/ani11102861>
- Salisbury BA, Pungliya M, Choi JY, Jiang R, Sun XJ and Stephens JC, 2003. SNP and Haplotype variation in the human genome. *Mutation Research* 15: 53-61. [https://doi.org/10.1016/s0027-5107\(03\)00014-9](https://doi.org/10.1016/s0027-5107(03)00014-9)
- Shin S.K, Kim SM, Llyod S and Cho GJ, 2020. Prevalence of hoof disorders in horses in South Korea. *The Open Agriculture Journal* 14: 25-29.
- Sun W, Chang H, Ren HZ, Yang ZP, Geng RQ, Lu SX, Du L and Tsunoda K, 2004. Genetic differentiation between sheep and goats based on microsatellite DNA. *Asian-Australian Journal of Animal Science* 17: 583-587.
- Sugiura N, Ochiai K, Yamamoto T, Kato T, Kawamoto Y, Omi T and Hayama SI, 2020. Examining multiple paternity in the raccoon dog (*Nyctereutes procyonoides*) in Japan using microsatellite analysis. *Journal of Veterinary Medical Science* 82: 479-482. <https://doi.org/10.1292/jvms.19-0655>
- Tozaki T, Kakoi H, Mashima S, Hirota KI, Hasegawa T, Ishida N, Choi MNH and Tomita M, 2001. Population study and validation of paternity testing for Thoroughbred horses by 15 microsatellite loci. *Journal of Veterinary Medical Science* 63: 1191-1197. <https://doi.org/10.1292/jvms.63.1191>
- Tryon RC, White SD and Bannasch DL, 2007. Homozygosity mapping approach identifies a missense mutation in equine cyclophilin B (PPIB) associated with HERDA in the American Quarter Horse. *Genomics* 90: 93-102. <https://doi.org/10.1016/j.ygeno.2007.03.009>
- Van Haeringen H, Bowling AT, Scott ML, Lenstra JA and Zwaagstra KA, 1994. A highly polymorphic horse microsatellite locus: VHL20. *Animal Genetics* 25: 207. <https://doi.org/10.1111/j.1365-2052.1994.tb00129.x>
- Watson E, Davis A, Splan R and Porr CAS, 2020. Characterization of horse use in therapeutic horseback riding programs in the United States: A pilot survey. *Journal of Equine Veterinary Science* 92: 103157. <https://doi.org/10.1016/j.jevs.2020.103157>
- Werner FA, Durstewitz G, Habermann FA, Thaller G, Krämer W, Kollers S, Buitkamp J, Georges M, Brem G, Mosner J and Fries R, 2004. Detection and characterization of SNPs useful for identity control and parentage testing in major European dairy breeds. *Animal Genetics* 35: 44-49. <https://doi.org/10.1046/j.1365-2052.2003.01071.x>
- Yoon DH, Kong HS, Oh JD, Lee JH, Cho BW, Kim JD, Jeon JK, Jo CY, Jeon GJ and Lee HK, 2005. Establishment of an individual identification system based on microsatellite polymorphisms in Korean cattle (Hanwoo). *Asian-Australian Journal of Animal Science* 18: 762-766.