



Detection of *Fusobacterium* Species in Female Dromedary Camels using qPCR and MicroSEQ 500

Ahmed Ali^{1,2*}, Derar R Derar^{1,2}, Sahar TM Tolba³ and Sahar A Allam⁴

¹Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Qassim University, Buraidah, Saudi Arabia

²Department of Theriogenology, Faculty of Veterinary Medicine, Assiut University, Assiut, Egypt

³Microbiology Department, Faculty of Science, Ain Shams University, Cairo, Egypt

⁴Infectious Disease Unit, Animal and Poultry Health Department, Animal and Poultry Production Division, Desert Research Center, Cairo, Egypt

*Corresponding author: drahmedali77@gmail.com

Article History: 23-271

Received: 21-Aug-23

Revised: 09-Sep-23

Accepted: 10-Sep-23

ABSTRACT

The goal of this study was to detect *Fusobacterium* spp. in apparently healthy dromedary camels using qPCR and Microseq 500 sequencing. Six pregnant dromedary camels were swabbed for vaginal, rectal, nasal, and ocular discharges. Genomic DNA was extracted. qPCR followed by microseq 500 was used for the detection of *Fusobacterium* spp. The purity of the extracted DNA from various swabs ranged from 1.5 to 2.1. Detection of *Fusobacteria* using qPCR indicated their presence in 91.7% of the collected swabs. *Fusobacterium* spp. was found in all vaginal, ocular, and fecal swabs, but in 66.7% of nasal swabs. The *Fusobacterium* gene was abundant in vaginal swabs. The isolate was identified by 16S rRNA sequence as *Fusobacterium gastrois* CAMSA16 OQ824900. In conclusion, this study sheds light on some naturally occurring microbes in camels that, under specific circumstances, can cause serious diseases. *F. gastrois* may have zoonotic potential.

Key words: Female camels, *Fusobacterium*, qPCR, Microseq 500.

INTRODUCTION

Fusobacterium spp. is a Gram-negative, non-spore-forming, non-motile, anaerobic, rod-shaped bacteria (Han 2015). Many reports considered *Fusobacterium* spp. as a normal flora in the gut and oral cavities, however, the current consensus is that *Fusobacterium* spp. are pathogens (Han 2015; Lee et al. 2022). In women, *Fusobacterium* spp. can produce ovarian abscess (Morrall and Schmidt 2022), premature labor and serious uterine infection (Vidaurrezaga et al. 2020), bacteremia and septic shock (Lee et al. 2022), thrombophlebitis of the internal jugular vein (Kherabi et al. 2020), stillbirth, and neonatal sepsis (Han, 2015). In animals, *Fusobacterium* spp. has been associated with abortion (Agerholm et al. 2007); digital dermatitis (Rosander et al. 2022); metritis (Galvão et al. 2019); salpingitis (Sadeghi et al. 2022) and pyometra (Knudsen et al. 2015). In dromedary camels, *Fusobacterium* has been found in cases with cervicovaginal adhesion (Ghoneim et al. 2021) and in the prepuce of infertile males (Waheed et al. 2022).

Despite serving as the first line of defense against pathogens that ascend in the genital system, vagina contains a variety of bacteria that have the potential to cause a disease (Moreno et al. 2021). Additionally, during pregnancy the uterus is not sterile. Cows and mares may become pregnant notwithstanding the existence of few possibly harmful microbes (Karstrup et al. 2017; van Heule et al. 2023).

An essential protocol in microbiology laboratories is the accurate automated identification and sensitivity testing of chemotherapeutic drug pathogens. VITEK 2 (bioMérieux) and the Phoenix (Becton Dickinson) are two examples of automated identification systems. Unculturable bacteria cannot be detected by automated methods. However, genomic DNA analysis and identification based on the sequencing of the 16S rRNA gene could be an optional strategy. The Microseq 500 was recommended in several reports for sequencing the 16S rRNA gene in samples that are difficult to access by conventional techniques. (Woo et al. 2003; Fontana et al. 2005).

Cite This Article as: Ali A, Derar DR, Tolba STM and Allam SA, 2023. Detection of *Fusobacterium* species in female dromedary camels using qPCR and MicroSEQ 500. International Journal of Veterinary Science x(x): xxxx. <https://doi.org/10.47278/journal.ijvs/2023.088>

This study aimed to use qPCR and Microseq⁵⁰⁰ sequencing to detect *Fusobacterium* spp. in the vagina, feces, and nasal and ocular discharges of apparently healthy pregnant dromedary camels.

MATERIALS AND METHODS

Compliance with Ethical Standards

This study was approved by the Animal Care and Welfare Committee, Deanship of Scientific Research, Qassim University, Kingdom of Saudi Arabia. The summary for the detection of *Fusobacterium* species in female dromedary camels is illustrated (Fig. 1).




Fig. 1: Summary of protocol for detection of *Fusobacterium* species in female dromedary camels using qPCR and MicroSEQ 500.

Sampling

Follow the Green and Correct Red and Similar

Vaginal, rectal, nasal, and ocular swabs were obtained from six pregnant dromedary camels (aged 6-7 years). Swabs were prepared according to the method described by Barati et al. (2022). A sucrose, phosphate, and glutamate (SPG transport) medium were used to collect each swab. SPG medium contained the following ingredients: streptomycin (0.05g/L), L-glutamic acid (0.721g/L), K₂HPO₄ (1.237g/L), KH₂PO₄ (0.512g/L), and sucrose (74.6g/L). A pH of 7.2 to 7.4 was adjusted. Centrifugation at 3000rpm for 15min was used to clear the samples. For DNA extraction, the supernatant was stored at -20°C.

Genomic DNA Isolation from Prepared Swabs

The DNeasy Blood and Tissue Kit (50) from QIAGEN (catalog number 69504) was used to extract DNA from the swabs. The OD A₂₆₀/A₂₈₀ ratio of the purified DNA ranged from 1.7 to 1.9. Thermo Scientific's Nanodrop 2000 spectrophotometer was used to measure the DNA concentration. The 16S rDNA region was amplified using two different methods, qPCR and Microseq500 (Wang et al. 2017).

Real-time Quantitative PCR

The primer and probe set sequences are shown (**Error! Reference source not found.**). The National Center for Biotechnology Information's (NCBI) blasts search for homology with unrelated sequences to validate primer and

probe sequences (<http://www.ncbi.nlm.nih.gov/BLAST>) (Altschul et al. 1997). The optimal concentrations of forward and reverse primers, as well as the probe were 300, 300, and 300nM, respectively. Purified genomic DNA from various swabs was used to test the specificity of the probe and primer sets for their target DNA. Tenfold dilutions of DNA were used in series as standard curves. Amplification was carried out as previously designated by Boutaga et al. (2003). Using Biosystem Step One Plus, the results from unidentified genomic DNA samples were automatically plotted on the standard curves of the target positive control.

MicroSEQ™ 500

The technique was carried out using ThermoFisher Scientific MicroSEQ™ 500 16S rDNA PCR Kit (Catalog Numbers 4348228) and MicroSEQ™ 500 16S rDNA Sequencing Kit (Catalog Numbers 4346480) supplied by Thermo Fisher Scientific (Waltham, USA). As described by (Fontana et al. 2005), amplification of 527bp fragment of the 16S rRNA gene was performed from the 5' end of swab DNA with suspected bacterial identity.

PCR Reactions

MicroAmp™ reaction tubes were used to prepare the samples and controls. The supplied negative control was mixed with 15 µL of PCR master. Positive control DNA is included in the kit. The 9600 thermal cycle from the GeneAmp™ PCR System's 9700-9600 emulation was used. The thermal cycling conditions were set to start at 95°C/10min. The following steps were included in each of the 30 cycles: melting at 95°C/30s, annealing at 60°C/30s, and extension at 72°C/45 seconds. The final extension lasted 10min at 72°C. The final step is 4°C for an infinite time. For separation, PCR products were loaded onto a 2% agarose gel. A standard ladder is used to estimate the PCR product yield. The actual fragment size depends on the bacterial species. PCR products were purified for sequencing.

Statistical Analysis

The data were presented in mean±SD. Analysis of variance (ANOVA) was used for comparison between means with Fisher's protected least significant difference (LSD) as the post-ANOVA test. IBM SPSS Statistics 21 for Windows was used for analysis. The significance level was set at P<0.05.

RESULTS

The extracted DNA purity from the various swabs ranged from 1.5 to 2.1. qPCR detected an increase in the *Fusobacterium* gene in 91.7% (22/24) of the collected swabs. Cycle quantitation (Cq) values ranged from 23 to 40 (Table 2).

Linear regression was calculated automatically by the Biosystem Step One Plus Instrument. The efficiency of the reaction was 88.15% (Fig.). The amplification plot of the target gene after a 10-fold serial dilution with camel swabs was shown (Fig. 3). *Fusobacterium* spp. was found in all vaginal, ocular, and rectal swabs, but only in 66.7% of nasal swabs. *Fusobacterium* gene was found in high concentrations in vaginal swabs compared to fecal, nasal, and ocular swabs (Table 2).

Table 1: Primers and fluorogenic probes for detection of *Fusobacterium* spp.

	Sequence (5 to 3)	Amplicon size (bp)
<i>Fusobacterium</i> spp.	Forward GGATTTATTGGGCGTAAAGC	162
	Reverse GGCATTCCTACAAATATCTACGAA	
	Probe FAM-CTCTACACTTGTAGTTCCG-BHQ	

Table 2: Cycle quantitation (Cq) values for the *Fusobacterium* spp. gene in vaginal (V), ocular (C), nasal (N), and rectal (F) swabs from pregnant female dromedary camels (n=6)

Cq/Swab,s type	1 st	2 nd	3 th	4 th	5 th	6 th	Mean±SD	*F	P-value
Cq/V	23.64	24.74	31.39	28.1	30	35.62	28.91±4.4 ^a	10.75	0.0003
Cq/C	36.97	38.31	36.42	38.45	35.41	37.13	37.11±1.2 ^b		
Cq/N	37.01	40.36	33.62	-ve	-ve	39	37.49±2.9 ^b		
Cq/F	31.2	35	35.39	35.45	36.18	36	34.87±1.9 ^b		

^{a-b} Means in the same column differ significantly.




Fig. 2: The cycle threshold (Cq) values were determined by serially diluting a standard positive control, which contained the target gene, tenfold at concentrations of 2×10^5 , 2×10^4 , 3×10^3 , 2×10^2 , and 20 and 2 copies/ μ L, respectively. The initial number of target genes in the sample is inversely correlated with the Cq values. The Applied Biosystems Step One Plus instrument automatically determined the target gene's copy number. The reaction's efficacy was 88.15%. The linear regression formula was $Y = -4.1946x + 38.741$. Slope = $-4.1946x$; Y-intercept = 38.741; $R^2 = 0.995$.




Fig. 3: Target gene amplification plot after a 10-fold serial dilution in the camel swabs. The fluorescent signal grows at each time point, measured in relative fluorescence units (RFU). Baseline was below (equal 10^1), with PCR cycles accumulating with undetected reporter fluorescent signal. When the fractional PCR cycle number (Ct) was 23 points above the threshold, real amplification began.

The MicroSEQ 500 technique is used for direct DNA sequencing (Fig. 4). Sequence data from camel vaginal swabs was submitted to GenBank as CAMSA16, accession number OQ824900. It is similar to the *Fusobacterium gastrois* strain.




Fig. 4: Band of *Fusobacterium* spp using Microseq500.

DISCUSSION

In this study, a high relative profusion of *Fusobacterium* spp. in the vaginal, rectal, ocular, and nasal secretion of apparently healthy pregnant camels, which is nearly homologous to *F. gastrois*. *F. gastrois* is a novel pathogen with the potential to cause gastric ulceration (De Witte et al. 2018). Earlier studies have exposed existence of *F. gastrois* in the oral and nasal microbiota of pigs and dogs (De Witte et al. 2019), in the stomach of wild boars (Cortez Nunes et al. 2022) and in the stool of humans (De Paepe et al. 2018), indicating that *F. gastrois*, like *F. necrophorum* and *F. nucleatum*, can inhabit a wide variety of mammalian hosts (De Witte et al. 2018).

Fusobacterium gene with high concentration were detected in vaginal swabs compared with fecal, nasal and ocular swabs. The main uterine pathogens were found in vaginal samples from cattle (Jeon et al. 2017). In addition, in cows, a strong association between fecal microbial populations and vaginal-uterine microbial populations was confirmed (Jeon et al. 2017). Galvão et al. (2019) clarified that *Fusobacterium* has the strongest association with metritis.

Despite its role as the frontline to mechanically guard the genital system, the vagina is rich with microbial flora with diverse commensal bacteria and fungus with potential pathogenicity (Moreno et al. 2021). In addition, *Fusobacterium* has been identified in cervicovaginal adhesion cases of dromedaries (Ghoneim et al. 2021).

F. gastrosuis bacteria clearly induced cell death (Thomson et al. 2012). These cell death findings support a role for *F. gastrosuis* in the development of porcine gastric ulceration. However, it is still possible that *F. gastrosuis* is only transient in camels. Future research is needed to determine whether *F. gastrosuis* is a member of the uterine microbiota and/or whether it can cause genital pathologies in dromedaries. The main sources of uterine contamination are thought to be bacteria from the vagina, feces, or environment (Jeon et al. 2017). It would be fascinating to investigate the effect of *F. gastrosuis* on mucosal explants from non-porcine species (De Witte et al. 2018).

Fusobacterium spp. genome analysis revealed the presence of a broad range of virulence-linked genes, which have been related to cell death (Ang et al. 2014). Furthermore, some *Fusobacterium* spp. can adhere to and invade host cells without the assistance of other factors (De Witte et al. 2018). Fusobacteria can produce lactamase, which protects organisms from lactam antibiotics, as well as proteolytic enzymes, which promote regional vein invasion (Karstrup et al. 2017).

Risks associated *Fusobacterium* infection included environmental stress, immunodeficiency, mucus abrasion, unhygienic obstetrical care, or the presence of other supporting microbes. When cows are exposed to a comparable number of pathogens, parturition heat stress disrupts the host immune utility and raises metritis risk (Morrall and Schmidt 2022). Healthy heifers and cows with uterine disease shared the vaginal microbiological league (Moreno et al. 2021). Within 20min of calving, cows have an established uterine microbiome that matches between metritic cows and healthy cows. In cows with metritis, the microbiome deviated two days postpartum in favor of a greater relative abundance of Bacteroidetes and Fusobacteria (Galvão et al. 2019). *Fusobacterium* was isolated from healthy and metritic cow uterine samples (Burfeind et al. 2014).

In conclusion, this study sheds light on a naturally occurring microbe in camels that can cause serious diseases under certain conditions. More research is needed to determine whether *F. gastrosuis* is a member of the uterine microbiota and/or whether it can cause genital pathologies in dromedaries. *F. gastrosuis* may also have zoonotic impending.

Conflict of Interest

The authors declare that they have no conflict of interest.

Acknowledgment

The authors extend their appreciation to the Deputyship for research and Innovation, Ministry of Education, Saudi Arabia, for funding this research work through the project number (QU -IF-1-1-2). The authors also thank to the technical support of Qassim of university.

Funding Information

This work was supported by Ministry of Education, Saudi Arabia through the project number (QU -IF-1-1-2)

Author Contributions

Saher Allam: conceptualization, sampling, methodology; draft writing. Ahmed Ali: writing, reviewing,

editing. Derar R. Derar: reviewing, editing. Sahar T.M. Tolba: methodology and reviewing.

REFERENCES

- Agerholm JS, Boye M and Aalbaek B, 2007. Ovine fetal necrobacillosis. *Journal of Comparative Pathology* 136: 213–221. <https://doi.org/10.1016/j.jcpa.2007.01.012>
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ, 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25: 3389-3402. <https://doi.org/10.1093/nar/25.17.3389>
- Ang MY, Heydari H, Jakubovics NS, Mahmud MI, Dutta A, Wee WY and Choo SW, 2014. Fusobase: an online *Fusobacterium* comparative genomic analysis platform. *Database (Oxford)* 2014: bau082. <https://doi.org/10.1093/database/bau082>
- Barati S, Bakhtiari NM, Shokoohizadeh L, Ghorbanpoor M and Momtaz H, 2022. Genotyping of *Chlamydia abortus* using multiple loci variable number of tandem repeats analysis technique. *BMC Veterinary Research* 18: 54. <https://doi.org/10.1186/s12917-022-03142-6>
- Boutaga K, Van Winkelhoff AJ, Vandenbroucke-Grauls CM and Savelkoul PH, 2003. Comparison of real-time PCR and culture for detection of *Porphyromonas gingivalis* in subgingival plaque samples. *Journal of Clinical Microbiology* 41: 4950-4954. <https://doi.org/10.1128/JCM.41.11.4950-4954.2003>
- Burfeind O, Bruins M, Bos A, Sannmann I, Voigtsberger R and Heuwieser W, 2014. Diagnosis of acute puerperal metritis by electronic nose device analysis of vaginal discharge in dairy cows. *Theriogenology* 82: 64-70. <https://doi.org/10.1016/j.theriogenology.2014.03.001>
- Cortez Nunes F, Letra Mateus T, Taillieu E, Teixeira S, Carolino N, Rema A, De Bruyckere S, Gärtner F, Haesebrouck F and Amorim I, 2022. Molecular detection of *Helicobacter* spp. and *Fusobacterium gastrosuis* in pigs and wild boars and its association with gastric histopathological alterations. *Veterinary Research* 53: 78. <https://doi.org/10.1186/s13567-022-01101-5>
- De Paepe K, Verspreet J, Verbeke K, Raes J, Courtin CM and Van de Wiele T, 2018. Introducing insoluble wheat bran as a gut microbiota niche in an in vitro dynamic gut model stimulates propionate and butyrate production and induces colon region specific shifts in the luminal and mucosal microbial community. *Environmental Microbiology* 20: 3406–3426. <https://doi.org/10.1111/1462-2920.14381>
- De Witte C, Demeyere K, De Bruyckere S, Taminiau B, Daube G, Ducatelle R, Meyer E. and Haesebrouck F, 2019. Characterization of the non-glandular gastric region microbiota in *Helicobacter suis*-infected versus non-infected pigs identifies a potential role for *Fusobacterium gastrosuis* in gastric ulceration. *Veterinary Research* 50: 39. <https://doi.org/10.1186/s13567-019-0656-9>
- De Witte C, Ducatelle R and Haesebrouck F, 2018. The role of infectious agents in the development of porcine gastric ulceration. *Veterinary Journal* 236: 56–61. <https://doi.org/10.1016/j.tvjl.2018.04.015>
- Fontana C, Favaro M, Pelliccioni M, Pistoia ES and Favalli C, 2005. Use of the MicroSeq 500 16S rRNA gene-based sequencing for identification of bacterial isolates that commercial automated systems failed to identify correctly. *Journal of Clinical Microbiology* 43: 615-619. <https://doi.org/10.1128/JCM.43.2.615-619.2005>
- Galvão KN, Bicalho RC and Jeon SJ, 2019. Symposium review: The uterine microbiome associated with the development of uterine disease in dairy cows. *Journal of Dairy Science* 102: 11786–11797. <https://doi.org/10.3168/jds.2019-17106>

- Ghoneim IM, Al-Ahmad JA, Fayez MM, El-Sabagh IM, Humam NA and Al-EknaH MM, 2021. Characterization of microbes associated with cervico-vaginal adhesion in the reproductive system of camels (*Camelus dromedaries*). *Tropical Animal Health and Production* 53: 132. <https://doi.org/10.1007/s11250-020-02538-6>
- Han YW, 2015. *Fusobacterium nucleatum*: a commensal-turned pathogen. *Current Opinion in Microbiology* 23: 141–147. <https://doi.org/10.1016/j.mib.2014.11.013>
- Jeon SJ, Cunha F, Vieira-Neto A, Bicalho RC, Lima S, Bicalho ML and Galvão KN, 2017. Blood as a route of transmission of uterine pathogens from the gut to the uterus in cows. *Microbiome* 5: 1-13. <https://doi.org/10.1186/s40168-017-0328-9>
- Karstrup CC, Agerholm JS, Jensen TK, Swaro LRV, Klitgaard K, Rasmussen EL, Krogh KM and Pedersen HG, 2017. Presence and localization of bacteria in the bovine endometrium postpartum using fluorescence in situ hybridization. *Theriogenology* 92: 167–175. <https://doi.org/10.1016/j.theriogenology.2017.01.026>
- Kherabi Y, Chevrel G, Roux D and Federici L, 2020. Syndrome de Lermierre inversé: à propos d'un cas et revue de la littérature [Gynecological Lermierre's syndrome: A case report and literature review]. *La Revue de Médecine Interne* 41: 493–495. <https://doi.org/10.1016/j.revmed.2020.02.012>
- Knudsen LR, Karstrup CC, Pedersen HG, Agerholm JS, Jensen TK and Klitgaard K 2015. Revisiting bovine pyometra--new insights into the disease using a culture-independent deep sequencing approach. *Veterinary Microbiology* 175: 319–324. <https://doi.org/10.1016/j.vetmic.2014.12.006>
- Lee SJ, Baek YJ, Kim JN, Lee KH, Lee EH, Yeom JS, Choi JY, Ku NS, Ahn JY, Kim JH and Jeong SJ, 2022. Increasing *Fusobacterium* infections with *Fusobacterium varium*, an emerging pathogen. *PloS One* 17: e0266610. <https://doi.org/10.1371/journal.pone.0266610>
- Moreno CG, Luque AT, Galvão KN and Otero MC, 2021. Bacterial communities from vagina of dairy healthy heifers and cows with impaired reproductive performance. *Research in Veterinary Science* 142: 15–23. <https://doi.org/10.1016/j.rvsc.2021.11.007>
- Morrall A and Schmidt U, 2022. *Fusobacterium Necrophorum* Septicemia Secondary to an Ovarian Abscess: A Case Report. *Cureus* 14: e26047. <https://doi.org/10.7759/cureus.26047>
- Rosander A, Albinsson R, König U, Nyman A and Frosth S, 2022. Prevalence of bacterial species associated with ovine footrot and contagious ovine digital dermatitis in Swedish slaughter lambs. *Acta Veterinaria Scandinavica* 64: 6. <https://doi.org/10.1186/s13028-022-00625-2>
- Sadeghi M, Azari M, Kafi M, Nourani H, Ghaemi M, Najafi M and Eshghi D, 2022. Bovine salpingitis: Histopathology, bacteriology, cytology and transcriptomic approaches and its impact on the oocyte competence. *Animal Reproduction Science* 242: 107004. <https://doi.org/10.1016/j.anireprosci.2022.107004>
- Thomson JR and Friendship RM, 2012. The Stomach: Gastric Ulceration. In: Zimmerman JJ, Karriker LA, Ramirez A, Schwartz KJ, Stevenson GW (eds) *Diseases of Swine*, 10th edn. Wiley-Blackwell, Oxford, pp: 208–211.
- Van Heule M, Monteiro HF, Bazzazan A, Scoggin K, Rolston M, El-Sheikh Ali H, Weimer BC, Ball B, Daels P and Dini P, 2023. Characterization of the equine placental microbial population in healthy pregnancies. *Theriogenology* 206: 60–70. <https://doi.org/10.1016/j.theriogenology.2023.04.022>
- Vidaurrezaga MM, McKinley GF and Camins BC, 2020. Postpartum *Fusobacterium gonidiaformans* bacteremia. *Anaerobe* 62: 102168. <https://doi.org/10.1016/j.anaerobe.2020.102168>
- Waheed MM, Ghoneim IM, Fayez MM, El-Bahr SM and Meligy AMA, 2022. Diversity of bacteria and fungi in the prepuce of camels (*Camelus dromedarius*). *Journal of Camel Practice and Research* 29: 77-81. <https://doi.org/10.5958/2277-8934.2022.00011.X>
- Wang J, Gu J, Wang Y, Lin K, Liu S, Lu H, Zhang T and Yu G 2017. 16S rDNA Gene sequencing analysis in functional dyspepsia treated with fecal microbiota transplantation. *Journal of Pediatric Gastroenterology and Nutrition* 64: e80–e82. <https://doi.org/10.1097/MPG.0000000000001476>
- Woo PC, Ng KH, Lau SK, Yip KT, Fung AM, Leung KW, Tam DM, Que TL and Yuen KY 2003. Usefulness of the MicroSeq 500 16S ribosomal DNA-based bacterial identification system for identification of clinically significant bacterial isolates with ambiguous biochemical profiles. *Journal of Clinical Microbiology* 41: 1996–2001. <https://doi.org/10.1128/JCM.41.5.1996-2001.2003>