

Hatching Performance of Kokok Balenggek Chicken (G1): Formation of Superior Local Chicken in West Sumatra

Husmaini, Linda Suhartati, Rusfidra, Fajri Aulia Rachman and Ananda

Department of Animal Science, Faculty of Animal Science Universitas Andalas, West Sumatra, Indonesia

*Corresponding author: lindasuhartati@ansci.unand.ac.id

Article History: 23-352

Received: 08-Nov-23

Revised: 30-Dec-23

Accepted: 10-Jan-24

ABSTRACT

This study aims to determine the hatching performance of Kokok Balenggek chicken (KBC) using different males as the formation of superior local meat-type chickens. The study used an experimental method, randomized block design (RBD), with eight treatments and three groups. The treatment was male KBC (G0) and the group was the hatching period. Chickens are mated using Artificial Insemination (AI) with a sex ratio of 1:5. A total of 594 KBC egg were selected. The variables of this study consisted of fertility rate (FR%), embryonic mortality (EM%), hatchability of fertile egg (HRF%), and day-old chick (DOC) viability. Data were analyzed using Minitab version 19. The results of the study showed that individual differences in males had non-significant ($P>0.05$) effect on FR, EM, HRF and DOC viability. The average FR was (73.38%), EM (30.96%), HRF (69.04%) and DOC viability (90.73%). The macroscopic examination of KBC-G0 semen indicated volume as 0.31 ± 0.11 mL, consistency to be thick creamy with a milky white color and pH as 7.35 ± 0.17 . The microscopic examination of KBC-G0 semen indicated average ++ mass movement and sperm motility, viability and abnormality of spermatozoa to be 71.25 ± 8.35 , 78.34 ± 6.22 and $11.67\pm 2.90\%$, respectively whereas concentration of spermatozoa was $210.46\pm 43.68 \times 10^7$ sperm/mL spermatozoa. In conclusion, the eight males KBC have the same and good hatching performance so can be used as males to produce KBC (G1) in the formation of superior local meat type chickens.

Key words: DOC Viability, Fertility, Hatchability, Mortality, Semen Quality.

INTRODUCTION

Kokok Balenggek chicken (KBC) is one of native chicken from West Sumatra, which originates from Tigo Lurah District, Solok Regency, Indonesia (Masfi and Mafardi 2022). KBC has a multilevel crowing sound and is classified as a singing chicken. The price of KBC is in line with the number and frequency of crowing. However, not all KBC broodstock can produce progeny that have multilevel cocks. Husmaini et al. (2022) stated that 59.69% of breeders in in-situ areas had slaughtered KBC and 52.94% of the KBC has been discarded because these did not have a multilevel crowing. KBC can be classified based on body size and coat color. Based on body size, KBC is divided into three types, namely Yungkilok Gadang chickens (body weight (BW) >2 kg), *Ratih* chickens (BW <2 kg) and Randah Batu chickens (short legs like *Kate* chickens (Rukmana 2003). Yungkilok Gadang type KBC, which does not have multilevel crowing, can be used as a meat-type chicken. The average BW of roosters and hens reported to be 2024.50 ± 291.23 and 1429.34 ± 228.06 g, respectively (Husmaini et al. 2023).

The development of KBC into superior local meat-type chickens can be done by selection. This development program has been successfully carried out on several local chickens, including the Kampung Unggul Balitbangtan chicken (KUB-1). KUB-1 chickens are the result from developing free-range chickens (*Gallus-gallus domesticus*) from Depok, Cianjur and Majalengka (Masito 2021). The development of free-range chickens (*Gallus-gallus domesticus*) produces superior characteristics, including, at the age of 20 weeks, male KUB-1 chickens have a body weight of between 1.60-1.80kg (SNI 2017).

KBC selection can be done based on the quality of male and female. The ability to produce healthy offspring is a characteristic, one of which involves the mating behavior of males and females as well as genetic components (Wolc et al. 2019). The quality of the male can be determined through the hatching performance of eggs produced. The male's quality can determine the egg fertility level (Ridwan and Rusdin, 2008). According to Bandu et al. (2015), egg hatchability can also be influenced by the male's ability to fertilize eggs. Fertilization can occur if the

Cite This Article as: Husmaini, Suhartati L, Rusfidra, Rachman FA and Ananda, 2024. Hatching performance of Kokok Balenggek chicken (G1): Formation of superior local chicken in West Sumatra. International Journal of Veterinary Science x(x): xxx. <https://doi.org/10.47278/journal.ijvs/2024.152>

rooster has good sperm quality and even mortality is closely related to fertility and hatchability (Astomo et al. 2016). A fertile egg has two possibilities: successfully hatching and being mortal. If the egg hatchability is high, the chick mortality rate will be low. Sperm quality has been shown to predict male fertility. By adding external factors like semen collection, processing, and storage, exposure to environmental factors and semen application to a receptive hen, AI further complicates the reproductive process through human intervention (Wolc et al. 2019). So, it is necessary to know the quality of KBC-G0 males through the hatching performance. This difference in male quality can be a reference in developing KBC into superior meat-type chicken to increase reproductive and reproductive performance of this chicken.

MATERIALS AND METHODS

Ethical Approval

Animal experiments were conducted following the Republic of Indonesia Law No. 18 of 2009 (section 66), which addressed animal keeping, raising, killing and proper treatment and care.

Material and Experimental Design

This study used 48 Kokok Balenggek chicken (8 male and 40 female) with male and female with an age of 48 and 36 weeks and body weight 2 and 1.5kg, respectively. Female Kokok Balenggek chicken were in egg production. Chicken were raised intensively using battery cages at the Faculty of Animal Husbandry, Andalas University. Chickens were numbered using wing-bands for males J01 to J08 and for females B01 to B40. KBC-G0 was procured from Solok Regency, West Sumatra. KBC was given the ND vaccine. Chickens were artificially inseminated twice a week with sex ratio of 1:5. A total of 594 KBC egg were selected (storage <7 d). Eggs were selected based on criteria that the eggshell was not cracked, broken, has a normal shape and egg weight ranged from 39-45g. Day-old chicks were maintained for a week to determine its suitability.

The equipment used were three egg hatching machines with a capacity of 200 eggs, 5 watt lamps, thermometers (Gea Medical), hygrometers (HTC-1) with a humidity range, thermostat (STC-1000), mini LED flashlight (Surya SYT L017), 1mL injection syringes (Onemed), 1.5mL microtubes and digital scale (Camry).

Measurement and Methods

This research was conducted using randomized block design (RBD). The treatment was eight male KBC (G0) and the groups were three hatching periods with three egg hatching machines. After the complete end of incubation, the total number of fertility rate, embryonic mortality, hatchability of fertile eggs and DOC viability were recorded according to the groups. Temperature (°C) and humidity (%) were recorded for 21 days.

Hatching results were calculated according to the formula (Alasahan and Copur 2016):

Fertility Rate (FR %) = (number of fertile eggs/numbers of set egg) x 100

Embryonic mortality (%) = (number of embryonic mortality/numbers of fertile egg) x 100

Hatchability of Fertile Egg (HRF %) = (number of hatched chick/number of fertile egg) x 100

DOC Viability (%) = (number of initial DOC observation/total DOC at the end of the observation) x 100.

DOC viability was observed for a week after hatching (Suprijatna et al. 2005). The characteristics of a quality DOC include having fur that looks smooth and full, the DOC's body is not deformed, agile, aggressive, has a high appetite, avoids dry feet and does not have omphalitis (SNI 2017).

Semen collection was by massage method in a microtube twice weekly from each cock. Fresh semen was evaluated macroscopically (volume, color, consistency, pH) and microscopically (mass movement, concentration, motility, abnormalities). The evaluation procedure refers to Arifiantini (2012) which was adapted for poultry semen.

Statistical Analyses

Data were expressed as mean±SD from three hatching machines and analyzed statistically using Minitab version 19. One-way ANOVA was used to analyze the effects of male on hatching performance such as fertility rate, embryonic mortality, hatchability of fertile egg and DOC viability. The percentage was arcsine transformed before analysis. $P < 0.05$ was regarded as statistically significant. Sperm quality was analyzed descriptively. Recording temperature and humidity are displayed in graphs.

RESULTS AND DISCUSSION

Sperm Quality of KBC-G0

The macroscopic examination of KBC-G0 semen indicated volume as 0.31 ± 0.11 mL, consistency to be thick creamy with a milky white color and pH as 7.35 ± 0.17 (Table 1). Ananda et al. (2023) stated that fresh semen of KBC produced a volume ranging from 0.2-0.3mL/ejaculate. Andarusiwo and Yuniati (2021) reported that the semen of the Green Jungle chicken was white to cloudy white. Color and consistency illustrate the concentration of spermatozoa.

The microscopic examination of KBC-G0 semen indicated average ++ mass movement and sperm motility, viability and abnormality of spermatozoa to be 71.25 ± 8.35 , 78.34 ± 6.22 and $11.67 \pm 2.90\%$, respectively whereas concentration of spermatozoa was $210.46 \pm 43.68 \times 10^7$ sperm/mL spermatozoa (Table 2).

Hatchability Performance

The effect of males on hatching performance and DOC viability are showed in Table 3. The male of KBC had no significant ($P > 0.05$) effect on the fertility rate, embryonic mortality, hatching rate of fertile egg and viability of KBC-G1 DOC. Average eight male of KBC percentage of fertility, embryonic mortality, hatching of fertile egg and day-old chicks viability was 73.38, 30.96, 69.04 and 90.73%, respectively (Table 3).

In this study, the KBC-G0 males were selected based on age, body weight, and perfect or flawless appearance. Hatching performance between males does not differ in the present study. This could be due to the same ration fed, namely 524 TA mixed with bran, which can cause the quality of the semen produced and the ability to fertilize to be relatively the same. The level of protein given can affect

Table 1: Macroscopic quality of male KBC-G0 semen

Male	Volume (mL)	Consistency	Color	pH
J01	0.12	Thick creamy	Milky white	7.33
J02	0.38	Thick creamy	Milky white	7.33
J03	0.45	Thick creamy	Milky white	7.17
J04	0.40	Thick creamy	Milky white	7.33
J05	0.32	Thick creamy	Milky white	7.33
J06	0.25	Thick creamy	Milky white	7.67
J07	0.38	Thick creamy	Milky white	7.17
J08	0.20	Thick creamy	Milky white	7.50
Average	0.31±0.11	Thick creamy	Milky white	7.35±0.17

Table 2: The microscopic quality of male KBC-G0 semen

Male	Mass movement	Motility of spermatozoa (%)	Concentration of spermatozoa (10 ⁷ sperm/mL)	Viability (%)	Abnormality of spermatozoa (%)
J01	++	60	162.50	73.17	13.66
J02	+++	80	228.75	79.71	7.25
J03	+++	80	251.25	88.84	10.70
J04	+++	80	278.75	86.12	9.09
J05	++	70	223.75	73.30	9.50
J06	++	70	208.75	75.36	14.98
J07	++	70	171.25	77.97	13.66
J08	++	60	158.70	72.25	14.54
Average	++	71.25±8.35	210.46±43.68	78.34±6.22	11.67±2.90

Table 3: Effect of males on hatching performance and viability of KBC-G1 Day Old Chicks

Treatment	Fertility Rate (%)	Embryonic Mortality (%)	Hatchability of Fertile Egg (%)	Day Old Chicks Viability (%)
J01	67.68±3.00	40.55±6.38	59.45±6.38	91.67±14.43
J02	74.60±9.91	38.33±12.58	61.67±12.58	88.89±19.24
J03	77.18±15.21	33.13±4.47	66.87±4.46	85.00±13.22
J04	84.87±4.50	31.94±6.36	68.06±6.36	78.06±10.55
J05	63.06±3.37	21.67±10.20	78.33±10.20	100.00±0.00
J06	76.35±14.21	18.89±10.09	81.11±10.09	100.00±0.00
J07	68.45±15.19	32.78±7.52	67.22±7.52	88.89±19.25
J08	74.81±7.14	30.36±6.44	69.64±6.44	93.33±11.55
Average	73.38±5.29	30.96±2.69	69.04±2.69	90.73±7.51

the quality of semen in male chickens and in turn, affect fertility and hatchability (Tadondjou et al. 2013). In addition, the eggs collected were selected based on egg weight. Hatching egg quality and hatchability can be affected by egg weight (Ipek and Sozcu 2017).

The average fertility (73.38%) KBC in the present study was lower compared to results 87.70 and 85% reported by Ritonga (2016) and Asri and Surtina (2020), respectively. In this study, mating was carried out using the AI method, whereas in the research conducted by Ritonga (2016) and Asri and Surtina (2020), the natural mating method was used. Sutyono (2001) stated that AI can reduce fertility and hatchability. Implementing AI requires persistence, thoroughness, and caution. If there are errors in the implementation of AI, it can cause damage to the mucosa of the reproductive organs, which can affect the quality of semen and chicken eggs. According to Sadid et al. (2016), percentage of fertility of local chicken eggs using natural mating methods was higher compared to AI. The fertility rate of chicken eggs using the AI method using fresh semen ranges from 62-77% (Long and Kulkarni 2004).

The factors that most influence embryo mortality include egg age, temperature, and humidity in the hatching machine (Ningtyas et al. 2013) and egg handling during hatching (Wicaksono et al. 2013). Napirah and Has (2017) stated that differences in the storage time of hatching eggs can affect embryo mortality. Kostaman et al. (2020) stated that the longer the egg is stored, the quality of the egg

decreases, causing the egg's nutrients used for embryo development to decrease and resulting in failure to hatch. Ningtyas et al. (2013) stated that the temperature of the hatching machine can affect embryo mortality. Eggs that fail to hatch can be caused by the humidity of the hatching machine being too low and the temperature of the hatching machine being too high (Ningtyas et al. 2013). Air humidity functions to maintain fluid in the egg. Inconstant increases and decreases in temperature during hatching can cause embryo death (Ningtyas et al. 2013).

The mortality rate in the present study (30.96%) was lower than that reported (47.2%) by Ritonga (2016). Embryo mortality can be influenced by several factors, including the age of the egg, temperature and humidity of the hatching machine (Ningtyas et al. 2013; Iraqi et al. 2024) egg weight, egg shape index, storage period (Dey et al. 2019) breeder flock age, setter, hatcher type and genotype (Grochowska et al. 2019) and handling of the eggs during hatching (Wicaksono et al. 2013; Ozi et al. 2021). Higher mortality rate (65.83%) has also been reported by Agustira and Yayuk (2017) which could be due to more storage time of hatching eggs as Agustira and Yayuk (2017) stored eggs for nine days while in the present it was only five days. This statement was in accordance with the opinion of Kostaman et al. (2020) that the longer storage of eggs can decrease the quality of the eggs resulting in decreased egg's nutrients necessary for embryo development thus resulting in failure to hatch.




Fig. 1: KBC-G0 hatching temperature (°C). Three groups were three hatching periods.




Fig. 2: Hatch humidity (%) recording. Three groups were three hatching periods.

The average hatchability in the present study ranged from 59.45 ± 6.38 to $81.11 \pm 10.09\%$. Hatchability is also influenced by the length of time the eggs are stored before entering the hatching machine (Ayeni et al. 2020; Melo et al. 2021; Kolokolnikova 2021; Adriaensen et al. 2022; Shirley and Ardener 2023). In this study, the eggs were kept for five days and then started hatching. Short storage can improve hatchability of eggs from young breeders, but not from older breeders (Nasri et al. 2019). The hatchability rate for KBC eggs in this study was 69.04%. The average hatchability in this study was lower than reported (75.9%) by Syamsudin et al. (2016) in three days storage of Sentul Chicken eggs at Warso Unggul Farm. This can happen because hatching eggs stored for three days are fresh than eggs stored for five days. Fresh eggs have small pores to prevent bacteria from entering the egg so that egg quality can be maintained properly (Susanti et al. 2015). Eggs that are contaminated with bacteria cause the nutrients contained in the egg to be damaged so that the nutritional needs for embryo development are not met and this will have an impact on the egg's hatchability (Mahmoud et al. 2022). Bacterial contamination, such as *Enterococcus faecalis* and *Escherichia coli*, can colonize the internal organs of chicken embryos, potentially affecting their viability (Chiang et al. 2022).

In the present study, treatment has no effect on DOC viability (Table 3). The average Day Old Chicks Viability in this study ranged from 78.06 to 100%. DOC survival can be influenced by genetic factors, feeding, breeding management, and disease (Pratiwi et al. 2013). DOC survivability in the study was not 100% due to DOC mortality during one week of rearing. DOC deaths occurred due to physical defects that prevented the DOC from surviving. These physical defects included omphalitis, abnormal leg, and beak shape. Santosa (2016) stated that

omphalitis is the leading cause of increased DOC mortality. Omphalitis in chicks, also known as yolk sac infection, is a common hatchery-born disease that leads to high rates of early chick mortalities (El-Sawah et al. 2016). It is characterized by the infection of the yolk sac, which can result in the deterioration and decomposition of the yolk, leading to nutrient deficiency and a decrease in the transfer of maternal antibodies (Jalob et al. 2016).

Hatching Temperature and Humidity

The range of KBC-G0 hatching temperature data for 21 days was 38 to 38.8°C (Fig. 1) while KBC-G0 hatching humidity for 21 days ranged from 59.7 to 70.72% (Fig. 2).

Temperature is one of the most important factors during incubation and drives embryonic growth and development. Incubation conditions have a substantial impact on embryonic development, hatchability, chick quality and post hatch performance (Hans et al. 2022; Rocha et al. 2022; Fares et al. 2023; Al-Zghoul et al. 2023). Incubation condition need to be adjusted to meet embryonic requirement to obtain optimal chick quality and hatchability (Meijerhof 2009). Egg Shell Temperature lower than 37.8°C from E15 onward appears to be beneficial for optimal embryo development (Maatjens et al. 2016).

Conclusion

In conclusion, the eight males KBC have the same and good hatching performance so can be used as males to produce KBC (G1) in the formation of superior local meat type chickens.

Acknowledgment

The author would like to thank the Directorate of Community Service Technology Research, Directorate General of Higher Education, Research and Technology, Ministry of Education, Culture, Research and Technology for funding this research through the Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) scheme with contract. T/13/UN.16.19/PT.01.03/PDUPT-Pangan/2023 Fiscal Year 2023. Thank you to Andalas University through LPPM for article improvements on Scopis Camp 5 activities.

Author's contribution

All authors contributed equally to the manuscript

Conflict of interest

The authors have declared no conflict of interest

REFERENCES

- Adriaensen H, Parasote V, Castilla I, Bernardet N, Halgrain M, Lecompte F and Rehault-Godbert S, 2022. How egg storage duration prior to incubation impairs egg quality and chicken embryonic development: Contribution of imaging technologies. *Frontiers in Physiology* 13: 1-12. <https://doi.org/10.3389/fphys.2022.902154>
- Agustira R and Yayuk KR, 2017. Storage time and hatching temperature on the hatchability of village chicken eggs. *Jurnal Ilmiah Peternakan* 5(2): 95-101.
- Alasahan S and Copur AG, 2016. Hatching Characteristics and Growth Performance of Eggs with Different Egg Shapes. *Brazilian Journal of Poultry Science* 18(1): 001-008

- <http://dx.doi.org/10.1590/1516-635x1801001-008>
Al-Zghoul MB, Jaradat ZW, Ababneh MM, Okour MZ, Saleh KMM, Alkofahi A and Alboom MH, 2023. Effects of embryonic thermal manipulation on the immune response to post-hatch *Escherichia coli* challenge in broiler chicken. *Veterinary World* 16(5): 918-928. <https://doi.org/10.14202/vetworld.2023.918-928>
- Ananda, Jaswandi, Rusfidra and Gusnidar H, 2023. Sperm Longevity and Motility in Ringer's Lactate Solution with addition of egg yolk among five phenotypes of Kokok Balenggek Chicken. *Bulletin of Animal Science* 47(3): 127-135. <https://doi.org/10.21059/buletinpeternak.v47i3.83647>
- Andaruworo S and Yuniati E, 2021. Teknologi IB pada ayam Hutan Hijau jantan dengan ayam Kampung betina dalam upaya pelestarian plasma nuftah ayam Bekisar. *Jurnal Ahli Muda Indonesia* 2: 57-62. <https://doi.org/10.46510/jami.v2i1.59>
- Arifiantini RI, 2012. Teknik Koleksi dan Evaluasi Semen pada Hewan. Penerbit IPB Press, Bogor.
- Asri A and Surtina D, 2020. Effect of sex ratio in mating Kokok Balenggek Chicken to fertility, Hatchability and hatching weight. *Jurnal Peternakan Mahaputra* 1(1): 16-23. <https://doi.org/10.36665/jpm.v1i1.22>
- Astomo W, Septinova D and Kurtini T, 2016. The effect of sex ratio arabic chicken on fertility, hatchability, and DOC weight. *Jurnal Ilmiah Peternakan Terpadu* 4(1): 6-12. <http://dx.doi.org/10.23960/jipt.v4i1.p%25p>
- Ayeni AO, Agbede JO, Igbasan FA, Onibi GE and Adegbenro M, 2020. Effects of storage periods and positioning during storage on hatchability and weight of the hatched chicks from different egg sizes. *Bulletin of the National Research Centre* 44(101): 1-6. <https://doi.org/10.1186/S42269-020-00362-4>
- Bandu IS, Sutodjo H and Jelantik IGN, 2015. Influences of strain broodcock on hatchability and Weight of doc from laying hens strain cp 909. *Jurnal Nukleus Peternakan* 2(2): 179-185. <https://doi.org/10.35508/nukleus.v2i2.774>
- Chiang CT, Wu YH and Chao CH, 2022. A real-time artificial intelligence recognition system on contaminated eggs for egg selection. *IEEE International Conference on Mechatronics and Automation* pp. 1057-1061. <https://doi.org/10.1109/ICMA54519.2022.9856045>
- Dey S, Samanta R, Pan S, Debnath B and Debbarma A, 2019. Factors associated with embryonic mortality of eggs of vanaraja breeder chicken reared on deep litter system. *International Journal of Current Microbiology and Applied Sciences* 8(4): 525-532. <https://doi.org/10.20546/ijcmas.2019.804.057>
- El-Sawah AA, Dahshan AHM, Nasef SA, El-Nahass ES and Nayel AI, 2016. Characterization of *E. coli* and *Salmonella* spp. isolates associated with omphalitis in baby chicks. *Journal of Veterinary Medical Research* 23 (1): 61-70. <https://doi.org/10.21608/JVMR.2016.43223>
- Fares WA, Ahmed MRM, Rizk RE, Shahein EHA, Boutrous NG and El-Sabroun K, 2023. Influence of non-ventilating intervals during early incubation stage on egg hatching process. *Veterinary World*, 16(7): 1534-1540. <https://doi.org/10.14202/vetworld.2023.1534-1540>
- Grochowska E, Kinal A, Sobek Z, Siatkowski I and Bednarczyk M, 2019. Field study on the factors affecting egg weight loss, early embryonic mortality, hatchability, and chick mortality with the use of classification tree technique. *Journal Poultry Science* 98(9): 3626-3636. <https://doi.org/10.3382/ps/pez180>
- Hans A, Vanille P, Ines C, Nelly B, Maeva H, François L and Sophie RG, 2022. How egg storage duration prior to incubation impairs egg quality and chicken embryonic development: contribution of imaging technologies. *Frontiers in Physiology* 13: 1-12. <https://doi.org/10.3389/fphys.2022.902154>
- Husmaini, Putra RA, Juliyarsi I, Edwin T, Suhartati L, Alianta AA and Harmaini, 2022. Population structure of Kokok Balenggek chicken in in-situ area as indigenous chicken of Indonesia. *Advances in Animal and Veterinary Science* 10(3): 1-6. <http://dx.doi.org/10.17582/journal.aavs/2022/10.5.993.998>
- Husmaini, Suhartati L and Rusfidra, 2023. Qualitative and quantitative characteristics of G0 kokok balenggek chicken: the formation superior local meat-type chicken. *International Journal of Veterinary Science* 12(4): 554-558. <https://doi.org/10.47278/journal.ijvs/2023.001>
- Ipek A and Sozcu A, 2017. Comparison of hatching egg characteristics, embryo development, yolk absorption, hatch window, and hatchability of Pekin Duck eggs of different weights. *Journal Poultry Science* 96(10): 3593-3599. <https://doi.org/10.3382/ps/pex181>
- Iraqi E, Hady AA, Nadia E, Khalil H, El-Saadany A and El-Sabroun K, 2024. Effect of thermal manipulation on embryonic development, hatching process, and chick quality under heat-stress conditions. *Poultry Science* 102(1): 1-10. <https://doi.org/10.1016/j.psj.2023.103257>
- Jalob ZK, Farhan WH, Ibrahim ZY and Jumaa BN, 2016. Bacteriological And Pathological Study Of Omphalitis In Broiler Chicks. *Kufa Journal for Veterinary Medical Sciences*, 6 (2).
- Kolokolnikova TN, 2021. The storage of turkey hatching eggs. *Glavnyj zootehnik* 44-51. <https://doi.org/10.33920/SEL-03-2107-06>
- Kostaman T, Sopiyanas S, Soewandi BDP and Komarudin, 2020. Percentage of fertility and hatchability of cemani and white leghorn chickens based on size of egg weight. *Jurnal Agripet* 20(2): 118-125. <https://doi.org/10.17969/agripet.v20i2.16411>
- Long JA and Kulkarni G, 2004. An effective method for improving the fertility of glycerol-exposed poultry semen. *Journal Poultry Science* 83: 1594-1601. <https://doi.org/10.1093/ps/83.9.1594>
- Maatjens CM, Roovert-Reijrink IAMV, Engel B, Van Der Vol CW, Kemp B and Brand HVD, 2016. Temperature during the last week of incubation. I. Effects on hatching pattern and broiler chicken embryonic organ development. *Poultry Science* 95: 956-965. <http://dx.doi.org/10.3382/ps/pev447>
- Mahmoud BY, Semida DA, Elnesr SS, Elwan H and El-Full EA, 2022. Approaches of egg decontamination for sustainable food safety. *Sustainability* 15(1): 464. <https://doi.org/10.3390/su15010464>
- Masfi R and Mafardi, 2022. The symbolization of the Kukuak Balenggek chicken statue, Solok Regency, West Sumatra Province: Analysis of cultural mythology. *International Journal of Science and Society* 4 (3): 251-264. <https://doi.org/10.54783/ijssoc.v4i3.517>
- Masito M, 2021. The effect of Kampung Unggul Balitbangtan (KUB) chicken innovation characteristics toward the development of farmers perceptions in South Sumatra. *IOP Conference Series: Earth and Environmental Science* 1001: 1-9. <http://doi.org/10.1088/1755-1315/1001/1/012027>
- Meijerhof R, 2009. The influence of incubation on chick quality and broiler performance. Pages 167-176 in Proc. 20th Austral Poultr Sci Symp, 9-11 February 2009. New South Wales, Sydney, Australia.
- Melo EF, Itallo C, Sousa-de, Araújo MV, Triginelli FLS, Castro N, Carneiro B, Leonardo J and Camargos L, 2021. Effect of egg storage duration and egg turning during storage on egg quality and hatching of broiler hatching eggs. *Animal* 15(2): 1-5. <https://doi.org/10.1016/j.animal.2020.100111>
- Napirah A and Has H, 2017. Pengaruh lama penyimpanan terhadap fertilitas dan daya tetas telur ayam kampung persilangan. *Seminar Nasional Riset Kuantitatif Terapan* 1(1): 167-170.

- Nasri HHV, Brand D, Najjar T and Bouzouaia M, 2019. Egg storage and breeder age impact on egg quality and embryo development. *Journal of Animal Physiology and Animal Nutrition* 104(1): 257-268. <https://doi.org/10.1111/jpn.13240>
- Ningtyas MS, Ismoyowati and Sulistyawan IH, 2013. The effect of temperature on hatchability and egg hatching yield duck (*Anas platyrinchos*). *Jurnal Ilmiah Peternakan* 1(1): 347-352.
- Ozl S, Uçar A, Erkuş T, Nicholson AD and Elibol O, 2021. Research Note: Effects of turning and short period of incubation during long-term egg storage on embryonic development and hatchability of eggs from young and old broiler grandparent flocks. *Journal Poultry Science* 100 (4): 1-6. <https://doi.org/10.1016/j.psj.2021.101026>
- Pratiwi RN, Wahyuni HI and Murningsih W, 2013. Pengaruh pemberian vitamin A dan E dalam ransum terhadap daya tunas, daya tetas, bobot tetas dan daya hidup DOC Ayam Kedu Hitam yang dipelihara in situ. *Animal Agriculture Journal* 2 (1): 240-246.
- Ridwan and Rusdin, 2008. Conservation of bureau chicken semen using various dilutions on fertility and the fertility period spermatozoa post artificial insemination. *Jurnal Agroland* 15 (1): 63-67.
- Ritonga ISR, 2016. The effect of egg weight on hatching weight and performance of Balenggek cockerels when providing different protein rations. Thesis. Faculty of Animal Science Andalas University, Padang.
- Rocha ACG, Patrone LGA, Caroline CS, Kaoma SCS, Bicego KC, Szawka RE and Gargaglioni LH, 2023. Metabolic and respiratory chemosensitivity and brain monoaminergic responses to cold exposure in chicks subjected to thermal manipulation during incubation. *Journal of Thermal Biology* 109: 103317. <https://doi.org/10.1016/j.jtherbio.2022.103317>
- Rukmana R, 2003. Intensifikasi dan Pengembangan Ayam Buras. Yogyakarta. Penerbit Kanisius
- Sadid SI, Tanwiriah W and Indrijani H, 2016. Fertilitas, daya tetas, dan bobot tetas Ayam Lokal Jimmy's Farm Cipanas Kabupaten Cianjur Jawa Barat. Student E-journal. Fakultas Peternakan Universitas Padjajaran 5(4): 1-11.
- Santosa PE, 2016. Effectivity of antibiotics against omphalitis cases in broiler. *Jurnal Ilmiah Peternakan Terpadu* 4(4): 319-322. <http://dx.doi.org/10.23960/jipt.v4i4.p%25p>
- Shirley and Ardener, 2023. Chicken egg storage and transport. Burleigh Dodds series in Agricultural Science 141-168. <https://doi.org/10.19103/as.2022.0118.12>
- Standar Nasional Indonesia, 2017. Bibit ayam umur sehari/kuri - bagian 1: KUB-1. Jakarta. Badan Standarisasi Nasional.
- Suprijatna E, Atmomarsono U and Kartasudjana R, 2005. Ilmu Dasar Ternak Unggas. Penebar Swadaya. Jakarta.
- Susanti I, Kurtini T and Septinova D, 2015. The effect of storage duration to fertility, weight loss, hatching eggs, and hatching weight of the Arabic chicken egg. *Jurnal Ilmiah Peternakan Terpadu* 3(4): 185-190. <http://dx.doi.org/10.23960/jipt.v3i4.p%25p>
- Sutiyono, 2001. Teknologi Inseminasi Buatan Pada Ayam Bekisar. Perhutani, Semarang.
- Syamsudin GH, Tanwiriah W and Sujana E, 2016. The fertility, hatchability and DOC weight of Sentul chicken at warso unggul gemilang farm-Bogor. Student E-journal Universitas Padjajaran 5(4): 42-56.
- Tadondjou CD, Ngoula F, Kana JR, Defang HF, Mube HK and Tegula A, 2013. Effect of dietary energy levels on body weight, testicular development and semen quality of local barred chickens of the western highlands of Camerron. *Advance Reproduction Science* 1: 38-43. <http://dx.doi.org/10.4236/arsci.2013.13006>
- Wicaksono D, Kurtini T and Nova K, 2013. Comparison of fertility and, losses, power, and weight hatching native chicken hatching eggs on combination. *Jurnal Ilmiah Peternakan Terpadu* 1(2): 1-8. <http://doi.org/10.23960/jipt.v1i2.159>
- Wolc A, Arango J, Settar P, Fulton JE, O'Sullivan NP and Dekkers JCM, 2019. Genetics of male reproductive performance in White Leghorns. *Journal Poultry Science* 28(7): 2729-2733. <http://dx.doi.org/10.3382/ps/pez077>