

Investigation into *Trypanosoma evansi* Infection in Horses in East Sumba-Indonesia

Ida Ayu Pasti Apsari¹, Ida Bagus Ngurah Swacita², Nyoman Sadra Dharmawan^{3,4}, Ida Bagus Oka Winaya⁵, Umbu Yabu Anngung Praing⁶, Kadek Karang Agustina² and I Wayan Masa Tenaya^{2*}

¹Department of Parasitology; ²Department of Veterinary Public Health; ³Center for Study on Animal Diseases;

⁴Department of Veterinary Clinical Pathology; ⁵Department of Pathology, Faculty of Veterinary Medicine, Udayana University. PB. Sudirman St. Campus, Denpasar City, 80225; Bali Indonesia

⁶Livestock Service Office of East Sumba Regency, Jend Soeharto St. Waingapu City, 87112; East Nusa Tenggara Province-Indonesia

*Corresponding author: wayanmasatenaya@unud.ac.id

Article History: 24-431	Received: 01-Mar-24	Revised: 31-Mar-24	Accepted: 05-Apr-24	Online First: 09-May-24
-------------------------	---------------------	--------------------	---------------------	-------------------------

ABSTRACT

Trypanosoma evansi (*T. evansi*) is a blood parasite, the causative agent of Trypanosomiasis (Surra) in many animal species primarily horses. In Indonesia, surra is a major disease of horses causing a catastrophic outbreak in Sumba Island killed thousands of horses during 2010-2012. Diagnosis of the disease is frequently based on parasitological technique. The aims of this study were to investigate *T. evansi* infection using both serological, parasitological and hematological techniques in horses in East Sumba. A total of 270 blood samples were randomly collected from both sexes of the healthy-looking animals for serum tested with Card Agglutination Test for Trypanosomiasis (CATT), and blood smear for detecting the presence of the parasite and for the leukocyte sub-population respectively. The results showed that 24.81% (67/270) of the samples were positive antibody to *T. evansi* antigen, and only 2.2% (6/270) of the serologically positive samples were also confirmed positive with the parasite. The leukocytes sub-population of the parasite-positive animals consisted of lymphocytes 60-98% ($82.17 \pm 14.43\%$), monocyte 0-4% ($1.5 \pm 1.76\%$), neutrophils 0-40% ($16.17 \pm 14.57\%$), eosinophils 0% and basophils 0-1% ($0.17 \pm 0.41\%$) respectively. It was concluded that the seroprevalence of investigated animals were almost 25%, 2.2% of them suffered with parasitemia, lymphocytosis and neutropenia. This data suggesting a positive correlation between the applied tests, and it was considered as a novel diagnostic confirmation regarding Surra infection in the region.

Key words: *Trypanosoma evansi* seroprevalence, Surra, CATT, Leukocyte sub-population

INTRODUCTION

Trypanosoma evansi (*T. evansi*) the causative agent of Surra is a protozoan of the flagellate class that has a predilection for blood, infecting its host via the fly intermediate host. Generally, horses are most often affected by the disease compared to other animals, causing economic losses mainly due to morbidity and mortality (OIE 2013; Aregawi et al. 2019). Moreover, horses are the most susceptible to *T. evansi* infection, followed by camels and dogs, and buffaloes that are referred to as reservoir animals but in pet animals, dogs are considered to be the most susceptible to *T. evansi* infection (Aregawi et al. 2019; Khan et al. 2022). The incubation period of Surra in horses is 1-4 weeks followed by common but not specific clinical signs such as fluctuating fever, weakness, lethargy, anemia, severe weight loss, petechial hemorrhages sometime with nervous signs which are different from one

host and one place to another associated with its immunosuppressive effects (Desquesnes et al. 2013; Wardhana and Savitri 2018). The grates catastrophic outbreak of Surra in Sumba Island, as the largest source of horse breeding in Indonesia, was reported during 2010-2012 causing more than 2000 animal death, primarily horses followed by cattle (Sawitri and Wardhana 2024). This condition was very detrimental to horse breeders from economic and cultural aspects (Anggung et al. 2019). With the high mortality, as no vaccine for Surra, suggesting no effective anti surra treatment was applied during the outbreak, although prolonged survival of the infected animals may helped by the increased production of anti-inflammatory IL-10 (Nguyen et al. 2023). Therefore, the only appropriate prevention strategy was suggested by understanding and implementing epidemiological approach, including an appropriate control vectors (Kim et al. 2024).

Cite This Article as: Apsari IAP, Swacita IBN, Dharmawan NS, Winaya IBO, Praing UYA, Agustina KK and Tenaya IWM, 2024. Investigation into *Trypanosoma evansi* infection in horses in East Sumba-Indonesia. International Journal of Veterinary Science 13(6): 794-798. <https://doi.org/10.47278/journal.ijvs/2024.171>

Clinical signs of Surra are not always sufficiently specific for confirmation clinical diagnoses, so that laboratory tests are required to support definitive diagnosis (Desquesnes et al. 2022). The diagnosis of trypanosomiasis can be based on finding the parasite itself, although with limited sensitivity and specificity compared to serological assay (Apsari et al. 2024), so that detecting its molecular properties with more sensitivity and high specificity were advised (Desquesnes et al. 2022). The specific potency of every test should be evaluated carefully for effective application based on the different epidemiological conditions. Many researchers have compared several diagnostic techniques such as using PCR, CATT and serological assays (Tehseen et al. 2017; Kim et al. 2024). Detection of *T. evansi* through blood smear is very dependent on the amount of parasitemia that occurs. Serologically, ELISA and CATT both give good results in cattle, buffalo and horses (Laha and Sasmal 2009; Apsari et al. 2024). For field investigations in horses, it was suggested to use serological diagnosis with Card Agglutination Test for *T. evansi* (CATT) because it provides more sensitive, it also showed a higher prevalence for a wide range of many animal species than other traditionally used diagnostic methods (Tehseen et al. 2017; Junco et al. 2024). However, no single test could be used to detect active infections and/or trypanosome species or subspecies and further corrections are required to find specific gaps in diagnostic methods and the sustainable control or elimination of the disease, mainly the use of molecular based diagnostic tests (Desquesnes et al. 2022; Villena et al. 2023). For this reason, in this study, the developed diagnostic method namely CATT and microscopy were coupled with the analysis of leukocyte sub-population, to get a comparative diagnosis.

MATERIALS AND METHODS

Ethical clearance

This research was approved by Ethical Commission of The Faculty of Veterinary Medicine Udayana University with letter No. B/26/UN14.2.9/PT.01.04/2023.

Blood samples collection

A total of 270 blood samples were originated from nine purposively selected districts in East Sumba including Lewa, Katala Hamu Lingu, Waingapu, Pandawai, Kahaungu Eti, Rindi, Pahunga Lodu, Ngadu Ngala, and Wula Waijelu. The samples were collected randomly from both sexes of apparently healthy horses, using K3 EDTA vacutainer tubes (Arkan Medical) for microscopy examination and Plain tube for serological assays respectively. The CATT/ *T. evansi* kit was purchased commercially (<http://www.itg.be/production@itg.be>).

Microanalysis

The microanalysis of blood samples were based on a published-hematological method (Deshpande et al. 2021) with a minor modification. Briefly, a drop of blood sample (3–5 µL) from an animal was placed on one end of a clean glass object for thin blood, air dried for 5 minutes, fixed with absolute methanol and immediately stained with 10% Giemsa. The blood smear preparations were washed under running water and dried and examined under a microscope

with a magnification of 400 – 1000x with immersion oil. Counting of leukocyte-sub populations was done using the automatic hematology analyzer Abaxis HM 5, counting each type of leukocyte per 100 leukocyte cells and analyzed statically using published methods (Conboy and Zajac 2012).

Serological testing

The Surra serological test was carried out using the card agglutination technique for *T. evansi* (CATT/ *T. evansi*) (CATT kit) based on a standard method (Schlenker 1997) with a slight modification. Briefly, serum was diluted in CATT kit buffer with a ratio of 1: 8. Serological testing was carried out by mixing 20 µL of diluted serum samples with one drop (45 µL) of *T. evansi* antigen on the circle on the test card. Each serum sample and antigen were homogenized using a disposable-plastic stirrer. After each sample in the circle was mixed, CATT /T card was placed in a rotating machine at a speed of 70rpm for 5min. The positive reactions were scored based on a publish method (Hagos 2010), indicated by the presence of blue agglutination (sand-like sediment) and scored into five categories namely +3 (very clumpy), +2 (a clear sand-like precipitate), +1 (a sand-like precipitate that can be seen evenly dotted on the circle), +/- (the visible reaction is very faint and almost invisible, - (no agglutination reaction).

RESULTS AND DISCUSSION

Of the total 270 serum samples tested, 24.81% (67/270) were positive using the CATT/ *T. evansi* kit. Only one of the tested samples 1.5% (1/67) showed very strong agglutination (+++), 18% (12/67) with strong agglutination (++), and majority (80.6%; 54/67) of samples classified as weak agglutination (+) (Table 1).

Table 1: Serological test result of 67 samples tested using CATT

Total samples	Agglutination with CATT			
	(+++)	(++)	(+)	(-)
67	1	12	54	-

Note: +++ (very strong agglutination), ++ (Strong agglutination), + (weak agglutination, - (no agglutination)

The microscopy of blood smear examination revealed that only 6 of 270 (2.2%) tested samples were positive parasite of interest (Fig. 1), and these samples were coincidentally positive antibody using CATT ranged from + to ++++. It has been proven that CATT was very sensitive to demonstrate the presence of positive antibody to *T. evansi* in buffalo (Farida et al. 2022). However, CATT results cannot confirm whether the infection is active or not, in some cases the animal had infected by *T. evansi* and the agent was no longer present in the animal, therefore the CATT should be confirmed with PCR for an eradication program purpose (Tejedor-Junco et al. 2023). Likewise, antibodies can last for 2.3-22.6 months after trypanocide treatment, so that serological reactions are not necessarily the result of an active *T. evansi* reaction at that time (Monzon et al. 2003; Elshafie et al. 2013). The positive CATT test results have also been associated with the quantity of parasite circulating in the blood, during parasitemia phase, when trypanosomes exceed 2.5x 10⁶par/ml of blood (Chappuis et al. 2005). Similarly, in this study the serological-positive




Fig. 1: An example of blood smear sample of a horse positive for *Trypanosoma evansi* with strong agglutination status (100x, the parasites are pointed with black arrows).

horses with CAAT were also found positive to contain trypanosomes microscopically, suggesting the horses under study had infected with huge number of detectable trypanosomes. Moreover, the Giemsa staining of thin blood smear method can detect *T. evansi* during parasitemia if 105 trypanosomes/ml blood are circulating in the blood (Reid et al. 2001), although microscopy observation was considered less sensitive than serological assay using CATT and real-time PCR (Nurcahyo et al. 2019; Habeeba et al. 2022). Several risk factors are believed to influence the high or low prevalence of *T. evansi* in horses such as location of sampling, gender, age, horse breed, season when samples were collected, rainy season - summer, when the number of vectors and blood-sucking activity by vectors increases, so that such condition should be considered when doing investigation (Sumbria et al. 2017).

The seroprevalence of *T. evansi* in horses in different regions was found to vary, also depending on the presence of risk factors that influence the region. In this study, which was conducted from March to July (dry season), and a seroprevalence of 24.81% was found. This result was slightly higher than those reported previously (Nurcahyo et al. 2019) when the study was done during wet season (January to March) who found 12.9%. The two different seroprevalence within the same location of study and using the same assay of CATT, may be associated with seasonal reason, as one factor, although no association mainly between sex and age was reported (Benaissa et al. 2020; Sana et al. 2022). Using several methods for diagnosing *T. evansi* in horses, the CATT method was considered the most appropriate choice and sensitive for serological surveys, it can give the highest rate (14.4%) compared to PCR (1.3%) and WOOS test (0.5%), suggesting the PCR and the WOOS test were more specific, so that the seropositive status of animals

should be further confirmed using the PCR method on satellite DNA targets (Tehseen et al. 2017; Kim et al. 2024). However, in this study there was 100% association between the microscopy and the CATT analysis, indicating that the animals under investigated were in acute parasitemia phase with a high quantity of *T. evansi* in the circulating blood, and the CATT predominantly detect IgM during this stage of disease progression (Chandu et al. 2021).

The results of the leukocytes-sub population originated from the serology and microscopy positive horses were lymphocytes (82.17±14.43%), neutrophils (16.17±14.57%), monocytes (1.5±1.76%), basophils and (0.17±0.41%) and eosinophils (0%) respectively. The significance of results with the negative animals is presented in Table 2.

The leukocyte sub-population investigation becomes a reference for diagnosing the cause of the disease for monitoring the course of it. The results of this study (Table 2) showed that only basophils had a significant difference between positive and negative parasitemia. There was a significant decrease in basophils ($P \leq 0.05$) during parasitemia, compared with the non-parasitemia animals. This condition was strongly associated with the function of the cells to protective immunity against parasites infection including helminths, ticks, mites, and protozoan parasites (Eberle and Voehringer 2016). So that the significant reduction of basophils reported here suggesting strong immune responses against the parasites. Interestingly, blood glucose observed from the same animal with parasitemia was significantly lower ($P < 0.01$) than the non-parasitemia horses, but other blood biochemical components were in normal ranges (unpublished data). The reduction of blood glucose during acute infection of Surra as reported elsewhere (Garba and Mayaki 2018), may be associated with the increased utilization of host glucose and depletion by horse body cell during febrile condition.

A relatively high lymphocytosis during parasitemia compared to the normal values was observed, although with no significant difference with $P \geq 0.05$ as demonstrated (Table 3). This phenomenon indicating that the animals with lymphocytosis were in the acute phase of disease progression as reported in experimentally *T. evansi* infected rabbit and sheep (Sivajothi et al. 2015; Olatunde et al. 2021). However, in normal condition, lymphocytosis/leukocytosis in young horses frequently was due to the release of adrenaline as a result of fear, excitement or physical exercise that triggers an increased blood pressure and heart rate (Rossdale et al. 1992). The neutrophil value observed in parasitemia animals was lower compared to the normal value (Table 3), but no significant different ($P \geq 0.05$) illustrated in Table 1.

Table 2: Leukocyte sub-populations (% mean±SD) and significance of positive and negative serological examination.

Leukocyte sub-population	Positive	Negative	Significance	Level of significance
Lymphocyte	82.17±14.43	76.30±11.90	0.392NS	$P \geq 0.05$
Neutrophil	16.17±14.57	15.70±12.96	0.948NS	$P \geq 0.05$
Basophils	0.17±0.41	3.60±3.24	0.023*	$P \leq 0.05$
Monocyte	1.50±1.76	4.40±3.37	0.073NS	$P \geq 0.05$
Eosinophils	0	0	-	-

Note: NS (no significance), *(Significance), Positive and Negative: detected with serological and microscopic analysis.

Table 3: Comparison values (%) leukocyte-subsets during parasitemia with normal condition

Leukocytes subset	Parasitemia	Values ¹	Values ²	Values ³
Lymphocyte	60-98	21-42	18-55	5.0-33.1
Neutrophil	0-40	52-70	36-79	56.19-65.49
Basophils	0-1	0-2	0-3	0
Monocyte	0-4	0-6	0-7	3.97-8.69
Eosinophils	0	0-7	0-16	3.56-6.44

¹Merck's Manual (2010); ²Advia (2010); ³Data of local horses (Radityas 2013).

The reduction of neutrophil in this study, although was not significant was accordance with a condition that when parasitemia by *T. evansi* occurs, neutropenia occurs (Olatunde et al. 2021). This condition is associated with severe acute inflammatory phase and infection as endotoxemia (Mun et al. 2010). Moreover, the neutropenia occurs due to the effect of endotoxemia, 25-35% of horses that show symptoms of colic (as a symptom of surra in horses) experienced endotoxemia, the marginalization of neutrophils in small diameter blood vessels, resulting in a decrease in circulating neutrophils in the peripheral blood (Cuervo 2019). So that the reduction of the neutrophils during the acute phase of *T. evansi* infection may indicate a typical sign of leukocyte subsets.

Conclusion

The investigation into *Trypanosoma evansi* infection in horses in East Sumba, the largest source of horse breeding in Indonesia was done using serological, parasitological and hematological techniques, although additional data regarding significant reduction of blood glucose in horses with parasitemia was not reported here. The seroprevalence of the samples tested was 25 and 2.2% of the seropositive animals suffered with parasitemia, which were also coincidentally experienced lymphocytosis and neutropenia. This data suggesting a positive correlation between the applied tests, and it was considered as a novel diagnostic confirmation regarding Surra infection in the region. One limitation of this study is that the use of molecular technique primarily real-time PCR was not applied to further confirmed diagnostic purposes that may be used to build national policy in controlling Surra in the region.

Acknowledgement

Thank you to the Chancellor, Chair of LPPM Unud and Dean of the Faculty of Veterinary Medicine for the funds (DIPA PNB) of Udayana University for 2023 in accordance with the Research Implementation Assignment Agreement Letter Number B/1.616/UN14.4.A/PT.01.03/2023, dated 02 May 2023 and the facilities provided. Thanks, are also expressed to the research members, students and all parties who have helped carry out this research until it can be published.

Author's contribution

All authors were actively involved with different responsibilities. Ida Ayu Pasti Apsari, Ida Bagus Ngurah Swacita and Nyoman Sadra Dharmawan: preparing research proposal. Ida Bagus Oka Winaya and Umbu Yabu Anngung Praing: Sample collection and conducted laboratory works. Kadek Karang Agustina and I Wayan Masa Tenaya: statistical analyses and write manuscript.

Conflict of Interest: None

REFERENCES

- Advia, 2010. Siemens Health Care Diagnostics Inc. 2010. Operator's Guide. Tarrytown, NY, USA, pp: 9–33.
- Anggung, Yabu PU, Suatha, Ketut I, Sampurna and Putu I, 2019. Keragaman Morfometri Kuda Pacu Sandalwood (*Equus Caballus*) Di Pulau Sumba. *Indonesia Medicus Veterinus* 8(1): 106. <https://doi.org/10.19087/imv.2019.8.1.106>
- Apsari IAP, Suratma NA, Swacita IBN, Soma IG, Sari TK, Putra IPC and Sudipa PH, 2024. Parasitological and serological detection of *Trypanosoma evansi* on Bali Cattle at the Pesangaran Slaughterhouse, Denpasar, Indonesia using Hematological Profile. *Biodiversitas* 25(3): 1057-1062. <https://doi.org/10.13057/biodiv/d250319>
- Aregawi WG, Agga GE, Abdi RD and Büscher P, 2019. Systematic review and meta-analysis on the global distribution, host range, and prevalence of *Trypanosoma evansi*. *Parasites and Vectors* 12(1): 67. <https://doi.org/10.1186/s13071-019-3311-4>
- Benaissa MH, Mimoune N, Bentría Y, Kernif T, Boukheikh A, Youngs CR, Kaidi R, Faye B and Halis Y, 2020. Seroprevalence and risk factors for *trypanosoma evansi*, the causative agent of surra, in the Dromedary Camel (*Camelus Dromedarius*) population in Southeastern Algeria. *Onderstepoort Journal of Veterinary Research* 87(1): 1–9. <https://doi.org/10.4102/ojvr.v87i1.1891>
- Chandu AGS, Sengupta PP, Jacob SS, Suresh KP, Borthakur SK, Patra G and Roy P, 2021. Seroprevalence of *Trypanosoma evansi* in cattle and analysis of associated climatic risk factors in Mizoram, India. *Journal of Parasitic Diseases* 45(1): 244–251. <https://doi.org/10.1007/s12639-020-01301-w>
- Chappuis F, Loutan L, Simarro P, Lejon V and Büscher P, 2005. Options for Field Diagnosis of Human African Trypanosomiasis. *Clinical Microbiology Reviews* 18(1): 133–146. <https://doi.org/10.1128/CMR.18.1.133-146.2005>
- Conboy and Zajac, 2012. *Veterinary Clinical Parasitology*. Edited by Zajac Anne. M, Conboy Gary. A, Little Sussan. E, and Mason Reichard. V, 9ed ed. India: John Wiley and Sons Inc.
- Cuervo, 2019. *Practical Hematology and Biochemistry: How to Interpret Blood Work*. In *Equine Internal Medicine*. www.voorjaarsdagen.eu.
- Deshpande NM, Gite S and Aluvalu R, 2021. A review of microscopic analysis of blood cells for disease detection with AI Perspective. *PeerJ Computer Science* 7: 1–27. <https://doi.org/10.7717/peerj-cs.460>
- Desquesnes M, Gonzatti M, Sazmand A, Thévenon S and Bossard G, 2022. A review on the diagnosis of Animal Trypanosomoses. *Parasites & Vectors* 5(1): 64. <https://doi.org/10.1186/s13071-022-05190-1>
- Desquesnes M, Holzmüller P, Lai DH, Dargantes A, Lun ZR and Jittapapong S, 2013. “*Trypanosoma evansi* and Surra: A Review and perspectives on origin, history, distribution, taxonomy, morphology, hosts, and pathogenic effects. *BioMed Research International* 2013: 194176. <https://doi.org/10.1155/2013/194176>
- Desquesnes Marc, Sazmand A, Gonzatti M, Boulangé A, Bossard G, Thévenon S, Gimonneau G, Truc P and Herder S, 2022. Diagnosis of Animal Trypanosomoses : Proper use of current tools and future prospects. *Parasites & Vectors* 15(1): 235. <https://doi.org/10.1186/s13071-022-05352-1>

- Eberle JU and Voehringer D, 2016. Role of basophils in protective immunity to parasitic infections. *Seminars in Immunopathology* 38(5): 605–613. <https://doi.org/10.1007/s00281-016-0563-3>
- Elshafie EI, Sani RA, Hassan L, Sharma R, Bashir A and Abubakar IA, 2013. Seroprevalence and risk factors of *Trypanosoma evansi* infection in horses in Peninsular Malaysia. *Research in Veterinary Science* 94(2): 285–289. <https://doi.org/10.1016/j.rvsc.2012.09.004>
- Farida A, Amira K and Fahrimal Y, 2022. 11. Diagnostic Surra (*Trypanosoma evansi*) on buffalo slaughtered in banda aceh and aceh besar slaughterhouses based on card agglutination test for *Trypanosoma evansi* (CATT). *Jurnal Medika Veterinaria* 15(1): 75–80. <https://doi.org/10.21157/j.med.vet.v14i2.10164>
- Garba U M and Mayaki AM, 2018. Fluctuations in blood glucose level of donkeys infected with *Trypanosoma evansi*. *ARC Journal of Animal and Veterinary Sciences* 4(1): 14–21. <https://doi.org/10.20431/2455-2518.0401003>
- Habeeba S, Khan RA, Zackaria H, Yammahi S, Mohamed Z, Sobhi W, Abdelkader A, Alhosani MA and Muhairi SA, 2022. Comparison of microscopy, card agglutination test for *Trypanosoma evansi*, and real-time pcr in the diagnosis of trypanosomiasis in Dromedary Camels of the Abu Dhabi Emirate, UAE. *Journal of Veterinary Research (Poland)* 66(1): 125–129. <https://doi.org/10.2478/jvetres-2022-0002>
- Hagos A, 2010. Equine Trypanosomiasis in Ethiopia: Epidemiology, characterization and control. Ph.D Thesis. Katholieke Universiteit Leuven.
- Junco T, Gonz M, Doreste MM, Mart S, Paone M, Cecchi G and Corbera JA, 2024. An Atlas of Surra in Spain: A Tool to Support Epidemiological Investigations and Disease Control, 1–13.
- Khan SZ, Umm-e-Aimen, Rizwan M, Ali A, Khan I, Safiullah, Abidullah, Imdad S, Waseemullah and Khan A, 2022. Epidemiological survey of trypanosomiasis in the dromedary camels raised in Dera Ismail Khan, Khyber Pakhtunkhwa, Pakistan. *Agrobiological Records* 7: 10-17. <https://doi.org/10.47278/journal.abr/2021.010>
- Kim J, Li Z, Radwanska M and Magez S, 2024. Recent progress in the detection of surra , a neglected disease caused by *Trypanosoma evansi* with a one health impact in large parts of the tropic and sub-tropic world. *Journal of Microorganisms* 12(1): 44. <https://doi.org/10.3390/microorganisms12010044>
- Laha R and Sasmal NL, 2009. Short report detection of *Trypanosoma evansi* infection in clinically ill cattle, buffaloes and horses using various diagnostic tests. *Epidemiology and Infection* 137(11): 1583-1585. <https://doi.org/10.1017/S095026880900260x>
- Merck's Manual, 2010. MSD Manual Veterinary Manual. In: Schalm's Veterinary Hematology, 6th Ed. Merck & Co., Inc.
- Monzon CM, Mancebo OA and Russo AM, 2003. Antibody levels by indirect ELISA test in *Trypanosoma evansi* infected horses following treatment with Quinapyrimine Sulphate. *Veterinary Parasitology* 111(1): 59–63. [https://doi.org/10.1016/S0304-4017\(02\)00331-X](https://doi.org/10.1016/S0304-4017(02)00331-X)
- Mun A, Riber C and Trigo P, 2010. Case study hematology and clinical pathology data in chronically starved horses. *Journal of Equine Veterinary Science* 30(10): 581–589. <https://doi.org/10.1016/j.jevs.2010.09.002>
- Nguyen HTT, Magez S and Radwanska M, 2023. From Helping to Regulating – A Transcriptomic profile of Ifng+ IL10+ IL21+ Cd4+ Th1 cells indicates their role in regulating inflammation during experimental trypanosomiasis. *Frontiers in Tropical Diseases* 4: 1127022. <https://doi.org/10.3389/fitd.2023.1127022>
- Nurcahyo W, Marlin RK, Yowi SH and Prastowo J, 2019. The prevalence of horse trypanosomiasis in sumba island, indonesia and its detection using card agglutination tests. *Veterinary World* 12(5): 646–652. <https://doi.org/10.14202/vetworld.2019.646-652>
- OIE, 2013. *Trypanosoma evansi* Infections (Including Surra). OIE Terrestrial Manual 1–4.
- Olatunde OA, Jegede HO and Ameen SA, 2021. Hematological, serum biochemical and histopathological effects of selected herbs and combinations on *Trypanosoma brucei* infected West African Dwarf Sheep. *Asian Journal of Natural Product Biochemistry* 19(1): 10–16. <https://doi.org/10.13057/biofar/f190102>
- Radityas H, 2013. Kajian Hematologi Kuda (Equus Caballus) Lokal Indonesia. Gadjah Mada University Indonesia.
- Reid SA, Husein A and Copeman DB, 2001. Evaluation and improvement of parasitological tests for *Trypanosoma evansi* infection. *Veterinary Parasitology* 102(4): 291–297. [https://doi.org/doi.org/10.1016/S0304-4017\(01\)00539-8](https://doi.org/doi.org/10.1016/S0304-4017(01)00539-8)
- Rossdale PD, Mcgladdery AJ, Ousey JC, Holdstock N, Grainger L and Houghton E, 1992. Increase in plasma progesterone concentrations in the mare after foetal injection with CRH, ACTH or Betamethasone in late gestation. *Equine Veterinary Journal* 24(5): 347–350. <https://doi.org/10.1111/j.1042-3306.1992.tb02853.x>
- Sana K, Monia L, Ameni BS, Haikel H, Imed BS, Walid C, Bouabdella H, Bassem BHM, Hafedh D, Samed B, Makram O, Atef BH, Mohsen B, Taib K, Ammar J, Chedia S and Habib JM, 2022. Serological survey and associated risk factors analysis of trypanosomiasis in camels from Southern Tunisia. *Parasite Epidemiology and Control* 16: e00231. <https://doi.org/10.1016/j.parepi.2021.e00231>
- Sawitri DH and Wardhana AH, 2024. Detection of *Trypanosoma evansi* in healthy horses, cattle and buffaloes in East Sumba : Eight years after outbreak. *IOP Conference Series: Earth and Environmental Science* 192: 012039. <https://doi.org/10.1088/1755-1315/1292/1/012039>
- Schlenker S, 1997. Standard operating procedure. *Textile Chemist and Colorist* 29(7): 283–286. <https://doi.org/10.5055/jem.2005.0060>
- Sivajothi SVC, Rayulu B and Reddy S, 2015. Haematological and biochemical changes in experimental *Trypanosoma evansi* infection in rabbits. *Journal of Parasitic Diseases* 39(2): 216–220. <https://doi.org/10.1007/s12639-013-0321-6>
- Sumbria D, Singla LD, Kumar R, Bal MS and Kaur P, 2017. Comparative seroprevalence and risk factor analysis of *Trypanosoma evansi* infection in equines from different agro-climatic zones of Punjab (India). *OIE Revue Scientifique et Technique* 36(3): 971–980. <https://doi.org/10.20506/rst36.3.2729>
- Tehseen S, Jahan N, Desquesnes M, Shahzad MI and Qamar MF, 2017. Field investigation of *Trypanosoma evansi* and comparative analysis of diagnostic tests in horses from Bahawalpur, Pakistan. *Turkish Journal of Veterinary and Animal Sciences* 41: 288–293. <https://doi.org/10.3906/vet-1504-87>
- Tejedor-Junco MT, Melián Henríquez A, Peláez Puerto P, Ramos MD, González-Martín M, Morales Doreste M, Gimonneau G, Desquesnes M, Martín Martel S and Corbera JA, 2023. Surveillance and control of *Trypanosoma evansi* in the Canary Islands: A descriptive analysis. *Acta Tropica* 246: 106990. <https://doi.org/10.1016/j.actatropica.2023.106990>
- Villena FE, Puicón VH, López AM, Rivera K, Pannebaker D, Valdivia HO and Arévalo H, 2023. Parasitological and molecular detection of *Trypanosoma evansi* in a dog from Tocache, San Martin, Peru. *Veterinary Parasitology: Regional Studies and Reports* 42: 1–6. <https://doi.org/10.1016/j.vprsr.2023.100895>
- Wardhana AH and Savitri DH, 2018. Surra: Trypanosomiasis in livestock is potential as zoonotic disease. *Indonesian Bulletin of Animal and Veterinary Sciences* 28(3): 139. <https://doi.org/10.14334/wartazoa.v28i3.1835>