This is an open-access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)



Review Article

https://doi.org/10.47278/journal.ijvs/2024.192

Comparison Performance and Carcass Characteristics of Native Chicken Breeds under Free-Range and Intensive Rearing System: A Meta-Analysis

Jonathan Anugrah Lase^{1,4}, Rudi Afnan²*, Zakiah Wulandari², Sri Estuningsih³, Tike Sartika⁴, Surya⁴, Sari Yanti Hayanti⁴, Mohammad Miftakhus Sholikin⁴ and Cece Sumantri²

¹Graduate School of Animal Production and Technology, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia ²Department of Animal Production and Technology, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia ³Division of Veterinary Pathology, School of Veterinary and Biomedicine, IPB University, Bogor 16680, Indonesia ⁴Research Center for Animal Husbandry, Research Organization for Agriculture and Food, National Research and Innovation Agency of the Republic of Indonesia (BRIN), Cibinong, Bogor District 16915, Indonesia ***Corresponding author:** <u>rudi afnan@apps.ipb.ac.id</u>

Article History: 24-472	Received: 16-Apr-24	Revised: 09-Jun-24	Accepted: 13-Jun-24	Online First: 20-Jun-24
-------------------------	---------------------	--------------------	---------------------	-------------------------

ABSTRACT

Poultry-rearing systems generally consist of intensive, semi-intensive, or free-range systems. Currently, free-range is a system that guarantees healthy poultry products. This meta-analysis aims to assess the effects of intensive and free-range rearing systems on live weight (LW), feed intake (FI), feed conversion ratio (FCR), body weight gain (BWG), carcass weight (CW), breast meat weight (BM), water holding capacity (WHC), shear force (SF), flavor (Flav), tenderness (Tend), juiciness (Juic), thigh meat (TM), meat water content (MW), protein content (Prot), color (Col), and mortality (Mort) in native chickens. Literature search served as the data source using searches in platforms such as Elsevier, Google Scholar, Springer, Wiley, and Oxford University Press. Twenty-seven (27) articles were identified, covering seven parameters related to the growth performance and carcass characteristics of 30 free-range chickens. The influence of the maintenance system significantly affects the live weight (g) parameter (SMD=-1.21; C.I. 95%=-1.73 to -0.687) (P<0.001) and the carcass weight (g) parameter (SMD=-3.02; C.I. 95%=-4.59 to -1.45) (P<0.001). Regarding breast meat quality parameters, there is a significant influence on part b* (SMD=3.048; C.I. 95%=1.31 to 4.79) (P<0.001). The meta-analysis results concluded that performance and carcass characteristics are better in the intensive system. At the same time, the breast meat quality parameter is better in the free-range system.

Key words: Carcass, Meta-analysis, Native chicken, Performance

INTRODUCTION

Organic and natural foods are currently prevalent food sources and the main target for consumers worldwide (Wang et al. 2009). Poultry products such as meat and eggs are also important food sources. Currently, poultry-rearing systems are receiving much attention due to the demand for food products, including poultry products that are free from chemical residues, antibiotics, and other synthetic materials. There is a growing number of consumers interested in poultry products, particularly those raised in free-range systems (Miao et al. 2005; Wang et al. 2009). Free-range rearing systems are becoming increasingly popular because they affect the welfare of livestock, production efficiency, and the health of poultry products (Chen et al. 2013; Tong et al. 2014). Research by Yamak et al. (2016) shows that free-range rearing positively affects the welfare and quality of wild chicken meat. Consumers are interested in buying and even willing to pay more for free-range raised animal products because they believe the resulting products have better sensory quality and taste (Mcfadden et al. 2017; Zhang et al. 2018).

The demand for animal products produced from freerange rearing is believed to expand globally as people become more aware of consuming animal products raised with animal welfare principles (Tong et al. 2014). The issue of animal products raised with consideration for welfare has been ongoing for a long time. Since the 1990s, countries such as the United States, Australia, and Europe have initiated and promoted animal welfare, focusing on eliminating complete confinement systems for animals, especially those intended for meat and egg consumption

Cite This Article as: Lase JA, Afnan R, Wulandari Z, Estuningsih S, Sartika T, Surya, Hayanti SY, Sholikin MM and Sumantri C, 2024. Comparison performance and carcass characteristics of native chicken breeds under free-range and intensive rearing system: A meta-analysis. International Journal of Veterinary Science 13(6): 970-979. https://doi.org/10.47278/journal.ijvs/2024.192 (Hansen and Østerås 2019; Jin et al. 2019). One form of animal husbandry that upholds animal welfare is the free-range system. Free-range rearing allows animals to exhibit natural behaviors, move freely, receive natural sunlight, and consume natural feed (Pettersson et al. 2018; Fitra et al. 2021). Since this issue has been raised, market segmentation for animal products produced from free-range rearing has increased in these countries (Scrinis et al. 2017).

Poultry products from free-range rearing are generally healthier than those from intensive rearing systems (Rehman et al. 2016; Scrinis et al. 2017). Free-range chickens have better meat quality, composition, and flavor (Yamak et al. 2016). Free-range chickens exhibit better growth performance, meat quality, carcass characteristics, and immune function than cage-raised chickens (Stadig et al. 2016; Tong et al. 2014). Free-range chickens without confinement have better carcass quality and meat flavor (Fanatico et al. 2007; Lewis et al. 1997). On the other hand, poultry raised in confinement systems tend to be more stressed, resulting in physiological responses and behaviors that deviate from their natural tendencies and poorer growth performance (Marin et al. 2001).

Previous studies have discussed the effects of freerange and intensive rearing systems on the growth performance and meat quality of native chickens. However, the results vary, so a meta-analysis study is needed to review and obtain valid conclusions from various studies comprehensively. Therefore, this research aims to compare the growth performance and carcass characteristics of thigh and breast parts of native chickens raised in two rearing systems, namely free-range and intensive-rearing systems, using the meta-analysis method.

MATERIALS AND METHODS

Metadata

The articles were obtained from various electronic databases such as Elsevier, Google Scholar, Springer, Wiley, and Oxford University Press. The keywords used are "chicken," "native," "free-range and "rearing systems." The search results on the database and the selection process, from identification to inclusion (Fig. 1), are used to complete the grammatical qualifications. Articles that have been collected from several web publishers were then coded. Fill in the coding in the form of the author's name, year of publication, publisher, article title, free-range chicken breed, free-range chicken sex, cage size, number of samples, research parameters, and research results (intensive rearing as control and free-range rearing as treatment). Articles that have been coded were then checked by all authors for double checking. The results of the minutes of the study search and selection process can be seen in Table 1. After completing the checking process, it continued to the inclusion and exclusion evaluation selection stage.

Selection criteria

The selection process applies an inclusion and exclusion evaluation of the articles that have been collected. Inclusion criteria were the general characteristics of research subjects from a target study to be researched, such as studies of free-range rearing systems of native chickens; studies compared intensive and free-range rearing systems of native chickens; studies performance measuring growth and carcass characteristics, quality of breast, and thigh meat; the results of statistical analysis were in the form of average values and SD (if the statistical results are in the form of SEM, they were converted to SD for uniformity of data); and published in English with the Digital Object Identifier (DOI) and indexed by Scopus. Meanwhile, exclusion criteria included removing research subjects from a target study that were researched because they did not meet the inclusion criteria of the study, such as studies of free-range rearing systems other than native chickens (such as laying hens and broiler chickens); studies that only examine intensive or semi-intensive or free-range rearing systems for native chickens; and the results of the statistical analysis did not have mean values and SD or SEM.

Meta-Analysis approach

The meta-analysis approach used is a cumulative meta-analysis approach to see changes in the cumulative effect size observed for each parameter. Meta-analysis was conducted using OpenMEE (Makmur et al. 2022; Ariyanti et al. 2023). All analyses on each parameter were carried out with the same method; the author has marked as the Study ID column, and the number of samples, in the control category (intensive) and in the treatment category (free-range), has been converted into count format. Meanwhile, the average value and SD (Standard Deviation) for the control category (intensive) and treatment category (free-range) were changed to a continuous format. Next, the differences in all the data collected in a calculated effect size wizard were measured with data types in the form of means and SD and effect size Hedges' d (Palupi et al. 2012). Hedges effect size method was calculated using Eq.

$$d = \frac{(\bar{x}^T - \bar{x}^C)}{\bar{x}^T} J \tag{1}$$

Where X^T is the average value of the treatment category group (free-range), \overline{X}^C is the average value of the control category group (intensive), S is the combined standard deviation, and J is the correction factor for sample size calculated using Eq. (2) and (3):

$$S = \sqrt{\frac{(N^T - 1)(s^T)^2 + (N^C - 1)(s^C)^2}{(N^T + N^C - 2)}}$$
(2)

$$J = 1 - \frac{3}{(4(N^{C} + N^{T} - 2) - 1)}$$
(3)

Where N^T is the number of samples from the treatment category group (free-range), N^C is the number of samples from the control category group (intensive), S^T is the standard deviation of the treatment category group (freerange), and S^C is the standard deviation of the control category group (intensive). Then, the inverse variance of fixed effect method for estimating effect size C.I. 95% use Eq. (4). This range is the confidence interval (CI) which is estimated on the basis of a desired confidence level:

$$CI = \theta \pm Z . SE \tag{4}$$

Where CI is the Confidence Interval, θ is the value of the combined effect estimate from the meta-analysis (the weighted average of the effects of the analyzed studies),



Fig. 1: Flow of the article search and selection process

Z is the z score corresponding to the 95% confidence level (usually around 1.96 to 95% confidence) and SE is the Standard Error of the combined effect estimate, where this SE measures the extent to which the average effect estimate can vary.

In the OpenMEE software, control group and treatment group categories are filled in based on the mean (x), SD (sd), and sample size (n) codes. Subgroup Meta-Analysis is seen based on the variables sex and cage area. The random-effects method uses DL (DerSimonian-Liard) and a Confidence level of 95%. Standard Meta-Analysis analyzes df and effect size C.I. 95% (SMD, lower, upper, SE, and p-value). Furthermore, the standard tool for checking publication bias uses a funnel plot based on data exploration. The article search and selection process flow is presented in Fig. 1, and the studies included in the meta-analysis are presented in Table 1.

RESULTS

Growth performance and carcass characteristics

A total of 27 articles with seven parameters related to growth performance and carcass characteristics of 30 native chickens and 49 studies that cover eligibility rules are to be accepted in the meta-analysis. In this case, we are measuring intensive rearing systems with free-range rearing systems in native chickens on performance growth and carcass characteristics. Table 2 shows that native chickens have evidence of an effect on the rearing system on live weight parameters (SMD=-1.21; C.I. 95%=-1.73 to -0.687) and carcass weight in grams (SMD=-3.02; C.I. 95%=-4.59 to -1.45). Meanwhile, the parameters that were not proven to show a significant difference in the rearing system were the weight gain parameters (SMD=-0.355; C.I. 95%=-0.701 to -1.-10), carcass weight in percent

(SMD=-0.362; C.I. 95%=- 0.946 to 0.221), feed consumption (SMD=-4.19; 95% C.I.=-8.38 to 0.007), FCR (SMD=-0.798; 95% C.I.=-1.68 to 0.08), and mortality (SMD=4.7; 95% C.I.=1.3 to 8.13).

Quality of breast meat

Table 3 shows the meta-analysis of breast meat quality between free-range and intensive-rearing systems. The results obtained on free-range chickens have evidence of the effect of the rearing system on meat quality parameters for b* (SMD=3.048; C.I. 95%=1.31 to 4.79). Meanwhile, parameters that have no evidence of an effect on the rearing system are breast weight parameters in percent (SMD=0.682; C.I. 95%=0.072 to 1.29), meat quality for L* (SMD=1.37; C.I. 95%=0.129 to 2.61), a * (SMD=1.49; C.I. 95%=0.492 to 2.49), pH (SMD=-0.281; C.I. 95%=-1.11 to 0.547), shear force (SMD=-0.469; C.I. 95%=-2.04 to 1.11), WHC (SMD=-0.262; 95% C.I.=-0.758 to 0.234), meat water (SMD=0.808; 95% C.I.=0.128 to 1.487), as well as chemical composition for protein (SMD=0.642; C.I. 95%=-0.105 to 1.39) and fat (SMD=-1.66; C.I. 95%=-3.03 to -0.299), as well as on sensory for aroma (SMD=0.957; C.I. 95%=-1.52 to 3.44), taste (SMD=-0.144; C.I. 95 %=-0.53 to 0.243), juiciness (SMD=-0.517; C.I. 95%=-1.45 to 0.414), and tenderness (SMD=-0.026; C.I. 95%=-0.411 to 0.359).

Quality of thigh meat

Table 4 shows the thigh meat quality meta-analysis results between free-range and intensive-rearing systems. The parameters that were not proven to show a significant difference in the rearing system were the thigh weight parameters (SMD=0.228; C.I. 95%=-0.304 to 0.761), meat quality for L* (SMD=1.19; C.I. 95%=0.109 to 2.27), a* (SMD = 0.842; C.I. 95%=-0.092 to 1.78), b*

Table	1: Studies included in th	e meta-analysis					
Study	Reference	Breed	Sex	Cage (m ² /head)	Area	Rearing Systems	Parameters
1	Martínez-Pérez et al. (2023)	Rhode Island Red	Ma	0.1667		FR and IN	LW, FI, FCR and BWG
2	Ahmad et al. (2019)	Rhode Island Red x Naked Neck, Black Australorp x Naked Neck, Naked Neck	Mix	0.2300		FR and IN	LW
3	Rehman et al. (2016)	Lakha, Mushki, Peshawari, Sindhi	Fe	0.3528		FR and IN	LW and BWG
4	Cheng et al. (2008)	Taiwan Black	Fe	0.1300		FR and IN	CW, LW, BM (We, WHC, SF, pH, Flav, Tend, Juic), and TM (We, WHC, SF, Ph, Flav, Tend, Juic)
5	(Dou et al., 2009)	Gushi	Fe	0.1429		FR and IN	LW, BWG, MW, BM (We, Prot, Fat, WHC, SF, pH) and TM (We)
6	Li et al. (2016)	Lingnanhuang	Fe	0.1250		FR and IN	CW, LW, BM (We, WHC, SF, pH) and TM (We)
7	Mikulski et al. (2011)	Hybrid	Ma	0.1300		FR and IN	CW, LW, FCR, BM (We, Prot, Fat, WHC, pH, Col, Flav, Tend, Juic) and TM (We, WHC, pH, Col)
8	Wang et al. (2009)	Gushi	Fe	0.1429		FR and IN	CW, MW, LW, BWG, BM (We, Prot, Fat, WHC, SF, pH) and TM (We)
9	Bosco et al. (2014)	Native chicken	Ma	0.1000		IN	LW, FI, FCR, BWG, and Mort
10	Jiang et al. (2011)	Chinese Local Chicken (Local x Broiler)	Ma	0.3800		FR and IN	CW, LW, FI, FCR, BWG, BM (We, SF, pH, Col) and TM (We)
11	Niranjan et al. (2008)	C1 cross, C2 cross, Vanaraja 234, Gramapriya 212	Ma	-		FR	LW
12	Jin et al. (2019)	Yellow chicken (wannan)	Ma	0.6667		FR	CW, LW, FI, FCR, BWG, Mort, BM (We, SF, pH, Col) and TM (We)
13	Sogunle et al. (2012)	Harco black. Novogen	Ma	0.2500		FR and IN	LW, FI, FCR, BWG, Mort
14	Evaris et al. (2020)	Rhode Island Red	Fe	2.9400		FR and IN	CW, LW, and BM (We, Fat)
15	Rehman et al. (2022)	Aseel	Ma	1.1577		FR and IN	CW, LW, BWG, BM (We), and TM (We)
16	Ahmad et al. (2019)	Rhode Island Red x Naked Neck, Black Australorp x Naked Neck, Naked Neck	Ma	0.2300		FR and IN	CW and LW
17	Duran (2004)	Azul	Ma	2.7174		IN	CW, LW, BM (We, Flav, Tend, Juic) and TM (We)
18	Lin et al. (2014)	Taiwan	Fe	1.2500		FR and IN	BM (Flav, Tend, Juic) and TM (Flav, Tend, Juic)
19	Volk et al. (2011)	Slovenian hybrid Prelux-G	Ma	8.0000		FR	CW, BM (Ph and Col), and TM (pH and Col)
20	Calik (2017)	Yellowleg Partridge breed (Z-33)	Ma	1.0000		IN	LW, BM (We, WHC, pH, Col, Flaf, Tend, Juic) and TM (We, WHC, pH, Col, Flav, Tend, Juic)
21	Cerolini et al. (2019)	Milanino	Fe and Ma	2.0000	and	IN	CW, and LW
22	Miguel et al. (2008)	Castellana Negra	Ma	1.0000		IN	CW, LW, BM (pH, Col, Flav, Tend, Juic), and TM (pH, and Col)
23	Mosca et al. (2018)	Milanino	Fe and Ma	8.0000		FR	CW, LW, BM (WHC, pH, Col), and TM (WHC, pH, Col)
24	Tong et al. (2014)	Suqin yellow	Ma	2.4000		FR	CW, LW, FI, FCR, BWG, Mort, BM (We, WHC, SF, pH, Col) and TM (We)
25	Yamak et al. (2016)	Alectoris chukar	Fe and Ma	3.5000		FR and IN	CW, LW, BM (We, pH, Col) and TM (We, pH, Col)
26	Olaniyi et al. (2012)	Harco black, Novogen	-	0.2500		FR and IN	LW, FI, FCR, and Mort
27	Sosnówka-Czajka et al. (2017)	Yellowleg Partridge (Z-33), Rhode Island Red (R-11)	Mix	1.0000		FR and IN	LW, BM (WHC, pH, Col), and TM (WHC, pH, Col)

Ma=male; Fe=female; Mix=mix; FR=free-range; IN=intensive; LW=live weight; FI=feed intake; FCR=feed conversion ratio; BWG=body weight gain; CW=carcass weight; BM=breast meat; We=weight; WHC=water holding capacity; SF=shear force; Flav=flavor; Tend=Tenderness; Juic=Juiciness; TM=thigh meat; MW=meat water; Prot=protein; Col=colour; Mort=mortality

(SMD=0.391; C.I. 95%=-0.712 to 1.5), pH (SMD=-1.01; C.I. 95%=-2.24 to 0.217), WHC (SMD=0.011; C.I. 95%=-0.433 to 0.456), sensory for flavor (SMD=0.179; C.I. 95%=-0.24 to 0.598), juiciness (SMD=0.117; C.I. 95%=-0.301 to 0.536), and tenderness (SMD=0.18; C.I. 95%=-0.239 to 0.599).

DISCUSSION

The meta-analysis results showed a significant influence between the rearing system for native chickens on live weight and carcass weight in grams but not significantly on the parameters of body weight gain, carcass weight in percent, feed intake, FCR, and mortality. This situation is caused by the different types or breeds of native chickens and other types of feed in each study. Studies have shown that chicken breeds vary in various regions, so the phenotype and genotype can differ depending on the breed. For example, Saudi native chickens have different morphological appearances and colors and feather patterns (black, black striped, dark brown, light brown, and grey), where the black breed has a lighter body weight than other breeds of the Saudi native chicken (Fathi et al. 2017).

According to research by Wang et al. (2023), ten local chicken breeds native to China show genetic and performance variations related to body weight and different body sizes (small, medium and large). The same can also be seen in the growth curves for four local Italian chicken breeds and two crosses showing different performances

 Table 2: Comparison of growth performance and carcass characteristics between free-range and intensive system

Parameters	Specification		df	Effect Size (C.I. 95%)							
			2		SMD (lower;	upper)	SE	P Value			
Live weight (g)	General		30	-1.21	-1.73	-0.687	0.266	< 0.001			
0	Cage area (m ² /h)	0.125		-8.53	-9.34	-7.73	0.41	NA			
	0	0.13		-0.057	-0.358	0.244	0.154	0.712			
		0.143		-1.48	-2.22	-0.742	0.376	< 0.001			
		0.167		-0.088	-0.446	0.27	0.183	NA			
		0.23		-0.767	-1.56	0.021	0.402	0.056			
		0.25		-1.47	-1.65	-1.29	0.091	< 0.001			
		0.353		-0.173	-0.756	0.41	0.297	0.56			
		0.38		0.085	-1.16	1.33	0.633	NA			
		1		0.627	0.446	0.808	0.092	< 0.001			
		1.16		-0.288	-0.62	0.04	0.168	NA			
		2.94		-0.677	-1.12	-0.237	0.224	NA			
		3.5		-3.62	-5.07	-2.161	0.743	< 0.001			
Body weight gain (g/h/d)	General		10	-0.355	-0.701	-0.010	0.176	0.044			
	Sex	Ma		-0.412	-0.733	-0.092	0.164	0.012			
		Fe		-0.44	-1.33	0.445	0.451	0.33			
	Cage area (m ² /h)	0.143		-2.3	-3.14	-1.45	0.43	< 0.001			
		0.167		-0.009	-0.367	0.349	0.183	NA			
		0.25		-0.72	-0.885	-0.555	0.084	< 0.001			
		0.353		0.221	-0.152	0.595	0.19	0.245			
		0.38		0.053	-1 19	1 29	0.633	NA			
		11.6		-0.215	-0 543	0.112	0.055	NA			
Carcass weight (%)	General	11.0	7	-0.362	-0.946	0.221	0.298	0 224			
	Sex	Ma	,	-0.295	-1 24	0.647	0.290	0.539			
	bea	Fe		-0.449	-0.702	-0.195	0.129	<0.001			
	Cage area (m^2/h)	0.125		-0.46	-0.716	-0.204	0.12)	NA			
	Cage area (III /II)	0.125		-0.028	-1.01	0.952	0.151	NA			
		0.143		-0.01	-1.61	1.59	0.817	NA			
		0.145		0.167	-0.773	1.11	0.017	0.728			
		0.38		0.107	-0.773	1.02	0.48	NA			
		1.16		-1.87	-0.758	-1.2	0.346	NΔ			
Carcass weight (g)	General	1.10	7	-1.87	-2.55	-1.2	0.340	<0.001			
Carcass weight (g)	Sev	Ma	/	-3.02	-4.39	-1.45	0.8	<0.001			
	SCA	Fo		-4.5	-0.04	-2.39	0.831	0.001			
	$C_{ace} area (m^2/h)$	0.13		-2.07	-3.7	-0.449	0.318	0.012 NA			
	Cage area (III /II)	2.04		-0.31	-0.935	1.06	0.223	NA			
		2.94		4.27	5.61	2.0	0.223	<0.001			
Food intaka (g)	Conoral	5.5	5	-4.27	-5.01	-2.9	0.085	< 0.001			
reed intake (g)	Care area (m^2/h)	0.167	5	-4.19	-0.30	0.007	2.14	0.05 NA			
	Cage area (III /II)	0.107		-0.025	-0.362	0.333	0.165	INA <0.001			
		0.23		-0.30	-9.1	-4.05	1.29	<0.001 NA			
$ECP(\emptyset)$	Comoral	0.58	6	0.143	-1.1	1.39	0.055	NA 0.075			
FCR (%)	General (m^2/h)	0.12	0	-0.798	-1.08	0.08	0.448	0.075 NA			
	Cage area (m ² /n)	0.13		0.213	-0.132	0.558	0.176	INA NA			
		0.10/		-0.30/	-0.728	-0.000	0.184	INA -0.001			
		0.25		-1.54	-2.44	-0.632	0.461	<0.001			
M	Comon 1	0.38	2	0.19	-1.05	1.452	0.634	NA 0.007			
Mortality (%)	General	0.05	3	4.7	1.5	8.13	1./4	0.007			
	Cage area (m ² /h)	0.25		4./	1.3	8.13	1./4	0.007			

Ma=male; Fe=female; C.I.=confidence interval; *df*=degrees of freedom, equal to n (number of observation) – 1; SE=standard error; SMD=standardized mean difference.

 Table 3: The quality of breast meat in native chicken between free-range and intensive

Parameter	Specification		df	Effect Size (C.I. 95%)						
				SMD (lov	wer; upper)		SE	p Value		
Breast weight (%)	General		13	0.682	0.072	1.29	0.311	0.028		
• · ·	Sex	Ma		0.168	-0.489	0.826	0.336	0.616		
		Fe		1.02	0.078	1.97	0.483	0.034		
	Cage area (m ² /h)	0.125		-0.573	-0.831	-0.315	0.132	NA		
		0.13		0.342	-2.02	2.7	1.2	0.776		
		0.143		1.2	0.489	1.91	0.362	< 0.001		
		0.38		-0.183	-1.06	0.695	0.448	NA		
		1.16		0.191	-0.376	0.758	0.289	NA		
		2.94		1.61	1.11	2.1	0.251	NA		
		3.5		0.979	0.166	1.79	0.415	0.018		
Meat Quality										
L*	General		9	1.37	0.129	2.61	0.633	0.03		
	Sex	Ma		2.21	0.023	4.39	1.11	0.048		
		Fe		4.02	-1.93	9.97	3.04	0.186		
		Mi		-0.507	-1.51	0.499	0.513	0.323		
	Cage area (m ² /h)	0.13		-0.434	-1.43	0.557	0.506	NA		
	-	0.38		0.043	-0.834	0.919	0.447	NA		
		1		-0.507	-1.51	0.499	0.513	0.323		
		3.5		5.25	1.58	8.93	1.88	0.005		
a*	General		9	1.49	0.492	2.49	0.51	0.003		
	Sex	Ma		1.84	-0.023	3.7	0.95	0.053		
		Fe		3 4 1	0 164	6 65	1.66	0.039		

		Mi		0.36	-0.804	1.53	0.594	0.544
	~			0.50	0.001	1.00	0.571	0.511
	Cage area (m ² /h)	0.13		0.81	-0.209	1.83	0.52	NA
	e (0.29		0.000	0 779	0.076	0 4 4 7	NIA
		0.38		0.099	-0.778	0.970	0.447	INA
		1		0.36	-0.804	1.53	0.594	0.544
		2.5		0.54	1.05	6.00	1.07	0.005
		3.5		3.54	1.05	6.03	1.27	0.005
h*	General		9	3 048	1 31	479	0.887	<0.001
0	General			5.040	1.51	4.77	0.007	<0.001
	Sex	Ma		1.86	-0.308	4.03	1.11	0.093
		Fe		1 95	-1.6	115	3 3/	0.130
		10		4.75	-1.0	11.5	5.54	0.157
		Mi		3.8	1.01	6.59	1.4	0.008
	$C_{ace} area (m^2/h)$	0.13		0 572	0.428	1.57	0.51	NΛ
	Cage area (III /II)	0.15		0.572	-0.428	1.57	0.51	IN/A
		0.38		-0.387	-1.27	0.497	0.451	NA
		1		20	1.01	6 50	1 4 2	0.000
		1		3.0	1.01	0.39	1.42	0.008
		3.5		4.37	0.809	7.92	1.81	0.016
TT	C 1		10	0.001	1 1 1	0.547	0.400	0.506
рн	General		13	-0.281	-1.11	0.547	0.422	0.506
	Sex	Ma		-1.28	-4 15	1 58	1 46	0.38
	Bex	-		1.20	4.15	1.50	1.40	0.50
		Fe		-0.369	-1.64	0.901	0.648	0.569
		Mi		0.074	0.664	0.813	0 377	0.843
		1011		0.074	-0.004	0.015	0.577	0.045
	Cage area (m ² /h)	0.125		0.854	0.59	1.12	0.135	NA
	e (0.12		0.642	0.162	1 45	0.411	0.117
		0.15		0.045	-0.102	1.45	0.411	0.117
		0.143		-0.803	-1.48	-0.123	0.347	0.021
		0.29		0.200	0 671	1.00	0.449	NIA
		0.38		0.208	-0.0/1	1.09	0.448	NA
		1		0.074	-0.664	0.813	0.377	0.843
		2.5		2.5	0.00	0.1	0.00	0.001
		3.5		-3.5	-9.09	2.1	2.86	0.221
Shear force (Kg)	General		4	-0 469	-2.04	1.11	0.803	0.559
Shear force (Hg)	General			0.107	2.01	0.040	0.000	0.557
	Sex	Ma		-0.657	-1.56	0.243	0.459	NA
		Fe		-0.418	-2 37	1 53	0 99/	0.674
		10		-0.410	-2.51	1.55	0.774	0.074
	Cage area (m ² /h)	0.125		-2.2	-2.53	-1.89	0.164	NA
	e (0.13		1 56	0.85	2 27	0.361	NΛ
		0.15		1.50	0.05	2.21	0.501	
		0.143		-0.518	-1.18	0.146	0.339	0.126
		0.38		0.657	1 56	0.243	0.450	NΛ
		0.50		-0.057	-1.50	0.243	0.457	
WHC (%)	General		6	-0.262	-0.758	0.234	0.253	0.301
	Sex	Ma		-1 33	-2.41	-0 244	0 552	NΔ
	Ben	5		1.55	2.11	0.211	0.552	0.075
		Fe		0.201	-0.02	0.423	0.113	0.075
		Mi		-0.602	-18	0 593	0.61	0 323
	G (24)	0.105		0.002	0.017	0.596	0.12	20
	Cage area (m ² /n)	0.125		0.272	0.017	0.526	0.13	NA
		0.13		-0.71	-1 706	0.285	0 508	0.162
		0.15		0.71	1.700	0.205	0.500	0.102
		0.143		0.281	-0.376	0.937	0.335	0.402
		1		-0.602	-1 797	0 593	0.61	0 3 2 3
		1		-0.002	-1.///	0.575	0.01	0.525
Meat water (%)	General		1	0.808	0.128	1.487	0.347	0.02
	Sov	Ea		0 808	0.129	1 497	0 247	0.02
	Sex	re		0.000	0.120	1.40/	0.347	0.02
	Cage area (m ² /h)	0.143		0.808	0.128	1.487	0.347	0.02
Chamical Composition	e , ,							
Chemical Composition	~ .							
Protein (%)	General		2	0.642	-0.105	1.39	0.381	0.092
	Sov	Ma		1.42	0 322	2 52	0.550	NΛ
	SEX	Ivia		1.42	0.322	2.32	0.559	IN/A
		Fe		0.293	-0.364	0.949	0.335	0.383
	$C_{1} = c_{1} + c_{2} + c_{1} + c_{2} + c_{2$	0.12		1.40	0.222	0.51	0.550	NT A
	Cage area (III /II)	0.15		1.42	0.522	2.31	0.559	INA
		0.143		0.293	-0.364	0.949	0.335	0.383
$E_{ot}(0/)$	Comoral		2	1 66	2.02	0.200	0 606	0.017
Fat (%)	General		3	-1.00	-3.03	-0.299	0.090	0.017
	Sex	Ma		-0.189	1.17	0.793	0.501	NA
		P		2.2	2 525	0.067	0 (01	0.001
		ге		-2.2	-3.535	-0.867	0.681	0.001
	Cage area (m^2/h)	0.13		-0.189	-1.17	0.793	0.501	NA
	Cuge urea (III /II)	0.15		0.10)		0.775	0.501	0.4.47
		0.143		-2.17	-5.1	0.763	1.5	0.147
		2 94		-2.1	-2.63	-1.56	0.272	NΔ
G		2.74		2.1	2.05	1.50	5.272	1 12 1
Sensory								
Aroma	General		1	0.957	-1.52	3 44	1.27	0.449
. ii oiniu	a a a a a a a a a a a a a a a a a a a		-	0.00	1.02	2.54	0.640	NT 4
	Sex	Ма		2.28	1.03	3.54	0.643	NA
		Fe		-0 248	-0.87	0 374	0 317	NA
	G (21)	0.10		0.210	1.50	0.071	1.07	0.440
	Cage area (m ² /n)	0.13		0.957	-1.52	3.44	1.27	0.449
Flavor	General		2	-0.144	-0.53	0.243	0.197	0.466
	Corr	N	-	0 700	1 70	0.201	0 5 1 5	NIA
	Sex	Ivia		-0.709	-1.72	0.501	0.515	INA
		Fe		-0.047	-0.465	0.371	0.213	0.827
	$C_{0} \alpha_{0} \alpha_{0$	0.12		0 279	0.000	0.251	0.27	0.202
	Cage area (m ⁻ /n)	0.15		-0.278	-0.800	0.231	0.27	0.505
		1.25		0.01	-0.556	0.576	0.289	NA
Inicipass	General		n	0.517	1 45	0.414	0 475	0.277
JUICIIICSS	General		2	-0.517	-1.43	0.414	0.473	0.277
	Sex	Ma		-1.996	-3.195	-0.796	0.612	NA
		Ea		0.01	0 120	0.409	0.212	0.064
		re		-0.01	-0.428	0.408	0.215	0.904
	Cage area (m ² /h)	0.13		-0.956	-2.85	0.938	0.966	0.323
		1 25		0.03	_0 535	0 596	0 280	NA
	a .	1.23	-	0.05	-0.555	0.390	0.209	0.000
Tenderness	General		2	-0.026	-0.411	0.359	0.197	0.896
	Sev	Ma		-0 473	-1 47	0.521	0 507	NΔ
	JUA .	1v1a		-0.473	-1.++/	0.321	0.507	11/1
		Fe		0.053	-0.365	0.471	0.213	0.803
	Cage area (m^2/h)	0.13		-0.101	-0 627	0.425	0.268	0 706
	Cage area (III /II)	0.15		-0.101	-0.027	0.725	0.200	0.700
		1.25		0.061	-0.505	0.627	0.289	NA

 L^* =(Lightness) represents the lightness or darkness of meat; a*=(Redness-Greenness) represents the position of meat color on the red-green axis; b*=(Yellowness-Blueness) represents the position of meat color on the yellow-blue axis; Ma=male; Fe=female; Mi=Mix; WHC=water holding capacity; C.I.=confidence interval; df=degrees of freedom, equal to n (number of observation) – 1; SE=standard error; SMD=standardized mean difference. (Mancinelli et al. 2023). Furthermore, research by Chaiwang et al. (2023) reported that four different chicken breeds kept in the same conditions and with the same commercial feed also showed other growth performance in terms of growth performance (body weight, feed intake, body weight gain, feed conversion ratio), carcass trait (live weight, carcass percentage, carcass weight), meat quality traits, amino acid content, rancidity of chicken meat, and nucleotide content and derivatives. However, if the feed is given with different metabolic energy (ME) and crude protein (CP) contents, it will affect feed intake and mortality rates (Chang et al. 2023). Apart from that, the effect size value generally shows that the treatment is lower than the control, which means that the intensive maintenance system has better performance and carcass characteristics than the free-range maintenance system; one example of this is the higher amount of intramuscular fat in intensive raised chicken as opposed to free-range chickens (Yang et al. 2009; Yang et al. 2014). This is because controlled management is the primary factor that promotes improved chicken production in the intensive system. A review of indigenous chickens in East Africa with intensive, semi-intensive, and free-range rearing systems shows that they correlate with different performance and the highest body weight in intensive rearing systems (Mujyambere et al. 2022).

The results show a significant influence between the rearing system on native chickens only on the meat quality parameter for b*. Meanwhile, other parameters did not substantially affect the rearing system for native chickens. The meat quality in the research literature, especially the breast meat part, is more dominant in yellowness. According to (Mikulski et al. 2011), the feed consumed by the chicken causes the yellow breast meat in poultry. On the other side, access in and out of the cage led to greater amounts of polyunsaturated fat in the muscles of chickens (Sokołowicz et al. 2016). This is supported by several previous studies regarding yellowish meat color (Freitas et

al. 2015; Chai and Sheen 2021). Meat color in poultry is essential to consumer interest (Kim et al. 2020).

Intensive rearing and free-range rearing systems for native chickens have no evidence of an effect on thigh meat quality. These results were driven by reference studies using different native chicken breeds. According to Chaiwang et al. (2023), performance, carcass, meat quality traits, chemical content, nucleotides, and derivatives are different for each chicken breed (commercial broilers, Mae Hong Son Thai native chickens, Pradu Hang Dam Thai native chickens, and male layer chickens) in the section tight.

Several research results related to thigh meat have also been carried out for research objects on native chickens (Mahuang and Tuer) and commercial broilers (Deng et al. 2022); mixed-sex Thai indigenous crossbred chickens (Hang et al. 2018); Korean native chickens (Javasena et al. 2015); Korat hybrid chickens (Katemala et al. 2021); crossbred native chickens, namely LBC chickens (Layer-Broiler crossed with Chee), LSC chickens (Layer-Shanghai crossed with Chee), and LSRBC chickens (Laver-Shanghai Road Bar crossed with Chee) (Promket et al. 2016); two Chinese native chickens, namely Wuding chicken and Yanji silky fowl chicken (Xiao et al. 2021); and local Chinese chicken breeds, namely Beijing-you chickens (Zheng et al. 2019). Besides breed factors, meat quality is also influenced by genetic configuration, environment, feed, and stress (Chaiwang et al. 2023). The general effect size values obtained show that the treatment is higher than the control, where the free-range maintenance system has a better quality of thigh meat than the intensive maintenance system. According to Yamak et al. (2016), the quality of thigh meat with a free-range rearing system is better than that of an intensive rearing system. Various chicken species may cause performance variations in research studies, feeds, and rearing systems (Zheng et al. 2019). The freerange rearing system benefits animal welfare compared to other rearing systems (Stefanetti et al. 2023).

Table 4: The quality of thigh meat in native chicken between free-range and intensive

Parameter	Specification			Effect Size (C.I. 95%)						
				SMD (lo	wer; upper)		SE	p Value		
Thigh weight (%)	General		12	0.228	-0.304	0.761	0.272	0.4		
	Sex	Ma		0.07	-0.868	1.01	0.479	0.883		
		Fe		0.392	-0.232	1.02	0.318	0.218		
	Cage area (m ² /h)	0.125		0.406	0.15	0.661	0.13	NA		
		0.13		0.071	-0.454	0.595	0.268	0.791		
		0.143		1.05	0.357	1.75	0.356	0.003		
		0.38		0	-0.877	0.877	0.447	NA		
		1.16		-1.53	-2.17	-0.882	0.328	NA		
		3.5		0.535	-0.658	1.73	0.609	0.379		
Meat Quality										
L*	General		8	1.19	0.109	2.27	0.551	0.031		
	Sex	Ma		1.23	-1.2	3.65	1.24	0.321		
		Fe		3.23	0.768	5.68	1.25	0.01		
		Mi		0.128	-0.312	0.568	0.224	0.569		
	Cage area (m ² /h)	0.13		-1.54	-2.65	-0.42	0.569	NA		
		1		0.128	-0.312	0.568	0.224	0.569		
		3.5		2.85	0.914	4.79	0.987	0.004		
a*	General		8	0.842	-0.092	1.78	0.477	0.077		
	Sex	Ma		1.47	-1.62	4.57	1.58	0.351		
		Fe		0.758	-0.691	2.21	0.739	0.305		
		Mi		0.68	0.229	1.13	0.23	0.003		
	Cage area (m ² /h)	0.13		-0.356	-1.34	0.632	0.504	NA		
		1		0.68	0.229	1.13	0.23	0.003		

		3.5		1.38	-0.581	3.34	1	0.168
b*	General		8	0.391	-0.712	1.5	0.563	0.487
	Sex	Ma		0.129	-2.38	2.63	1.28	0.919
		Fe		0.871	-1.87	3.61	1.4	0.533
		Mi		0.795	0.021	1.57	0.395	0.044
	Cage area (m^2/h)	0.13		-0.889	-1.92	0.138	0.524	NA
		1		0.795	0.021	1.57	0.395	0.044
		3.5		0.744	-1.42	2.9	1.11	0.501
pH	General		9	-1.01	-2.24	0.217	0.626	0.107
WHC (%)	General		3	0.011	-0.433	0.456	0.227	0.96
	Sex	Ma		-0.328	-1.39	0.738	0.544	NA
		Fe		0.057	-0.563	0.677	0.316	NA
		Mi		0.071	-0.857	1	0.474	0.88
	Cage area (m^2/h)	0.13		-0.04	-0.576	0.495	0.273	0.882
		1		0.071	-0.857	1	0.474	0.88
Sensory								
Flavor	General		1	0.179	-0.24	0.598	0.214	0.403
	Sex	Fe		0.179	-0.24	0.598	0.214	0.403
	Cage area (m^2/h)	0.13		0.292	-0.331	0.915	0.318	NA
		1.25		0.086	-0.48	0.652	0.289	NA
Juiciness	General		1	0.117	-0.301	0.536	0.213	0.583
	Sex	Fee		0.117	-0.301	0.536	0.213	0.583
	Cage area (m^2/h)	0.13		0.173	-0.448	0.794	0.317	NA
		1.25		0.071	-0.495	0.637	0.289	NA
Tenderness	General		1	0.18	-0.239	0.599	0.214	0.399
	Sex	Fe		0.18	-0.239	0.599	0.214	0.399
	Cage area (m^2/h)	0.13		0.295	-0.329	0.918	0.318	NA
		1.25		0.086	-0.48	0.652	0.289	NA

L*=(Lightness) represents the lightness or darkness of meat; a*=(Redness-Greenness) represents the position of meat color on the redgreen axis; b*=(Yellowness-Blueness) represents the position of meat color on the yellow-blue axis; Ma=male; Fe=female; Mi=Mix; WHC=water holding capacity; C.I.=confidence interval; df=degrees of freedom, equal to n (number of observation) – 1; SE=standard error; SMD=standardized mean difference.

Conclusion

This meta-analysis provides critical scientific insights into the effects of rearing systems for native chickens in many regions on their growth performance and carcass characteristics. The conclusion is that the performance and carcass characteristics are better in the intensive system. Meanwhile, the better breast meat quality parameter is in the free-range system. Future research should concentrate on modern free-range rearing systems to evaluate the impact on chicken growth performance and meat quality in various regions.

Conflict of interest: There is no conflict of interest with any organization regarding the discussion in this manuscript.

Author's Contribution

JAL, S, and MMS: conceptualization, formal analysis, investigation, methodology, computing resources, validation, data curation, writing-original draft, writing, and editing. JAL, RA, ZW, SE, TS, CS, S, SYH, and MMS: investigation, data curation, formal analysis, validation, writing-review, and editing.

REFERENCES

- Ahmad S, Mahmud A, Hussain J and Javed K, 2019. Morphological and carcass traits of three chicken genotypes under free-range, semiintensive, and intensive housing systems. Turkish Journal of Veterinary and Animal Sciences 43(3): 342-352. https://doi.org/10.3906/vet-1902-9
- Ariyanti D, Sasongko NA, Fansuri MH, Fitriana EL, Nugroho RA and Pratiwi SA, 2023. Retrofitting of concrete for rigid

pavement using bacterial: A meta-analysis. Science of the Total Environment 902: 166019. <u>https://doi.org/10.1016/j.scitotenv.2023.166019</u>

- Bosco AD, Mugnai C, Rosati A, Paoletti A, Caporali S and Castellini C, 2014. Effect of range enrichment on performance, behavior, and forage intake of free-range chickens. Journal of Applied Poultry Research 23(2): 137– 145. <u>https://doi.org/10.3382/japr.2013-00814</u>
- Calik J, 2017. Effect of caponizing yellowleg partridge (z-33) cockerels on body weight and meat quality. Acta Scientiarium Polonorum Zootecnica 14(1): 51-60.
- Cerolini S, Vasconi M, Sayed AA, Iaffaldano N, Mangiagalli MG, Pastorelli G, Moretti VM, Zaniboni L and Mosca F, 2019. Free-range rearing density for male and female milanino chickens: carcass yield and qualitative meat traits. Journal of Applied Poultry Research 8(4): 1349-1358. <u>https://doi.org/10.3382/japr/pfz058</u>
- Chai HE and Sheen S, 2021. Effect of high pressure processing, allyl isothiocyanate, and acetic acid stresses on Salmonella survivals, storage, and appearance color in raw ground chicken meat. Food Control 30(40): 1-5. <u>https://doi.org/10.</u> 1016/j.foodcont.2020.107784
- Chaiwang N, Marupanthorn K, Krutthai N, Wattanakul W, Jaturasitha S, Arjin C, Sringarm, K and Setthaya P, 2023. Assessment of nucleic acid content, amino acid profile, carcass and meat quality of thai native chicken. Poultry Science 102(11): 1-5. <u>https://doi.org/10.1016/j.psj.2023.</u> 103067
- Chang C, Zhang QQ, Wang HH, Chu Q, Zhang J, Yan ZX, Liu HG and Geng AL, 2023. Dietary metabolizable energy and crude protein levels affect pectoral muscle composition and gut microbiota in native growing chickens. Poultry Science 102(2): 1-16. <u>https://doi.org/10.1016/j.psj.2022.102353</u>
- Chen X, Jiang W, Tan HZ, Xu GF, Zhang XB, Wei S and Wang XQ, 2013. Effects of outdoor access on growth performance, carcass composition, and meat characteristics of broiler

chickens. Poultry Science 92(2): 435-443. <u>https://doi.org/10.</u> 3382/ps.2012-02360

- Cheng FY, Huang CW, Wan TC, Liu YT, Lin LC and Lou CY, 2008. Effects of free-range farming on carcass and meat qualities of black-feathered taiwan native chicken. Asian-Australasian Journal of Animal Sciences 21(8): 1201-1206. https://doi.org/10.5713/ajas.2008.80080
- Deng S, Liu R, Li C, Xu X and Zhou G, 2022. Meat quality and flavor compounds of soft-boiled chickens: effect of chinese yellow-feathered chicken breed and slaughter age. Poultry Science 101(12): 1-13. <u>https://doi.org/10.1016/j.psj.2022.</u> 102168
- Dou TC, Shi SR, Sun HJ and Wang KH, 2009. Growth rate, carcass traits and meat quality of slow-growing chicken grown according to three raising systems. Animal Science Papers and Reports 27(4): 361-369.
- Duran M, 2004. The effect of caponization on production indices and carcass and meat characteristics in free-range extremeña zzul chickens. Spanish Journal of Agricultural Research 2(2): 211-216. <u>https://doi.org/10.5424/sjar/2004022-75</u>
- Evaris EF, Franco LS and Castro CS, 2020. Productive performance and carcass yield of egg type male chickens raised with outdoor access in the tropics. Tropical Animal Health and Production 52(6): 3225-3232. <u>https://doi.org/10.1007/s11250-020-02348-w</u>
- Fanatico AC, Pillai PB, Emmert JL and Owens CM, 2007. Meat quality of slow-and fast-growing chicken genotypes fed lownutrient or standard diets and raised indoors or with outdoor access. Poultry Science 86(10): 2245-2255. <u>https://doi.org/ 10.1093/ps/86.10.2245</u>
- Fathi MM, Al-Homidan I, Motawei MI, Abou-Emera OK and El-Zarei MF, 2017. Genetics and genomics: evaluation of genetic diversity of saudi native chicken populations using microsatellite markers. Poultry Science 96(3): 530-536. <u>https://doi.org/10.3382/ps/pew357</u>
- Fitra D, Ulupi N, Arief II, Mutia R, Abdullah L and Erwan E, 2021. Development of chicken production on free-range system. Wartazoa 31(4): 175-184. <u>http://dx.doi.org/10. 14334/wartazoa.v31i4.2683</u>
- Freitas ER, Borges AS, Pereira AL, Abreu VK, Trevisan MS and Watanabe PH, 2015. Effect of dietary ethanol extracts of mango (mangifera indica l.) on lipid oxidation and the color of chicken meat during frozen storage. Poultry Science 94(12): 2989-2995. <u>https://doi.org/10.3382/ps/pev295</u>
- Hang TT, Molee W and Khempaka S, 2018. Linseed oil or tuna oil supplementation in slow-growing chicken diets: Can their meat reach the threshold of a "high in n-3 polyunsaturated fatty acids" product? Journal of Applied Poultry Research 27(3): 389-400. https://doi.org/10.3382/japr/pfy010
- Hansen BG and Østerås O, 2019. Farmer welfare and animal welfare-exploring the relationship between farmer's occupational well-being and stress, farm expansion and animal welfare. Preventive Veterinary Medicine 170: 1-36. https://doi.org/10.1016/j.prevetmed.2019.104741
- Jayasena DD, Jung S, Bae YS, Park HB, Lee JH and Jo C, 2015. Comparison of the amounts of endogenous bioactive compounds in raw and cooked meats from commercial broilers and indigenous chickens. Journal of Food Composition and Analysis 37: 20-24. <u>https://doi.org/10.1016</u> /j.jfca.2014.06.016
- Jiang S, Jiang Z, Lin Y, Zhou G, Chen F and Zheng C, 2011. Effects of different rearing and feeding methods on meat quality and antioxidative properties in chinese yellow male broilers. British Poultry Science 52(3): 352-358. <u>https://doi.org/10.1080/00071668.2011.569926</u>
- Jin S, Yang L, Zang H, Xu Y, Chen X, Chen X, Liu P and Geng Z, 2019. Influence of free-range days on growth performance, carcass traits, meat quality, lymphoid organ indices and blood biochemistry of wannan yellow chickens.

Poultry Science 98(12): 6602-6610. <u>https://doi.org/10.3382/</u>ps/pez504

- Katemala S, Molee A, Thumanu K and Yongsawatdigul J, 2021. Meat quality and Raman spectroscopic characterization of Korat hybrid chicken obtained from various rearing periods. Poultry Science 100(2): 1248-1261. <u>https://doi.org/10.1016/j.psj.2020.10.027</u>
- Kim HJ, Kim HJ, Jeon JJ, Nam KC, Shim KS, Jung JH, Kim KS, Choi Y, Kim SH and Jang A, 2020. Comparison of the quality characteristics of chicken breast meat from conventional and animal welfare farms under refrigerated storage. Poultry Science 99(3): 1788-1796. <u>https://doi.org/ 10.1016/j.psj.2019.12.009</u>
- Lewis PD, Perry GC, Farme LJ and Pattersonc RL, 1997. Responses of two genotypes of chicken to the diets and stocking densities typical of uk and "label rouge" production systems: i. performance, behaviour and carcass composition. Meat Science 45(4): 501-516. <u>https://doi.org/10.1016/ S0309-1740(96)00084-8</u>
- Li Y, Luo C, Wang J and Guo F, 2016. Effects of different raising systems on growth performance, carcass, and meat quality of medium-growing chickens. Journal of Applied Animal Research 45(1): 326-330. <u>https://doi.org/10.1080/09712119.</u> 2016.1190735
- Lin CY, Kuo HY and Wan TC, 2014. Effect of free-Range rearing on meat composition, physical properties and sensory evaluation in taiwan game hens. Asian-Australasian Journal of Animal Sciences 27(6): 880-885. <u>https://doi.org/10.5713/ ajas.2013.13646</u>
- Makmur M, Zain M, Sholikin MM, Suharlina and Jayanegara A, 2022. Modulatory effects of dietary tannins on polyunsaturated fatty acid biohydrogenation in the rumen: a meta-analysis. Heliyon 8(7): 1-11. <u>https://doi.org/10.1016/j. heliyon.2022.e09828</u>
- Mancinelli AC, Menchetti L, Birolo M, Bittante G, Chiattelli D and Castellini C, 2023. Crossbreeding to improve local chicken breeds: predicting growth performance of the crosses using the gompertz model and estimated heterosis. Poultry Science 102(8): 1-11. <u>https://doi.org/10.1016/j.psj. 2023.102783</u>
- Marin RH, Freytes P, Guzman D and Jones B, 2001. Effects of an acute stressor on fear and on the social reinstatement responses of domestic chicks to cagemates and strangers. Applied Animal Behaviour Science 71(1): 57–66. <u>https://doi.org/10.1016/S0168-1591(00)00167-2</u>
- Martínez-Pérez M, Sarmiento-Franco L, Sandoval-Castro CA, Santos-Ricalde RH, Safwat AM and García Hernández Y, 2023. Effect of indoor and free-range raising systems on growth performance of male rhode island red chickens. Tropical and Subtropical Agroecosystems 26(2): 1-7. <u>http://doi.org/10.56369/tsaes.4312</u>
- Mcfadden JR, Huffman WE and Huffman WH, 2017. Willingness-to-pay for natural, organic, and conventional foods: the effects of information and meaningful labels. Food Policy 68: 214-232. <u>https://doi.org/10.1016/j.foodpol.2017.</u> 02.007
- Miao ZH, Glatz PC and Ru YJ, 2005. Free-range poultry production - a review. Asian-Australasian Journal of Animal Sciences 18(1): 113-132. https://doi.org/10.5713/ajas.2005.113
- Miguel JA, Ciria J, Asenjo B and Calvo JL, 2008. Effect of caponisation on growth and on carcass and meat characteristics in castellana negra native spanish chickens. Animal 2(2): 305-311. <u>https://doi.org/10.1017/S175173110</u> 7001127
- Mikulski D, Celej J, Jankowski J, Majewska T and Mikulska M, 2011. Growth performance, carcass traits and meat quality of slower-growing and fast-growing chickens raised with and without outdoor access. Asian-Australasian Journal of

Animal Sciences 24(10): 1407-1416. <u>https://doi.org/10.</u> 5713/ajas.2011.11038

- Mosca F, Zaniboni L, Stella S, Kuster CA, Iaffaldano N and Cerolini S, 2018. Slaughter performance and meat quality of Milanino chickens reared according to a specific free-range program. Poultry Science 97(4): 1148-1154. https://doi.org/10.3382/ps/pex439
- Mujyambere V, Adomako K, Olympio SO, Ntawubizi M, Nyinawamwiza L, Mahoro J and Conroy A, 2022. Local chickens in east african region: their production and potential. Poultry Science 101(1): 1-21. <u>https://doi.org/10. 1016/j.psj.2021.101547</u>
- Niranjan M, Sharma RP, Rajkumar U, Reddy B, Chatterjee RN and Battacharya TK, 2008. Comparative evaluation of production performance in improved chicken varieties for backyard farming. International Journal of Poultry Science 7(11): 1128-1131.
- Olaniyi OA, Oyenaiya OA, Sogunle OM, Akinola OS, Adeyemi OA and Ladokun OA, 2012. Free range and deep litter housing systems: effect on performance and blood profile of two strains of cockerel chickens. Tropical and Subtropical Agroecosystems 15(3): 511-523.
- Palupi E, Jayanegara A, Ploeger A and Kahl J, 2012. Comparison of nutritional quality between conventional and organic dairy products: a meta-analysis. Journal of the Science of Food and Agriculture 92(14): 2774-2781. <u>https://doi.org/10. 1002/jsfa.5639</u>
- Pettersson IC, Weeks CA, Wilson LM and Nicol CJ, 2018. Consumer perceptions of free-range laying hen welfare. British Food Journal 118(8): 1999-2013. <u>https://doi.org/10.</u> 1108/BFJ-02-2016-0065
- Promket D, Ruangwittayanusorn K and Somchan T, 2016. The study of carcass yields and meat quality in crossbred native chicken (chee). Agriculture and Agricultural Science Procedia 11: 84-89. <u>https://doi.org/10.1016/j.aaspro.2016.12</u>.014
- Rehman MS, Khan MT, Raziq F, Bughio E, Liaqat S, Gondal MA, Rauf M, Jalees MM, Farooq, Sarwar F, Ayesha AP, Azad A, Azhar M, Asad T, Arslan M, Abbas G and Shakir M, 2022. Evaluation of growth performance and slaughter traits of Aseel chicken grown with and without outdoor access. Research Square 1: 1-20. <u>https://doi.org/10.21203/rs.</u> 3.rs-1179992/v1
- Rehman MS, Mahmud A, Mehmood S, Pasha TN, Javed K, Hussain J and Khan MT, 2016. Production performance of aseel chicken under free range, semi intensive and confinement rearing systems. The Journal of Animal & Plant Sciences 26(6): 1589-1596.
- Scrinis G, Parker C and Carey R, 2017. The caged chicken or the free-range egg? the regulatory and market dynamics of layerhen welfare in the UK, Australia and the USA. Journal of Agricultural and Environmental Ethics 30(6): 783-808. https://doi.org/10.1007/s10806-017-9699-y
- Sokołowicz Z, Krawczyk J and Swiatkiewicz S, 2016. Quality of poultry meat from native chicken breeds - a review. Annals of Animal Science 16(2): 347-368. <u>https://doi.org/10.1515/ aoas-2016-0004</u>
- Sogunle OM, Olaniyi OA, Egbeyale LT, Akinola OS, Shittu TA, Abiola SS, Ladokun AO and Sobay RA, 2012. Free range and deep litter poultry production systems: effect on performance, carcass yield and meat composition of cockerel chickens. Tropical Animal Health and Production 45(1): 281–288. <u>https://doi.org/10.1007/s11250-012-0213-2</u>
- Sosnówka-Czajka E, Skomorucha I and Muchacka R, 2017. Effect of organic production system on the performance and meat quality of two purebred slow-growing chicken breeds.

Annals of Animal Science 17(4): 1197-1213. <u>https://doi.org/</u> 10.1515/aoas-2017-0009

- Stadig LM, Bas Rodenburg T, Reubens B, Aerts J, Duquenne B and Tuyttens FA, 2016. Effects of free-range access on production parameters and meat quality, composition and taste in slow-growing broiler chickens. Poultry Science 95(12): 2971-2978. <u>https://doi.org/10.3382/ps/pew226</u>
- Stefanetti V, Mancinelli AC, Pascucci L, Menchetti L, Castellini C, Mugnai C, Fiorilla E, Miniscalco B, Chiattelli D, Franciosini MP and Proietti PC, 2023. Effect of rearing systems on immune status, stress parameters, intestinal morphology and mortality in conventional and local chicken breeds. Poultry Science 102(12): 1-11. <u>https://doi.org/10. 1016/j.psj.2023.103110</u>
- Tong HB, Wang Q, Lu J, Zou JM, Chang LL and Fu SY, 2014. Effect of free-range days on a local chicken breed: growth performance, carcass yield, meat quality, and lymphoid organ index. Poultry Science 93(8): 1883-1889. <u>https://doi.org/10.3382/ps.2013-03470</u>
- Volk M, Malenšek J, Prevolnik M, Škrlep M, Šegula B, Čandek-Potokar M and Bavec M, 2011. Differences in carcass and meat quality between organically reared cocks and capons. Agriculturae Conspectus Scientificus 76(3): 153-155. <u>https://hrcak.srce.hr/72024</u>
- Wang H, Zhao X, Wen J, Wang C, Zhang X, Ren X, Zhang J, Li H, Muhatai G and Qu L, 2023. Comparative population genomics analysis uncovers genomic footprints and genes influencing body weight trait in chinese indigenous chicken. Poultry Science 102(11): 1-10. <u>https://doi.org/10.1016/j.psj.</u> 2023.103031
- Wang KH, Shi SR, Dou TC and Sun HJ, 2009. Effect of a freerange raising system on growth performance, carcass yield, and meat quality of slow-growing chicken. Poultry Science 88(10): 2219-2223. <u>https://doi.org/10.3382/ps.2008-00423</u>
- Xiao Z, Zhang W, Yang H, Yan Z, Ge C, Liao G and Su H, 2021. 1H NMR-based water-soluble lower molecule characterization and fatty acid composition of Chinese native chickens and commercial broiler. Food Research International 140: 110008. <u>https://doi.org/10.1016/j.foodres.</u> 2020.110008
- Yamak US, Sarica M, Boz MA and Ucar A, 2016. The effect of production system (barn and free-range), slaughtering age and gender on carcass traits and meat quality of partridges (aalectoris chukar). British Poultry Science 57(2): 185-192. <u>https://doi.org/10.1080/00071668.2016.1144920</u>
- Yang Y, Wen J, Fang GY, Li ZR, Dong ZY and Liu J, 2014. The effects of raising system on the lipid metabolism and meat quality traits of slow-growing chickens. Journal of Applied Animal Research 43(2): 147-152. <u>https://doi.org/10.1080/ 09712119.2014.928631</u>
- Yang Y, Gui-you FG, Li Zhongrong L, Furong L and Yulan F, 2009. Effects of different rearing modes on the muscular histological traits and meat tenderness of chicken. Animal Husbandry and Feed Science 1(3): 24-27.
- Zhang C, Razafindrabe RH, Chen K, Zhao X, Yang L, Wang L, Chen X, Jin S and Geng Z, 2018. Effects of different rearing systems on growth performance, carcass traits, meat quality and serum biochemical parameters of chaohu ducks. Animal Science Journal 89(4): 672-678. <u>https://doi.org/10.1111/asj. 12976</u>
- Zheng M, Mao P, Tian X, Guo Q and Meng L, 2019. Effects of dietary supplementation of alfalfa meal on growth performance, carcass characteristics, meat and egg quality, and intestinal microbiota in beijing-you chicken. Poultry Science 98(5): 2250-2259. <u>https://doi.org/10.3382/ps/ pey550</u>