



The Contact/Fumigant Adulticidal Effect of Egyptian Oils against the House Fly, *Musca domestica* (Diptera: Muscidae)

Mohamed M. Baz¹, Randa I. Eltaly², Mustapha Debboun³, Abdelfattah Selim⁴, Ibrahim T. Radwan⁵, Nazeer Ahmed⁶ and Hanem F. Khater*⁷

¹Department of Entomology, Faculty of Science, Benha University, Benha 13518, Egypt

²Entomology Department, Faculty of Science, Al-Azhar University, Girls Branch, Cairo 11884, Egypt

³Delta Mosquito & Vector Control District, 1737 W. Houston Ave, Visalia, CA 93291, USA

⁴Department of Animal Medicine (Infectious Diseases), Faculty of Veterinary Medicine, Benha University, Toukh 13736, Egypt; ⁵Supplementary General Sciences Department, Faculty of Oral and Dental Medicine, Future University in Egypt, Cairo 11835, Egypt; ⁶Department of Agriculture, University of Swabi, Anbar 23561, Khyber Pakhtunkhwa Pakistan

⁷Parasitology Department, Faculty of Veterinary Medicine, Benha University, Toukh 13736, Egypt

*Corresponding author: hanemkhater@gmail.com; hanem.salem@fvvm.bu.edu.eg

Article History: 22-600

Received: 23-Apr-22

Revised: 01-Jun-22

Accepted: 05-Jun-22

ABSTRACT

The common housefly, *Musca domestica* (Diptera: Muscidae) is a cosmopolitan pest of a major medical and veterinary importance acting as a mechanical vector of pathogens. This study evaluated the toxicity of six essential oils against adult *M. domestica* through contact/fumigant toxicity bioassays. The most effective oil was radish, followed by camphor, mustard, thyme, garlic, and rosemary. The mortality (MO)% 60 min, post-treatment (PT) with 2% of camphor, garlic, mustard, radish, rosemary, and thyme oils were 90.00, 83.33, 86.66, 93.33, 83.33, and 86.66%, respectively. After treatment for one hour, the median lethal concentration, LC₅₀ of camphor, garlic, mustard, radish, rosemary, and thyme were 0.305, 0.450, 0.366, 0.250, 0.475, and 0.405%, respectively. The corresponding LC₉₉ values were 2.54, 3.19, 2.92, 2.30, 3.14, and 2.96%, respectively. Their toxicity indices were 81.97, 55.56, 68.31, 100.00, 52.63, and 61.73%, respectively. After treatment with 2%, their median lethal time (LT₅₀) values were 16.52, 19.99, 18.29, 16.44, 18.73, and 17.92 min, respectively. The adulticidal effect of radish, mustard, camphor, and rosemary against house flies had a novelty record and the current investigation concluded that all applied oils were effective adulticides. Radish oil was the most potent oil followed by camphor, mustard, and thyme, whereas garlic and rosemary were less effective oils; therefore, the promising oils merit further investigations regarding their field application and ecotoxicological studies.

Key words: Camphor, Garlic, Mustard, Radish, Rosemary, Thyme.

INTRODUCTION

Arthropod-borne diseases have increased all over the world, especially those transmitted by flies, sandflies, mosquitoes, and ticks (Marcondes 2017). The common housefly, *Musca domestica* (Diptera: Muscidae), is a cosmopolitan pest known since ancient civilizations (Khater 2017). Housefly is a major medical and veterinary important insect as it transmits mechanically over and above 100 pathogens including bacteria, fungi, viruses, and parasites (Khamesipour et al. 2018).

House flies are difficult to be managed because of their high fecundity, short developmental periods, and resistance to insecticides (Khan et al. 2013). Conventional

insecticides cannot be used in some places and circumstances.

Botanicals mainly essential oils (EOs) have been used daily by the ancient Egyptians (Khater 2017). Alternative parasiticides of botanical origin have long been used throughout history until recently (Seddiek et al. 2011, 2014; Khater 2017; Abbas et al. 2020; Ahmed et al. 2021; Iqbal et al. 2021; Štrbac et al. 2021). Essential oils are pleasing alternatives to conventional insecticides because of their ease of use, public acceptability, besides their low mammalian and avian toxicities. They could be used with suitability in organic production systems if oils are produced organically (Khater 2012a, 2012b, 2013).

Cite This Article as: Baz MM, Eltaly RI, Debboun M, Selim A, Radwan IT, Ahmed N and Khater HF, 2022. The contact/fumigant adulticidal effect of Egyptian oils against the house fly, *Musca domestica* (diptera: muscidae). International Journal of Veterinary Science x(x): xxxx. <https://doi.org/10.47278/journal.ijvs/2022.180>

Plant extracts and EOs are widely used for pest control as adulticides (Khater et al. 2014; Abdel-Meguid et al. 2022; Radwan et al. 2022a) and larvicides (Khater and Shalaby 2008; Khater et al. 2011, 2013; Murugan et al. 2015; Roni et al. 2015; Govindarajan et al. 2016a,b; Baz et al. 2021, 2022a,b; Radwan et al. 2022a,b). Botanical insecticides have also been used for controlling the *cycloraphan* flies (Khater 2014; Khater et al. 2009, 2013, 2018; Khater and Geden 2018, 2019).

There are many studies deal with the larvicidal effect of essential oils, but there are few studies about the adulticidal effect of few oils such as thyme (Pavela 2007) and garlic (Nisar et al. 2021). In order to determine the eco-friendly plant oils that are sustainable for housefly management, this investigation aimed to estimate the toxicity of six EOs including four novel oils against adult housefly through contact/fumigant toxicity bioassays, and calculate their lethal concentration and time values, as well as their relative toxicities and toxicity indices.

MATERIALS AND METHODS

Oils

Essential oils (Table 1) were purchased from *EL CAPTAIN Company* for extracting natural oils, plants and cosmetics "Cap Pharm", El Obor, Cairo, Egypt.

Housefly Colony

House fly colony was reared in the insectary in the Medical and of Molecular Entomology Section, Entomology Department, Faculty of Science, Benha University, Egypt. Houseflies were maintained at $27\pm 2^\circ\text{C}$ and 75-85% RH in $30\times 30\times 30$ cm cages. Adult house flies were fed on yeast and granular sucrose. Adult flies were allowed to oviposit eggs on cotton pads soaked in full fat milk. The diet of larval rearing comprised 1:1 vermiculite and oatmeal, and 30 cm^3 yeast extract, then moisturized with full fat milk (Khater and Geden 2019).

Contact/fumigant Toxicity

The efficacy of the applied EOs against adult *M. domestica*, was estimated according to a previous protocol (Khater and Geden 2019) with a little modification. Seven concentrations (0.03, 0.06, 0.13, 0.25, 0.5, 1.0, and 2.0%) were diluted in a solvent (Tween 20@, 5%, plus distilled water). Glass bottles and their caps were treated with a thin layer of each tested concentration. Bottles were turned around to disperse oils. Ten flies were added to each bottle and then covered with a cap. The solvent was used to treat the control group. Thirty flies (three replicates) were used for each treated group. Dead flies were counted 5, 10, 20, 30, and 60 min after placement in the bottles. Bottles were preserved ($27\pm 2^\circ\text{C}$ along with $80\pm 5\%$ RH).

Data Analysis

Probit analysis was used to analyze the mortality data through SPSS V23 (IBM, USA) to determine lethal concentration (LC) plus lethal time (LT) values. The mortalities were compared via One- Way Analysis of Variance, followed by the Tukey's test, where $P>0.05$ is considered not significant. The toxicity index (TI) (Zidan and Abdel-Mageed 1988), relative toxicity (RT) (Khater

and Geden 2018) and Time Potency (TP) were calculated as follows:

$\text{TI} = \text{LC}_{50}$ of the most toxic plant oil $\times 100 / \text{LC}_{50}$ of each tested plant oil.

$\text{RT} = \text{LC}_{50}$ of the least toxic plant oil / LC_{50} of each tested plant oil.

$\text{TP} = \text{LT}_{50}$ of the least toxic plant oil / LT_{50} of each tested plant oil.

RESULTS

The present study investigated the *in vitro* efficacy of six plant oils against *M. domestica* (Table 1). The study revealed that all tested oils exhibited adulticidal efficiency against *M. domestica*, depending on the concentration and time of exposure. The most effective oil was radish, followed by camphor, mustard, thyme, garlic, and rosemary. After treatment with 0.06%, the mortality (MO)% of camphor, garlic, mustard, radish, rosemary, and thyme oils were 43.33, 23.33, 30.00, 30.00, 36.66, and 40.00%, respectively, 30 min post-treatment (PT) and 50.00, 36.66, 43.33, 46.66, 43.33, and 46.66%, respectively, 60 min PT. On the other hand, the analogous values PT for 60 min with 2% were 90.00, 83.33, 86.66, 93.33, 83.33, and 86.66%, respectively (Table 2).

After treatment for 60 min, LC_{50} of camphor, garlic, mustard, radish, rosemary, and thyme were 0.305, 0.450, 0.366, 0.250, 0.475 and 0.405%, respectively. The corresponding LC_{99} values were 2.54, 3.19, 2.92, 2.30, 3.14 and 2.96%, respectively. On the other hand, their TIs were 81.97, 55.56, 68.31, 100.00, 52.63 and 61.73%, respectively (Table 3).

After treatment with concentration 0.5%, the median lethal time (LT_{50}) of camphor, garlic, mustard, radish, rosemary, and thyme were 26.04, 30.53, 26.63, 23.26, 29.45, and 28.62 min, respectively. The corresponding LT_{50} values PT with 2% were 16.52, 19.99, 18.29, 16.44, 18.73, and 17.92 min, respectively. The time potency PT with 0.5% of camphor, mustard, radish, rosemary, and thyme were 1.17, 1.15, 1.31, 1.04, and 1.07 times, respectively, faster than garlic, whereas such values PT with 2% were 1.21, 1.09, 1.22, 1.07, and 1.12, respectively (Table 4).

DISCUSSION

EOs are used as an eco-friendly alternative for pest management as they are non-mutagenic, non-toxic to animals and more effective against insecticide-resistant pests (Khater 2012a, b, 2013; Ahmed et al. 2021). The efficacy of EOs against *M. domestica* in this study indicated that all tested plant oils exhibited adulticidal efficiency against depending on the concentration and time of exposure.

Radish oil was the most effective adulticide followed by camphor, mustard, thyme, and garlic and they were 1.9, 1.55, 1.29, 1.17, and 1.05 times more effective than rosemary oil. Radish, *Raphanus sativus*, is a common vegetable oil reputed to possess diverse medicinal properties and all parts of the plant are used in medicines (Ahmad et al. 2012).

This study indicated radish, as its LC_{50} value and toxicity index were 0.250 and 100%, respectively. The insecticidal activity of different fractions of *R. sativus* leaf

Table 1: Binomial, English and family names of the applied oils against adult house fly

Binomial name	English name	Family name
<i>Allium sativum</i>	Garlic oil	Amaryllidaceae
<i>Brassica juncea</i>	Mustard oil	Brassicaceae
<i>Cinnamomum camphora</i>	Camphor	Lauraceae
<i>Raphanus sativus</i>	Radish fegl	Brassicaceae
<i>Salvia rosmarinus</i>	Rosemary	Lamiaceae
<i>Thymus vulgaris</i>	Thyme	Lamiaceae

Table 2: Effect of essential oils against the adult stage of *Musca domestica*

Oil name	Time	Mortality (%) Mean± SE							
		Control	0.03%	0.06%	0.13%	0.25%	0.5%	1.0%	2.0%
Camphor	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
	10 min	0.00±0.00c	(3.33)	(16.66)	(23.33)	(23.33)	(26.66)	(33.33)	(40.00)
		0.33±0.57c	1.66±0.57bc	2.33±0.57ab	2.33±0.57ab	2.66±0.57ab	3.33±0.57ab	4.00±1.00a	
	20 min	(3.33)	(13.33)	(26.66)	(33.33)	(36.66)	(43.33)	(50.00)	(60.00)
		0.33±0.57d	1.33±0.57cd	2.66±1.15bcd	3.33±1.15bc	3.66±0.57abc	4.33±0.57ab	5.00±1.00ab	6.00±1.00a
	30 min	(3.33)	(26.66)	(43.33)	(50.00)	(53.33)	(56.66)	(66.66)	(76.66)
0.33±0.57e		2.66±1.15d	4.33±1.15cd	5.00±0.00bc	5.33±0.57bc	5.66±0.57abc	6.66±0.57ab	7.66±0.57a	
60 min	(6.66)	(30.00)	(50.00)	(60.00)	(63.33)	(73.33)	(80.00)	(90.00)	
	0.66±0.57e	3.00±1.73de	5.00±1.73cd	6.00±1.00bc	6.33±0.57abc	7.33±0.57abc	8.00±0.00ab	9.00±1.00a	
Garlic	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	
	10 min	0.00±0.00c	(3.33)	(6.66)	(13.33)	(26.66)	(26.66)	(30.00)	(36.66)
		0.33±0.57c	0.66±0.57c	1.33±0.57bc	2.66±0.57ab	2.66±0.57ab	3.00±0.00a	3.66±0.57a	
	20 min	(3.33)	(13.33)	(13.33)	(20.00)	(30.00)	3.33±0.57abc	(40.00)	(50.00)
		0.33±0.57e	1.33±0.57de	1.33±0.57de	2.00±1.00cde	3.00±0.00bcd	4.00±0.00ab	5.00±1.00a	
	30 min	(3.33)	(16.66)	(23.33)	(36.66)	(43.33)	(50.00)	5.66±0.57ab	6.66±0.57a
0.33±0.57f		1.66±0.57ef	2.33±0.57de	3.66±0.57cd	4.33±1.54bc	5.00±0.00abc			
60 min	(6.66)	(26.66)	(36.66)	(53.33)	(63.33)	(70.00)	(70.00)	(83.33)	
	0.66±0.57e	2.66±0.57d	3.66±0.57cd	5.33±0.57bc	6.33±0.57b	7.00±1.00ab	7.00±0.00ab	8.33±0.57a	
Mustard	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	
	10 min	0.00±0.00c	(6.66)	(10.00)	(20.00)	(20.00)	(26.66)	(33.33)	(40.00)
		0.66±0.57bc	1.00±1.00bc	2.00±1.00bc	2.00±1.00abc	2.00±0.00abc	2.66±1.15ab	3.33±0.57	4.00±1.00a
	20 min	(3.33)	(13.33)	(20.00)	(30.00)	(33.33)	(40.00)	(50.00)	(53.33)
		0.33±0.57e	1.33±0.57de	2.00±0.00cde	3.00±1.00bcd	3.33±0.57bcd	4.00±1.00abc	5.00±1.00ab	5.33±0.57a
	30 min	(3.33)	(23.33)	(30.00)	(43.33)	(50.00)	(56.66)	(63.33)	(70.00)
0.33±0.57e		2.33±1.52de	3.00±0.00cd	4.33±0.57bcd	5.00±0.00abc	5.66±0.57ab	6.33±0.57ab	7.00±1.00a	
60 min	(6.66)	(30.00)	(43.33)	(53.33)	(63.33)	(73.33)	(73.33)	(86.66)	
	0.66±0.57e	3.00±1.00de	4.33±0.57cd	5.33±0.57bcd	6.33±1.52abc	7.33±1.15ab	7.33±0.57ab	8.66±0.57a	
Radish	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	
	10 min	0.00±0.00d	(10.00)	(10.00)	(23.33)	(33.33)	(33.33)	(36.66)	(43.33)
		1.00±1.00cd	1.00±1.00cd	2.33±0.57bc	3.33±0.57ab	3.33±0.57ab	3.66±0.57ab	4.33±0.57a	
	20 min	(3.33)	(13.33)	(20.00)	(30.00)	(43.33)	(46.66)	(50.00)	(56.66)
		0.33±0.57d	1.33±0.57cd	2.00±1.00cd	3.00±1.00bc	4.33±0.57ab	4.66±0.57ab	5.00±1.00ab	5.66±0.57a
	30 min	(3.33)	(23.33)	(30.00)	(43.33)	(56.66)	(63.33)	(70.00)	(73.33)
0.33±0.57d		2.33±0.57cd	3.00±1.73c	4.33±1.15bc	5.66±0.57ab	6.33±0.57ab	7.00±1.00a	7.33±0.57a	
60 min	(6.66)	(33.33)	(46.66)	(60.00)	(70.00)	(73.33)	(80.00)	(93.33)	
	0.66±0.57e	3.33±1.52d	4.66±1.15cd	6.00±0.00bc	7.00±1.00abc	7.33±0.57ab	8.00±0.00ab	9.33±0.57a	
Rosemary	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	
	10 min	0.00±0.00d	(3.33)	(13.33)	(20.00)	(23.33)	(26.66)	(30.00)	(40.00)
		0.33±0.57cd	1.33±0.57bcd	2.00±1.00bc	2.33±0.57ab	2.66±0.00ab	3.00±1.00ab	4.00±1.00a	
	20 min	(3.33)	(10.00)	(23.33)	(30.00)	(33.33)	(40.00)	(46.66)	(53.33)
		0.33±0.57e	1.00±0.0de	2.33±0.57cd	3.00±1.007bc	3.33±0.57bc	4.00±1.00abc	4.66±0.57ab	5.33±0.57a
	30 min	(3.33)	(23.33)	(36.66)	(43.33)	(46.66)	(53.33)	(60.00)	(70.00)
0.33±0.57e		2.33±0.57de	3.66±0.57cd	4.33±0.57bcd	4.66±0.57bc	5.33±1.15abc	6.00±1.00ab	7.00±1.00a	
60 min	(6.66)	(30.00)	(43.33)	(50.00)	(56.66)	(66.66)	(73.33)	(83.33)	
	0.66±0.57f	3.00±1.00e	4.33±0.57de	5.00±1.00cd	5.66±0.57bcd	6.66±0.57abc	7.33±0.57ab	8.33±0.57a	
Thyme	5 min	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	
	10 min	0.00±0.00c	(3.33)	(16.66)	(23.33)	(23.33)	(26.66)	(33.33)	(40.00)
		0.33±0.57c	1.66±0.57bc	2.33±0.57ab	2.33±0.57ab	2.66±0.57ab	3.33±0.57ab	4.00±1.00a	
	20 min	(3.33)	(13.33)	(26.66)	(33.33)	(36.66)	(40.00)	(46.66)	(56.66)
		0.33±0.57d	1.33±0.57cd	2.66±1.15bcd	3.33±1.15abc	3.66±0.57abc	4.00±1.00ab	4.66±1.15ab	5.66±0.57a
	30 min	(3.33)	(23.33)	(40.00)	(46.66)	(50.00)	(56.66)	(63.33)	(70.00)
0.33±0.57d		2.33±0.57cd	4.00±1.00bc	4.66±1.52abc	5.00±1.00ab	5.66±0.57ab	6.33±0.57ab	7.00±1.00a	
60 min	(6.66)	(26.66)	(46.66)	(60.00)	(56.66)	(66.66)	(73.33)	(86.66)	
	0.66±0.57e	2.66±1.15de	4.66±1.15cd	6.00±0.00bc	5.66±0.57bc	6.66±1.15abc	7.33±0.57ab	8.66±0.57a	

Means followed by the same letter on the same row are not significantly different by ANOVA (P>0.05).

Table 3: Lethal concentration values of oils after treatment of adult *Musca domestica* for one hour

Essential oil	LC50	LC90	LC95	LC99	Relative toxicity	Toxicity index	Chi df	Equation R2
	Upper Lower	Upper Lower	Upper Lower	Upper Lower				
Camphor	0.305	1.54	1.89	2.54	1.55	81.97	30.95	Y=0.05+0.75xX 0.794
	-1.42	0.87	1.07	1.43				
Garlic	1.31	18.64	23.99	34.07	1.05	55.56	30.72	Y=0.15+0.63xX 0.678
	-1.95	1.095	1.33	1.761				
Mustard	4.80	104.09	132.87	186.86	1.29	68.31	30.23	Y=0.08+0.66xX 0.746
	-1.85	0.998	1.22	1.63				
Radish	1.92	35.56	45.68	64.68	1.9	100.00	31.923	Y=6.8E-3+0.8xX 0.828
	-2.47	0.77	0.96	1.29				
Rosemary	1.16	26.65	34.62	49.57	1.00	52.63	24.896	Y=0.18+0.66 0.811
	-0.35	1.148	1.39	1.83				
Thyme	1.63	11.83	14.90	20.68	1.17	61.73	27.860	Y=0.13+0.68xX 0.783
	-0.80	1.04	1.27	1.69				
	1.67	17.38	22.12	31.04			6	0.000 ^a

Relative toxicity = LC50 of the least toxic compound / LC50 of the tested compound: Toxicity index = LC50 of the most toxic compound \times 100 / LC50 of the tested compound: The most toxic oil has given 100 units on the toxicity index scale.

Table 4: Median lethal time values and time potency of essential oils against *Musca domestica* adults

Essential oil	Concentration 0.5%			Concentration 2%		
	LT ₅₀ (Lower-upper)	LT ₉₅ (Lower-upper)	Time potency	LT ₅₀ (Lower-upper)	LT ₉₅ (Lower-upper)	Time potency
Camphor	26.04	152.21	1.17	16.52	66.46	1.21
	20.42	89.94		13.28	47.47	
	34.76	410.66		20.24	116.05	
Garlic	30.53	199.48	1.00	19.99	99.99	1.00
	23.56	108.61		15.85	65.35	
	43.174	666.83		25.30	211.72	
Mustard	26.63	154.79	1.15	18.29	85.36	1.09
	20.89	91.21		14.55	57.70	
	35.67	420.48		22.85	167.71	
Radish	23.26	145.82	1.31	16.44	65.04	1.22
	18.08	85.82		13.26	46.54	
	30.84	397.01		20.12	113.46	
Rosemary	29.45	207.99	1.04	18.73	94.98	1.07
	22.57	110.71		14.77	62.40	
	41.84	739.37		23.64	198.74	
Thyme	28.62	196.80	1.07	17.92	83.34	1.12
	22.02	106.75		14.24	56.57	
	40.12	664.00		22.35	162.26	

Times potency = LT₅₀ of the least toxic plant oil / LT₅₀ of each tested plant oil: LT₅₀ of the lowest toxic oil was Garlic oil.

extracts against 3rd nymphal instars of *Phenacoccus solenopsis* was recorded (Ibrahim and Abdel-Mogib 2019). The insecticidal effect of fractions of radish root extracts against adults and nymphs of *Aphis gossypii* could be arranged as follows; methylene chloride with LC₅₀= 386.63 and 309.43 ppm, ethyl acetate with LC₅₀= 394.9 and 334.37 ppm, and petroleum ether with LC₅₀= 636.2 and 424.56 ppm, respectively (Ibrahim et al. 2020).

Mustard oil was effective as adulticide in this work (LC₅₀= 0.366%). Similar insecticidal effect was recorded as radish and mustard (*Brassica campestris*) oils induced larvicidal effects against the third larval instar of the blowfly, *Lucilia sericata* (LC₅₀= 6.93 and 7.92%, respectively) and hindered adult emergence PT with 12 and 8%, respectively (Khater and Khater 2009). Moreover, mustard (*Brassica juncea*) and arugula (*Eruca sativa*) plants repelled the sweet potato whitefly, *Bemisia tabaci* (Legaspi et al. 2016).

Garlic, *Allium sativum*, is a well-known herbal remedy of various health benefits (Tsai et al. 2012; Seddiek et al. 2014). This study revealed the multidirectional efficacy of garlic oil against house fly (LC₅₀= 0.450 PT for one hour). Parallel to our results, garlic juice was effective against the dipteran insects, *Delia radicum* and *M. domestica*; the recorded LC₅₀ values for *M. domestica* were 10.1 and 2.2% PT of larvae and adults for 24 h (Prowse et al. 2006). Aqueous garlic extract induced larvicidal effect and adversely affected oviposition, egg hatching, and fecundity of house flies (Nisar et al. 2021). Furthermore, complete larval mortalities of the oestrid fly, *Cephalopina titillator* (Clark); was reached 24h PT with 7.5% garlic oil (LC₅₀= 0.44%) (Khater 2014).

In the current work, camphor, *Cinnamomum camphora*, oil effectively controlled adult house flies (LC₅₀= 0.305%). Analogous studies showed that camphor oil was effective against *C. titillator* (Clark); the mortality

of 2nd larval instars PT for 24h with lavender, camphor, and onion oils was 100, 80, and 52%, respectively, whereas those of 3rd larval instars were 100, 68, and 52%, respectively. Furthermore, the LT₅₀, LT₉₀, and LT₉₉ values PT of 2nd larval instars were 3.60, 12.06, and 32.31 h for lavender oil; 6.58, 51.67, and 277.38 h for camphor oil; and 14.24, 120.73, and 689.66 h for onion oils, respectively (Khater et al. 2013). Camphor oil was also effective against the slender pigeon louse, *Columbicola columbae* and *in vitro* treated lice with 1% were died one hour PT (LC₅₀= 0.25%, and LT₅₀= 2.30 min PT with 0.004%). The louse infestations were eliminated seven days PT of infested pigeons (Khater et al. 2014).

This work demonstrated that the least toxic oil was rosemary and its toxicity index was 52.63%. Similarly, rosemary oil was less effective against the predacious mite, *Amblyseius zaheri*, but it was the most toxic to *Amblyseius barkeri* females (Momen and Amer 2000). Moderate efficacy of rosemary was also reported against *Culex pipiens* mosquito, rosemary oil induced a larvicidal effect (LC₅₀=71.37ppm) and adversely affected its pupation rates and adult emergences (Khater and Shalaby 2008); hexane extract of rosemary induced 100% mortality PT of 3rd and 4th instars larvae with 160 ppm (Shalaby and Khater 2005). Furthermore, rosemary oil was effective against *L. sericata* (LC₅₀=6.77%) and sublethal concentrations induced morphological abnormalities plus a dominance of males over females (4:1) which could lead to a population decline (Khater et al. 2011).

A comparable study recorded the repellent, lousicidal, and ovicidal effect of Egyptian EOs against flies and the buffalo louse, *Haematopinus tuberculatus*; four minutes PT of *H. tuberculatus*, *in vitro*, LC₅₀ values were 2.74, 22.79, 18.67, 7.28, and 12.35% for camphor, rosemary, chamomile, onion, and peppermint oils, respectively (LT₅₀= 0.89, 11.60, 21.32, 2.75, and 15.39 min, respectively) after treatment with 7.5%; *In vivo* treatments indicated complete mortality of lice 0.5–2 min and the number of lice infesting buffaloes was significantly reduced 3, 4, 6, and 6 days PT with camphor, chamomile, peppermint, and onion oils, respectively. Moreover, the oils and *d*-phenothrin significantly repelled flies, *M. domestica*, *Haematobia irritans*, *Hippobosca equine*, and *Stomoxys calcitrans* for 6 and 3 days, respectively, PT of infested buffaloes (Khater et al. 2009).

This study indicated that thyme (*Thymus vulgaris*) oil was an effective adulticide (LC₅₀= 0.305%). In agreement, a study indicated the lethal effects of thyme oil against *M. domestica* larvae and adults with 80% MO plus decreased longevity of both sexes (Pavela 2007). Thyme EO was highly effective against *M. domestica* larvae through residual film method followed by clove then basil oils (LC₅₀= 2.82, 3.79, and 5.99 mg/ml, respectively) (Chere et al. 2018).

Against *M. domestica*, findings of this study come along with other studies about using other botanicals against *M. domestica*; higher larvicidal and pupicidal efficacy of *Mentha piperita* than *Mentha citrata* EO in contact and fumigation toxicity assays was recorded (Kumar et al. 2012) and the EOs of mint and lavender have a larvicidal and pupicidal effect (Bosly 2013).

Parallel studies against *M. domestica* revealed that LC₅₀ values of 12 EOs and 17 terpenes ranged from 3.9 to 85.2 and from 3.3 to >100mg/dm³, respectively. *Citrus sinensis* EO was the most effective one (LC₅₀= 3.9 mg/dm³), then EOs of *Citrus aurantium* (LC₅₀= 4.8 mg/dm³) and *Eucalyptus cinerea* (LC₅₀= 5.5 mg/dm³). Moreover, 1, 8-cineole (56.86%), a major constituent of *E. cinerea* EO, was the most effective one (LC₅₀= 3.3 mg/dm³) (Palacios et al. 2009). Eucalyptus oil had repellent and toxic effect against house flies (Kumar et al. 2011).

Similar to the findings of this study, a corresponding study indicated toxic and repellent efficacy of vetiver, cinnamon, and lavender EOs and their blends against larvae and adult *M. domestica*. Such oils were moderately toxic for eggs. Mortality of 2nd instar larvae reached 94–100% in treated media, 57–78% in dipping assays, and 38–100% in contact assays. Contact/fumigant toxicity of lavender, vetiver, cinnamon, and sunflower oils was 100, 100, 100, and 67%, respectively. Against adults, oil blends were not more effective than the individual oils; nonetheless, diluted blends with sunflower oil were equally effective as the individual oils. Moreover, cinnamon and vetiver EOs were strong larval repellents (78 and 84%, respectively) in treated media, but all oils through olfactometer assays did not repel adult flies, whereas evaluating more materials against flies demonstrated significant repellency for neem oil, vanillin, and p-menthane-3,8-diol (PMD) (Khater and Geden 2019).

Besides their insecticidal effects, EOs prolonged larval and pupal durations, reduced pupation and adult emergency% and induced morphological anomalies of the developmental stages post treatment of the third larval instar of *M. domestica* (Khater 2003; Bosly 2013).

Conclusion

House fly is challenging pest and controlling adult flies would suppress the population shortly. In this work, the adulticidal outcome of radish, mustard, rosemary and camphor against house flies had a novelty record and the current investigation concluded that all applied oils were effective adulticides against housefly. Radish oil was the most potent oil followed by camphor, mustard, and thyme oils, whereas garlic and rosemary were less effective candidates; thus, the promising oils merit further investigations regarding their persistence, field application, and ecotoxicological studies.

Acknowledgments

The authors would like to thank the funding agency, the Science, Technology, and Innovation Funding Authority, Egypt [Grant number: 41608] for funding this work, project title: “Eco-friendly Pesticides against Pests of Medical, Veterinary, and Agricultural Importance.” The authors would like to thank Prof. Dr. Azza A. Moustafa, Research Institute of Medical Entomology, Egypt, for her valuable support and advice.

Funding

The Science, Technology, and Innovation Funding Authority, Egypt, via Grant number 41608, supported this work. Project title: “Eco-friendly Pesticides against Pests of Medical, Veterinary, and Agricultural Importance.”

Conflict of interest

The authors declare that there is no conflict of interest.

Authors' contribution

MB, HK: Helped with lab work and writing; RE, MD: Helped with writing; AS, NA: Helped with referencing and editing manuscript.

REFERENCES

- Abbas RZ, Zaman MA, Sindhu D, Sharif M, Rafique A, Saeed Z, Siddique F, Zaheer T, Khan MK and Akram MS, 2020. Anthelmintic effects and toxicity analysis of herbal dewormer against the infection of *Haemonchus contortus* and *Fasciola hepatica* in goat. Pakistan Veterinary Journal 40: 455-460. <https://doi.org/10.29261/pakvetj/2020.083>
- Abdel-Meguid AD, Ramadan MY, Khater HF and Radwan IT, 2022. Lousicidal efficacy of essential oils against the dog louse, *Trichodectes canis* (Mallophaga: Trichodectidae). Egyptian Academic Journal of Biological Sciences, Medical Entomology & Parasitology 14: 1-16.
- Ahmad F, Hasan I, Chishti DK and Ahmad H, 2012. Antibacterial activity of *Raphanus sativus* Linn. Seed extract. Global Journal of Medical Research 12(11): 25-34.
- Ahmed N, Alam M, Saeed M, Ullah H, Iqbal T, Al-Mutairi KA, Shahjeer K, Ullah R, Ahmed S, Ahmed NaaH, Khater H and Salman M, 2021. Botanical insecticides are a non-toxic alternative to conventional pesticides in the control of insects and pests. In: Global Decline of Insects. El-Shafie H (ed), IntechOpen, London, UK. <https://doi.org/10.5772/intechopen.100416>
- Baz MM, Hegazy MM, Khater HF and El-Sayed YA, 2021. Comparative Evaluation of Five Oil-Resin Plant Extracts against The Mosquito Larvae, *Culex pipiens* Say (Diptera: Culicidae). Pakistan Veterinary Journal 41: 191-196. <http://dx.doi.org/10.29261/pakvetj/2021.010>
- Baz M, Selim A, Radawn I and Khater H, 2022a. Plant oils in the fight against the West Nile Vector, *Culex pipiens*. International Journal of Tropical Insect Science. <https://doi.org/10.1007/s42690-022-00762-1>
- Baz MM, Selim A, Radwan IT, Alkhaibari AM and Khater HF, 2022b. Larvicidal and Adulticidal effects of Some Egyptian oils against *Culex pipiens*. Scientific Reports 12: 1-18. <https://doi.org/10.1038/s41598-022-08223-y>
- Bosly H, 2013. Evaluation of insecticidal activities of *Mentha piperita* and *Lavandula angustifolia* essential oils against house fly, *Musca domestica* L.(Diptera: Muscidae). Journal of Entomology and Nematology 5: 50-54. <https://doi.org/10.5897/JEN2013.0073>
- Chere JMC, Dar MA and Pandit RS, 2018. Evaluation of some essential oils against the larvae of house fly, *Musca domestica* by using residual film method. Advances in Biotechnology & Microbiology 9: 555752. <https://doi.org/10.19080/AIBM.2018.09.555752>
- Govindarajan M, Khater HF, Panneerselvam C and Benelli G, 2016a. One-pot fabrication of silver nanocrystals using *Nicandra physalodes*: A novel route for mosquito vector control with moderate toxicity on non-target water bugs. Research in Veterinary Science 107: 95-101. <https://doi.org/10.1016/j.rvsc.2016.05.017>
- Govindarajan M, Rajeswary M, Muthukumar U, Hoti S, Khater HF and Benelli G, 2016b. Single-step biosynthesis and characterization of silver nanoparticles using *Zornia diphylla* leaves: A potent eco-friendly tool against malaria and arbovirus vectors. Journal of Photochemistry and Photobiology B: Biology 161: 482-489. <https://doi.org/10.1016/j.jphotobiol.2016.06.016>
- Ibrahim HY and Abdel-Mogib M, 2019. Chemical constituents of radish, *Raphanus sativus* linn. (brassicaceae) leaves extracts and their insecticidal activity against mealybug, *solenopsis tinsley* (hemiptera: pseudococcidae). International Journal of Current Advanced Research 8(10): 20179-20184. <http://dx.doi.org/10.24327/ijcar.2019.20184.3934>
- Ibrahim HY, Abdel-Mogib M and Mostafa ME, 2020. Insecticidal activity of radish, *Raphanus sativus* Linn. (Brassicaceae) roots extracts. Journal of Plant Protection and Pathology 11: 53-58. <https://doi.org/10.21608/jppp.2020.79164>
- Iqbal T, Ahmed N, Shahjeer K, Ahmed S, Al-Mutairi KA, Khater HF and Ali RF, 2021. Botanical insecticides and their potential as anti-insect/pests: Are they successful against insects and pests? In: Global Decline of Insects. El-Shafie H (ed), IntechOpen, London, UK <https://doi.org/10.5772/intechopen.100418>.
- Khamesipour F, Lankarani KB, Honarvar B and Kwenti TE, 2018. A systematic review of human pathogens carried by the housefly (*Musca domestica* L.). BMC Public Health 18: 1-15.
- Khan HA, Shad SA and Akram W, 2013. Resistance to new chemical insecticides in the house fly, *Musca domestica* L., from dairies in Punjab, Pakistan. Parasitology Research 112: 2049-2054. <https://doi.org/10.1007/s00436-013-3365-8>
- Khater HF, 2003. Biocontrol of some insects. Parasitology. Egypt: Benha University.
- Khater HF, 2012a. Prospects of botanical biopesticides in insect pest management. Pharmacologia, 3: 641-656. <https://doi.org/10.5567/Pharmacologia.2012.641.656>
- Khater HF, 2012b. Ecosmart biorational insecticides: alternative insect control strategies. In: Insecticides-Advances in Integrated Pest Management. Preveen F (ed), Rijeka, Croatia, pp.17-61. <https://doi.org/10.5772/27852>
- Khater HF, 2013. Bioactivity of essential oils as green biopesticides: recent global scenario. Recent progress in medicinal plants. In: Essentials Oils II, Govil JN and Bhattacharya S (eds), Vol. 37; Studium Press LLC, USA, pp. 153-220.
- Khater HF, 2014. Bioactivities of some essential oils against the camel nasal botfly, *Cephalopina titillator*. Parasitology Research 113: 593-605. <https://doi.org/10.1007/s00436-013-3688-5>
- Khater H, 2017. Introductory Chapter: Back to the Future - Solutions for Parasitic Problems as Old as the Pyramids. In: Natural remedies in the fight against parasites. Khater HF, Govindarajan M, Benelli G (eds). IntechOpen, London, pp 4-19. <https://doi.org/10.5772/67554>.
- Khater HF and Shalaby AA-S, 2008. Potential of biologically active plant oils to control mosquito larvae (*Culex pipiens*, Diptera: Culicidae) from an Egyptian locality. Revista do Instituto de Medicina Tropical de São Paulo 50 :107-112. <https://doi.org/10.1590/S0036-46652008000200008>
- Khater HF and Khater D, 2009. The insecticidal activity of four medicinal plants against the blowfly *Lucilia sericata* (Diptera: Calliphoridae). International Journal of Dermatology 48: 492-497. <https://doi.org/10.1111/j.1365-4632.2009.03937.x>
- Khater HF and Geden CJ, 2018. Potential of essential oils to prevent fly strike and their effects on the longevity of adult *Lucilia sericata*. Journal of Vector Ecology 43: 261-270. <https://doi.org/10.1111/jvec.12310>
- Khater HF and Geden CJ, 2019. Efficacy and repellency of some essential oils and their blends against larval and adult house flies, *Musca domestica* L.(Diptera: Muscidae). Journal of Vector Ecology 44: 256-263. <https://doi.org/10.1111/jvec.12357>
- Khater HF, Ramadan MY and El-Madawy RS, 2009. Lousicidal, ovidical and repellent efficacy of some essential oils against lice and flies infesting water buffaloes in Egypt. Veterinary Parasitology 164: 257-266. <https://doi.org/10.1016/j.vetpar.2009.06.011>

- Khater HF, Hanafy A, Abdel-Mageed AD, Ramadan MY and El-Madawy RS, 2011. Control of the myiasis-producing fly, *Lucilia sericata*, with Egyptian essential oils. *International Journal of Dermatology* 50: 187-194. <https://doi.org/10.1111/j.1365-4632.2010.04656.x>
- Khater HF, Ramadan MY and Magedi ADA, 2013. In vitro control of the camel nasal botfly, *Cephalopina titillator*, with doramectin, lavender, camphor, and onion oils. *Parasitology Research* 112: 2503-2510. <https://doi.org/10.1007/s00436-013-3415-2>
- Khater HF, El-Shorbagy MM and Seddiek SA, 2014. Lousicidal efficacy of camphor oil, d-phenothrin, and deltamethrin against the slender pigeon louse, *Columbicola columbae*. *International Journal of Veterinary Science and Medicine* 2: 7-13. <https://doi.org/10.1016/j.ijvsm.2013.12.003>
- Khater HF, Ali AM, Abouelella GA, Marawan MA, Govindarajan M, Murugan K, Abbas RZ, Vaz NP and Benelli G, 2018. Toxicity and growth inhibition potential of vetiver, cinnamon, and lavender essential oils and their blends against larvae of the sheep blowfly, *Lucilia sericata*. *International Journal of Dermatology* 57: 449-457. <https://doi.org/10.1111/ijd.13828>
- Kumar P, Mishra S, Malik A and Satya S, 2012. Efficacy of *Mentha piperita* and *Mentha citrata* essential oils against housefly, *Musca domestica* L. *Industrial Crops and Products* 39: 106-112. <https://doi.org/10.1016/j.indcrop.2012.02.021>
- Kumar P, Mishra S, Malik A and Satya S, 2011. Repellent, larvicidal and pupicidal properties of essential oils and their formulations against the housefly, *Musca domestica*. *Medical and Veterinary Entomology* 25(3): 302-310 <https://doi.org/10.1111/j.1365-2915.2011.00945.x>
- Legaspi JC Miller N, Wolaver D, Haseeb M and Zaniccio J, 2016. Repellency of mustard (*Brassica juncea*) and arugula (*Eruca sativa*) plants, and plant oils against the sweetpotato whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Subtropical Agriculture and Environments* 67: 28-34.
- Marcondes CB, 2017. *Arthropod Borne Diseases*. Springer, Cham, Switzerland.
- Momen F and Amer S, 2000. Effect of rosemary and sweet marjoram on three predacious mites of the family Phytoseiidae (Acari: Phytoseiidae). *Acta Phytopathologica et Entomologica Hungarica* 34: 355-361. <https://doi.org/10.1556/aphyt.34.1999.4.11>
- Murugan K, Priyanka V, Dinesh D, Madhiyazhagan P, Panneerselvam C, Subramaniam J, Suresh U, Chandramohan B, Roni M, Nicoletti M, Alarfaj A, Higuchi A, Munusamy M, Khater H, Messing R and Giovanni B, 2015. Predation by Asian bullfrog tadpoles, *Hoplobatrachus tigerinus*, against the dengue vector, *Aedes aegypti*, in an aquatic environment treated with mosquitoicidal nanoparticles. *Parasitology Research* 114: 3601-3610. <https://doi.org/10.1007/s00436-015-4582-0>
- Nisar MS, Ismail MA, Ramzan H, Maqbool MM, Ahmad T, Ghramh HA, Khalofah A, Kmet J, Horvát M and Farooq S, 2021. The impact of different plant extracts on biological parameters of Housefly [*Musca domestica* (Diptera: Muscidae)]: Implications for management. *Saudi Journal of Biological Sciences* 28: 3880-3885. <https://doi.org/10.1016/j.sjbs.2021.03.070>
- Palacios SM, Bertoni A, Rossi Y, Santander R and Urzúa A, 2009. Efficacy of essential oils from edible plants as insecticides against the house fly, *Musca Domestica* L. *Molecules* 14: 1938-1947. [doi:https://doi.org/10.3390/molecules14051938](https://doi.org/10.3390/molecules14051938)
- Pavela R, 2007. Lethal and sublethal effects of thyme oil (*Thymus vulgaris* L.) on the house fly (*Musca domestica* Lin.). *Journal of Essential Oil Bearing Plants* 10: 346-356. <https://doi.org/10.1080/0972060X.2007.10643566>
- Prowse GM, Galloway TS and Foggo A, 2006. Insecticidal activity of garlic juice in two dipteran pests. *Agricultural and Forest Entomology* 8: 1-6. <https://doi.org/10.1111/j.1461-9555.2006.00273.x>
- Radwan IT, Baz MM, Khater H and Selim AM, 2022a. Nanostructured lipid carriers (NLC) for biologically active green tea and fennel natural oils delivery: Larvicidal and adulticidal activities against *Culex pipiens*. *Molecules* 27: 1939. <https://doi.org/10.3390/molecules27061939>
- Radwan IT, Baz MM, Khater H, Alkhaibari A, and Selim AM, 2022b. Mg-LDH nanoclays intercalated fennel and green tea active ingredient: Field and laboratory evaluation of insecticidal activities against *Culex pipiens* and their non-target organisms. *Molecules* 27: 2424. <https://doi.org/10.3390/molecules27082424>
- Roni M, Murugan K, Panneerselvam C, Subramaniam J, Nicoletti M, Madhiyazhagan P, Dinesh D, Suresh U, Khater HF and Wei H, 2015. Characterization and biotoxicity of *Hypnea musciformis*-synthesized silver nanoparticles as potential eco-friendly control tool against *Aedes aegypti* and *Plutella xylostella*. *Ecotoxicology and Environmental Safety* 121: 31-38.
- Seddiek SA, Ali MM, Khater HF and El-Shorbagy MM, 2011. Anthelmintic activity of the white wormwood, *Artemisia herba-alba* against *Heterakis gallinarum* infecting turkey poults. *Journal of Medicinal Plants Research* 5(16): 3946-3957.
- Seddiek SA, El-Shorbagy MM, Khater HF and Ali AM, 2014. The antitrichomonal efficacy of garlic and metronidazole against *Trichomonas gallinae* infecting domestic pigeons. *Parasitology Research* 113: 1319-1329.
- Shalaby A and Khater H, 2005. Toxicity of certain solvent extracts of *Rosmarinus officinalis* against *Culex pipiens* larvae. *Journal of the Egyptian-German Society of Zoology* E 48: 69-80.
- Štrbac F, Bosco A, Amadesi A, Rinaldi L, Stojanović D, Simin N, Orčić D, Pušić I, Krnjajić S and Ratajac R, 2021. Ovicidal potential of five different essential oils to control gastrointestinal nematodes of sheep. *Pakistan Veterinary Journal* 41: 353-358. <http://dx.doi.org/10.29261/pakvetj/2021.026>
- Tsai CW, Chen HW, Sheen LY and Lii CK, 2012. Garlic: Health benefits and actions. *Biomedicine* 2: 17-29. <https://doi.org/10.1016/j.biomed.2011.12.002>
- Zidan Z and Abdel-Megeed M, 1988. *New approaches in pesticides and insect control*. Arabian Publishing House and Delivery, Cairo, Egypt.