



Impact of *Curcuma longa* L. on Semen and Blood Parameters in Awassi Lambs

Khalid H Sultan

University of Mosul, College of Agriculture and Forestry, Animal Production Department, Iraq

*Corresponding author: dr.khalid.h@uomosul.edu.iq

Article History: 22-660

Received: 06-Jul-22

Revised: 16-Jul-22

Accepted: 20-Jul-22

ABSTRACT

The current study was conducted in a private field in Bibukht village, Mosul government, in order to evaluate the impact of *Curcuma longa* powder (CL) supplementation to the male Awassi ration on semen quality and some blood parameters. Twenty-four male Awassi lambs, with average body weight 47 ± 1.23 kg and 10-11 months old, were randomly assigned to two equal groups (12 lamb/group). The first group (control) was reared on a standard ration, whereas the second group was reared on a standard ration supplemented with CL (200mg/kg ration). The treatments lasted for 90 days. Semen and blood parameters were taken at 30, 60, and 90 days of treatment. Statistical analysis for the data revealed that CL supplementation caused a significant increase in ejaculate volume, concentration/ejaculate, concentration/mL, live sperm%, and individual motility compared with control, especially at the end of treatment (90 days), and a significant decrease in sperm abnormality% compared with control at $P < 0.05$. On the other hand, CL treatment increased testosterone values and reduces estrogen values significantly as compared with control. In terms of blood parameters, CL treatment significantly improved most of the blood picture indices when compared to control, as well as serum total protein and globulin values when compared to control. In conclusion, CL addition to male Awassi lambs ration improves testosterone values and sexual activity as shown by the improvement of serum quality.

Key words: Awassi, *Curcuma longa*, Blood parameters, Semen, Sexual activity.

INTRODUCTION

One of the feeding methods used in recent years is enhanced ruminant adaptability and yield items, as phytogetic additions have so been common. Consequently, using natural products has a positive impact influence on people's health (Karásková et al. 2015). After the dangers of their use on animals, a number of chemical preparations such as growth stimulants, antibiotics, or hormones were used to enhance the growth and, as a result, they affect human health when consumed the product of these animals (Schwarz et al. 2001), as these chemical preparations promote bacterium resistance in humans ingesting those animals' goods (Hameed et al. 2020). Benkő et al. (2008) used some additives, such as probiotics, on some reproductive parameters and the hormonal status of female quail breeders, likewise, Abdul-Majeed et al. (2021) used nettle plant to improve some physiological and biochemical parameters in broiler chickens.

On the other hand, according to Molosse et al. (2019), adding curcumin to the diet of nursing lambs caused an increase in weight gain as well as anti-inflammatory and antioxidant effects, both of which have biological benefits for improving performance. Wencelová et al. (2015) recorded that medicinal herbs, as well as extracts

containing essential oils, polyphenols, saponins, flavonoids and other secondary metabolites, can help to improve ruminal fermentation and metabolism.

As a result, feeding ruminants with herbal plants has become increasingly popular. However, the utilization of plants with nutraceutical qualities as synthetic promoters of growth replacements (García-Hernández et al. 2017) must be researched (FrAnKIC et al. 2009; Khan et al. 2016). Furthermore, every feed plant additive should be examined due to the composition and secondary metabolites, varied action mechanisms, and the optimum dose for each physiological situation (Mendel et al. 2017). When using high-performance liquid chromatography, two additional curcuminoid components, demethoxycurcumin and bisdemethoxycurcumin, were found in very tiny quantities (1.62 and 0.98 %, respectively) when following the method of Coradini et al. (2014), which is discussed in detail by Jaguezski et al. (2018). However, there is conflicting evidence about the effects of curcumin on male fertility indices. Several in vivo (Salahshoor et al. 2012) and in vitro (Bucak et al. 2010; Soleimanzadeh and Saberivand 2013) investigations strongly suggest that curcumin plays a role in the energy-promoting and protective actions on testicular tissue, spermatogenesis, and sperm cell oxidative balance. Curcumin has also been proven to counteract toxic

effects on male genital system caused by a number of physiological or environmental causes (Wei et al. 2009; El-Wakf et al. 2011; Khorsandi et al. 2013).

Turmeric supplementation of up to 2% dry matter concentrate has also benefited the growth performance and health of lambs (Odhaib et al. 2021). Curcumin has beneficial effects on the sexual glands, testis and ovary, which may be related to its antioxidative (Sahoo et al. 2008), anti-inflammatory (Farombi et al. 2007), anti-cancer (Cort et al. 2012), and anti-apoptotic properties (Aktas et al. 2012). The purpose of the current study is to determine and evaluate the efficiency of *Curcuma longa* on Awassi male lambs' semen quality and some blood parameters.

MATERIALS AND METHODS

Ethical Approval

All procedures were approved by the University of Mosul, College of Veterinary Medicine, Institutional Animal Care and Use Committee, ethical approval (UM.VET).

Experimental Animals

The research was conducted in a private field in the village at Baibukht, which is located northeast of Mosul, Iraq. In semi-open barns, twenty-eight Awassi male lambs 47 ± 1.23 kg body weight, 10–11 months old) were randomly assigned to two pens (n=12 each pen) and fed concentrate meals.

Experimental Design

Awassi lambs (24) were randomly divided into two groups (n=12). T1, the control group was fed a standard ration, whereas T2, the experimental group was fed a standard ration (Table 1) supplemented with 200mg/kg *Curcuma longa* powder (CL). CL was added to the diet on daily basis and mixed well, the experiment lasted 90 days. CL product was purchased from a local market (Alhadbaa, Mosul, Iraq).

Sampling of Blood

On days 30, 60, and 90 of the experiment, blood samples were obtained from lambs through jugular venipuncture into a vacuum tube containing (EDTA) as an anticoagulant as well as plan tubes. Blood samples were examined for red blood cells (RBC), white blood cells (WBC), hemoglobin (Hb), packed cell volume (PCV), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC) using an automatic hematology analyzer (Mytic18) from Orphee Swiss company. Blood samples in plan tubes were centrifuged at 3,000 for 15 minutes at 4°C. An automatic analyzer separated serum to assess biochemical blood profiles such as total protein (TP), albumin (ALB), and globulin (GLU). The kit prepared by Lake Forest, Ca 92630, Monobind, Inc. USA, was used to assess testosterone and estrogen concentrations.

Semen Collection

At 30, 60, and 90 days of the experiment, semen was collected from 24 Awassi lambs for all treatments by inserting an electrode of Electro ejaculator into the rectum

Table 1: Ingredients and nutritive composition of standard ration, determined on a dry matter*.

Ingredients of dry mater %	Chemical compound	%
Crushed barley	Dry matter	92.09
Soybean meal	Organic matter	94.52
Wheat bran	Crude protein	14.64
Wheat straw	Ether Extract	2.04
Food Salt	Crude fiber	8.15
CaCo ₃	Ash	5.48
	ME (Kcal/kg/DM)	2534

*(Al-Khawaja 1987)

and performing batches of electrical stimulation every 5-10 seconds to obtain an ejaculate, which was collected in a graduated tube and then located in a 37°C water bath for semen tests according to Fourie et al. (2004).

Semen Evaluation

Once a month, semen was collected. All collected & processing glassware were washed and sterilized with high-pressure steam, then dried and reheated at 35°C. The tube was immediately placed in a 37°C bath after ejaculation. The ejaculate volume (EV) was directly measured using the graduated collecting tube. Gallenham, No. 82TT8, Cat No. M/6-200-H HZ 60, England was used for microscopic investigation. Wave motion of fresh undiluted semen was assessed under a microscope $\times 10$ with a 5 score to determine mass motility (MM). 5 μ of diluted semen was placed on a warmed (37°C) slide to assess sperm individual motility. A drop of diluted (1:400) semen was deposited on a hemocytometer to count the number of spermatozoa. Evaluation of the survival of spermatozoa was evaluated using the eosin-nigrosin stain. On a clean glass slide, a small drop of semen and one drop of eosin-nigrosin stain were combined with a clean stick. A thin smear was created, dried with in air, and examined under a 400 \times magnification microscope. To assess the percentages of Live sperms (LS) and abnormality (AB), at least 200 spermatozoa were analyzed from each smear. pH-indicator paper was used to measure the pH of the semen.

Statistical Analysis

The data was analyzed using the Statistical Software Statics program (SAS, 2003), according to the Complete Randomized Design's two-way analysis of variance (CRD) utilizing (GLM) procedure. According to Steel and Torrie (1984), the differences between the means were determined using Duncan's multiple range test at the probability level ($P \leq 0.05$).

RESULTS

Effect of CL on Semen Parameters

Table 2 shows that after 30, 60 and 90 days of treatment, the CL caused a significant ($P < 0.05$) increase in EV (1.11, 1.18 and 1.26ml respectively) compared to control group (0.83, 1.05 and 1.12ml respectively), the superior increase was at 90 day of treatment. Also, a comparison between the control group with CL group exhibited a significant increase in CE at 60 and 90 day of treatment (1.99 and 2.37×10^9 respectively) compared to control group (1.67 and 1.78×10^9 respectively). A comparison between the CL group with control group exhibited a significant increase in CM at 90 day of treatment,

Table 2: Influence of *Curcuma longa* on semen parameters of Awassi males

Parameters	Treatments	Days		
		30	60	90
Ejaculate volume (ml)	T ₁	0.83±0.02d	1.05±0.06c	1.12±0.03bc
	T ₂	1.11±0.05bc	1.18±0.01ab	1.26±0.02a
Concentration /ejaculate ×10 ⁹	T ₁	1.55±0.02d	1.67±0.04dc	1.78±0.04c
	T ₂	1.58±0.04d	1.99±0.08b	2.37±0.01a
Concentration/ml × 10 ⁹	T ₁	1.87±0.05a	1.62±0.09b	1.61±0.07b
	T ₂	1.44±0.08b	1.61±0.09b	1.87±0.04a
Live sperms%	T ₁	55.50±0.86d	58.62±0.96cd	61.00±0.70bc
	T ₂	56.12±0.58d	64.12±2.27b	70.37±1.63a
Abnormality%	T ₁	1.65±0.02a	1.57±0.02ab	1.57±0.03ab
	T ₂	1.56±0.02ab	1.52±0.02b	1.20±0.04c
Mass motility	T ₁	2.46±0.08c	2.87±0.04b	3.19±0.06a
	T ₂	2.53±0.01c	2.90±0.04b	3.22±0.09a
Individual motility%	T ₁	60.50±1.63cd	58.62±1.47d	61.87±1.98bcd
	T ₂	63.25±0.83cb	65.50±0.48b	70.62±0.59a
pH	T ₁	7.07±0.01a	7.10±0.02a	7.14±0.01a
	T ₂	7.05±0.05a	7.04±0.01a	7.18±0.10a

T₁ – Control, T₂ – CL 200 mg/kg ration, ^{a,b,c} means different alphabets in a row differ significantly (P<0.05) for each parameter. EV- Ejaculate volume/ml, CE – Concentration /ejaculate, CM - Concentration /ml, LS - Live sperms%, AB - Abnormality%, MM - Mass motility/5.

Table 3: Influence of *Curcuma longa* on blood picture of Awassi males

Parameters	Treatments	Days		
		30	60	90
WBCs (×10 ³ /μL)	T ₁	7.91±0.11c	8.01±0.09c	8.04±0.11c
	T ₂	8.15±0.06c	8.55±0.10b	9.23±0.16a
RBCs (×10 ⁶ /μL)	T ₁	9.09±0.07cd	9.27±0.20cd	8.97±0.25d
	T ₂	9.58±0.16c	10.29±0.13b	11.12±0.09a
Hb (gm/dL)	T ₁	10.30±0.18c	10.32±0.16c	10.12±0.13c
	T ₂	11.24±0.14b	11.46±0.15ab	11.84±0.03a
PCV (%)	T ₁	28.81±0.29c	28.92±0.09c	29.93±0.19c
	T ₂	29.98±0.06b	30.90±0.17a	31.20±0.10a
MCV (fL)	T ₁	31.72±0.53b	31.30±0.80b	33.52±0.90a
	T ₂	31.36±0.61b	30.05±0.28b	28.04±0.17c
MCH (Pg)	T ₁	11.34±0.21ab	11.18±0.37ab	11.34±0.36ab
	T ₂	11.75±0.22a	11.15±0.16ab	10.64±0.08b
MCHC (%)	T ₁	35.76±0.60b	35.68±0.49b	33.83±0.52c
	T ₂	37.49±0.47a	37.11±0.49ab	37.96±0.12a

T₁ – Control, T₂–CL 200 mg/kg ration, ^{a,b,c} means different alphabets in a row differ significantly (P<0.05) for each parameter.

Table 4: Influence of *Curcuma longa* on some serum blood biochemical traits of Awassi males

Parameters	Treatments	Days		
		30	60	90
Total protein (gm/dL)	T ₁	6.42±0.04b	6.42±0.06b	6.40±0.03b
	T ₂	6.74±0.04a	6.87±0.05a	6.88±0.12a
Albumin (gm/dL)	T ₁	2.84±0.03a	2.85±0.05a	2.92±0.09a
	T ₂	2.83±0.03a	2.96±0.07a	2.98±0.11a
Globulin (gm/dL)	T ₁	3.58±0.06b	3.57±0.09b	3.47±0.10b
	T ₂	3.91±0.04a	3.90±0.08a	3.89±0.08a

T₁ – Control, T₂–CL 200mg/kg ration, ^{a,b,c} means different alphabets in a row differ significantly (P<0.05) for each parameter.

1.87×10⁹ and 1.61×10⁹ respectively. LS increased significantly in CL group at 60 and 90 day of treatment (64.12 and 70.37% respectively) in comparison with control (58.62 and 61.00% respectively). AB improved significantly (P<0.05) in CL group (1.20%) compared to control (1.57%) at 90 day of treatment. The data obtained from Statistical analysis of MM and IM demonstrated a significant increase in MM in CL group at 60 and 90 day of treatment (2.90 and 3.22, respectively) in comparison with control (2.87 and 3.19, respectively) and in IM (65.50 and 70.62%, respectively) compared to control (58.62 and 61.87%, respectively). No significant differences were recorded in pH between the treatments.

The results of hematological parameters of treated lambs at different days are shown in Table 3, a significant

increase was observed (P<0.05) in WBC in blood samples of CL group at 60 and 90 day of treatment (8.55 and 9.23 ×10³/μL) compared with control group (8.01 and 8.04 ×10³/μL, respectively) and in RBC (10.29 and 11.12 ×10⁶/μL) compared with control group (9.27 and 8.97 ×10⁶/μL, respectively).

Effect of CL on Hematology Parameters

CL treatment caused a significant increase in Hb at 30, 60 and 90 day of treatment (11.24, 11.46 and 11.84gm/dL, respectively) in comparison with control group (10.30, 10.32 and 10.12 gm/dL, respectively), also CL causes a significant increase in PCV% at 30, 60 and 90 day of treatment (29.98, 30.90 and 31.20%, respectively) compared to control group (28.81, 28.92 and 29.93, respectively).

DISCUSSION




Fig. 1: Effect of *Curcuma longa* on Testosterone in Awassi males.




Fig. 2: Effect of *Curcuma longa* on Estrogen in Awassi males.

MCV decreased significantly at 90 day of treatment in CL group (28.04 fL) in comparison with control (33.52 micron³), while MCHC increased significantly ($P < 0.05$) in CL group at 30 and 90 day of treatment (37.49 and 37.96%, respectively) compared to control (35.76 and 33.83%, respectively).

Effect of CL on Serum Blood Biochemical Traits

Table 4 shows that CL supplementation (200mg/kg ration) caused a significant increase in the TP values (6.74, 6.87 and 6.88gm/dL) and GLU values (3.91, 3.90 and 3.89mg/dL) at all periods of treatment (33, 60 and 90 days) respectively compared with control (6.42, 6.42 and 6.40 gm/dL) for TP and 3.58, 3.57 and 3.47 gm/dL for GLU respectively at $P < 0.05$. On the other hand, CL treatment had no effects on ALB along the whole treatment periods.

In regard for testosterone and estrogen values, Fig. 1 revealed that CL enhanced significantly testosterone concentration at 30, 60 and 90 day of treatment (6.43, 6.47 and 6.66ng/mL, respectively) in comparison with control (6.18, 6.39 and 6.48ng/mL, respectively) at $P < 0.05$, also CL reduced estrogen concentration (2.34, 2.49 and 2.58pg/mL, respectively) along the whole treatment periods (30, 60 and 90 days) compared with control (2.5, 2.60 and 2.79pg/mL, respectively).

Recently, medicinal plants have been utilized to lessen the impact of oxidative stress on spermatozoa, which reduces sperm's capacity to fertilize eggs. Several studies have also focused on enhancing sperm quality and minimizing the impact of oxidative stress by supplementing sperm with antioxidants (Abdelnour et al. 2020). Statically analyzed data obtained in the current study showed a significant improvement in all semen parameters of CL group (200mg/kg ration), which was consistent with the findings of Tvrdá et al. (2015), whom added 0, 1, 5, 10, 50 and 100 μ M/L of curcumin to bovine semen. Moreover, Abdelnour et al. (2020), conducted a study in twelve buck's rabbits, pooled semen was cryopreserved in tris-yolk fructose extender supplemented with CL at levels of 0.5, or 1.5 μ g/mL, respectively, and observed a significant increase in sperm progressive motility and viability compared with control group, also the percentages of dead sperms and abnormalities were reduced significantly. In a study on semen of five mature bulls, Arboud et al. (2020) revealed the effect of CL on semen traits in diluted semen enriched with CL 0, 100, 200 and 300 μ L/5mL, and observed a significant improvement in semen motility and abnormalities. The results of current study were in agreement with the results of Kazemizadeh et al. (2019), who recorded a significant improvement in different parameters including semen concentration, total sperm production and progressive motility were linearly improved by the increasing levels of curcumin supplementation in broiler breeder roosters. In their study on adult Baladi bucks, Ismail et al. (2020) claimed that the enhancement of advance motility and viability belong to the kind of phenolic and flavonoid components in CL.

According to the Petruska et al. (2014), CL contains a lipophenol in soluble form that scavenges free radicals, and inhibits the generation of ROS. The authors reported that CL has an anti-oxidative effect to decrease lipid oxidative in the cell membrane of sperm and through down-regulation of the H₂O₂ level, which leads to protect spermatozoa from being injured (Zhang et al. 2017). The improvement of semen parameters might be associated with the portability of CL to reduce the protein and lipid oxidations (Abdelnour et al. 2020).

These results are in agreement with those of Odhaib et al. (2021), who observed a substantial increase in several blood parameters in their investigation on sixteen male lambs fed rations containing 1, 1.5, and 2% curcumin. In the current study, CL raised WBC, RBC, Hb, PCV, MCV, and MCHC. RBC and Hb levels may have grown over the course of the trial due to changes in metabolic needs brought on by the lamb's increased body weight and requirement for high oxygen levels. Moreover, Noori and sultan (2020) did not reported any significant differences in GLU, ALB, GLU and cholesterol in their study on twenty-four lambs reared on ration supplemented with CL (200mg/kg ration). Al Nahari and Al Eisa (2016) conducted a study to investigate the effect of curcumin on testosterone concentration in rats. The increase of testosterone concentration belongs to curcuminoids ability to improve spermatogenic damage and induce antioxidant defense.

Conclusions

We can conclude that CL (200 mg/kg ration) induced improvements in semen and hematological traits, when added to ration, and also significantly increased testosterone concentration in Awassi lamb's serum blood.

Acknowledgments

The University of Mosul and the College of Agriculture and Forestry were helpful in the author's research, and they are acknowledged. I also want to express my gratitude to Dr. Saeb Younis Abdul-Rahman for his assistance with this study.

Author's Contribution

Khalid H Sultan designed the plan of study, supervised the experiment, revised the manuscript writing and carried out language editing and formatting the manuscript.

REFERENCES

- Abdelnour SA, Hassan MA, Mohammed AK, Alhimaidei AR, Al-Gabri N, Al-Khalidi KO and Swelum AA, 2020. The effect of adding different levels of curcumin and its nanoparticles to extender on post-thaw quality of cryopreserved rabbit sperm. *Animals* 10: 1-13. <https://doi.org/10.3390/ani10091508>
- Abdul-Majeed AF, Rahawi GA and AL-Chalabi AM, 2021. Effect of adding nettle plant on some physiological and biochemical parameters of broiler chickens. *Iraqi Journal of Veterinary Sciences* 35: 115-119. <https://doi.org/10.33899/ijvs.2021.131844.2010>
- Aktas C, Kanter M, Erbogga M and Ozturk S, 2012. Anti-apoptotic effects of curcumin on cadmium-induced apoptosis in rat testes. *Toxicology and Industrial Health* 28: 122-130. <https://doi.org/10.1177/0748233711407242>
- Al Nahari H and Al Eisa R, 2016. Effect of turmeric (*Curcuma longa*) on some pituitary, thyroid and testosterone hormone against aluminum chloride (Al [Cl. sub. 3]) induced toxicity in rat. *Advances in Environmental Biology* 10: 250-262.
- Al-Khawaja AK, Al-Bayati EA and Matti SA, 1978. Chemical composition and nutritional value of Iraqi feed materials. *Bulletin of the Nutrition Section. Directorate of Public Animal Resources. Ministry of Agriculture and Agrarian Reform. Baghdad (Iraq).*
- Benkő R, Matuz M, Viola R, Doró P, Hajdu E and Soós G, 2008. Quantitative disparities in outpatient antibiotic exposure in a Hungarian county. *Journal of Antimicrobial Chemotherapy* 62: 1448-1450. <https://doi.org/10.1093/jac/dkn385>
- Bucak MN, Sarıözkan S, Tuncer PB, Sakin F, Ateşşahin A, Kulaksız R Çevik M, 2010. The effect of antioxidants on post-thawed Angora goat (*Capra hircus ancyrensis*) sperm parameters, lipid peroxidation and antioxidant activities. *Small Ruminant Research* 89: 24-30. <https://doi.org/10.1093/biolreprod/83.s1.496>
- Coradini K, Lima FO, Oliveira CM, Chaves PS, Athayde ML, Carvalho LM and Beck RC, 2014. Co-encapsulation of resveratrol and curcumin in lipid-core nanocapsules improves their in vitro antioxidant effects. *European Journal of Pharmaceutics and Biopharmaceutics* 88: 178-185. <https://doi.org/10.1016/j.ejpb.2014.04.009>
- Cort A, Timur M, Ozdemir E, Kucuksayan E and Ozben T, 2012. Synergistic anticancer activity of curcumin and bleomycin: an in vitro study using human malignant testicular germ cells. *Molecular Medicine Reports* 5: 1481-1506. <https://doi.org/10.3892/mmr.2012.848>
- El-Wakf ME, Azza M, Elhabiby MW, El-kholy EG and El-Ghany EA, 2011. Use of turmeric and curcumin to alleviate adverse reproductive outcomes of water: Nitrate pollution in male rats. *Nature and Science* 11: 229-239.
- Farombi EO, Abarikwu SO, Adedara IA and Oyeyemi MO, 2007. Curcumin and kolaviron ameliorate di-n-butylphthalate-induced testicular damage in rats. *Basic and Clinical Pharmacology & Toxicology* 100: 43-48. <https://doi.org/10.1111/j.1742-7843.2007.00005.x>.
- Fourie PJ, Schwalbach LM, Naser EWC and Van der Westhuizen C, 2004. Scrotal, testicular and semen characteristics of young Dorper rams managed under intensive and extensive conditions. *Small Ruminant Research* 54: 53-59. <https://doi.org/10.1016/j.smallrumres.2003.10.011>
- FrAnKlIĆ T, Voljč M, Salobir J and Rezar V, 2009. Use of herbs and spices and their extracts in animal nutrition. *Acta Agriculturae Slovenica* 94: 95-102.
- García-Hernández C, Arece-García J, Rojo-Rubio R, Mendoza-Martínez GD, Albarrán-Portillo B, Vázquez-Armijo JF, Avendaño-Reyes L, Olmedo-Juárez A, Marie-Magdeleine C and López-Leyva Y, 2017. Nutraceutical effect of free condensed tannins of *Lysiloma acapulcensis* (Kunth) benth on parasite infection and performance of Pelibuey sheep. *Tropical Animal Health and Production* 49: 55-61. <https://doi.org/10.1007/s11250-016-1157-8>
- Hameed HM, Aga FK and Abdulrahman SY, 2020. Effect of β -mannanase, Lysolecithin and probiotic on some reproductive performance and hormone profile in female quail. *Iraqi Journal of Veterinary Sciences* 34: 87-93. <https://doi.org/10.33899/ijvs.2019.125587.1097>
- Ismail AA, Abdel-Khalek AK, Khalil WA, Yousif AI, Saadeldin IM, Abomughaid MM and El-Harairy MA, 2020. Effects of mint, thyme, and curcumin extract nanoformulations on the sperm quality, apoptosis, chromatin decondensation, enzyme activity, and oxidative status of cryopreserved goat semen. *Cryobiology* 97: 144-152. <https://doi.org/10.1016/j.cryobiol.2020.09.002>
- Jaguezeski AM, Perin G, Bottari NB, Wagner R, Fagundes MB, Schetinger MR, Morsch VM, Stein CS, Moresco RN, Barreta DA and Danieli B, 2018. Addition of curcumin to the diet of dairy sheep improves health, performance and milk quality. *Animal Feed Science and Technology* 246: 144-157. <https://doi.org/10.1016/j.anifeedsci.2018.10.010>
- Karásková K, Suchý P and Straková E, 2015. Current use of phyto-genic feed additives in animal nutrition: a review. *Czech Journal of Animal Science* 60: 521-430. <https://doi.org/10.17221/8594-CJAS>
- Kazemizadeh A, Zare Shahneh A, Zeinoaldini S, Yousefi AR, Mehrabani Yeganeh H, Ansari Pirsaraei Z and Akhlaghi A, 2019. Effects of dietary curcumin supplementation on seminal quality indices and fertility rate in broiler breeder roosters. *British Poultry Science* 60: 256-264. <https://doi.org/10.1080/00071668.2019.1571165>
- Khan RU, Shabana N, Kuldeep D, Karthik K, Ruchi T, Abdelrahman MM, Alhidary IA and Arshad Z, 2016. Directed microbial: beneficial applications, modes of action and prospects as a safe tool for enhancing ruminant production and safeguarding health. *International Journal of Pharmacology* 12: 220-231. <https://doi.org/10.3923/ijp.2016.220.231>
- Khorsandi L, Mirhoseini M, Mohamadpour M, Orazizadeh M and Khaghani S, 2013. Effect of curcumin on dexamethasone-induced testicular toxicity in mice. *Pharmaceutical Biology* 51: 206-212. <https://doi.org/10.3109/13880209.2012.716854>
- Mendel M, Chłopecka M, Dziekan N and Karlik W, 2017. Phyto-genic feed additives as potential gut contractility modifiers—A review. *Animal Feed Science and Technology* 230: 30-46. <http://dx.doi.org/10.1016/j.anifeedsci.2017.05.008>
- Molosse V, Souza CF, Baldissera MD, Glombowsky P, Campigotto G, Stefani LM and da Silva AS, 2019. Diet

- supplemented with curcumin for nursing lambs improves animal growth, energetic metabolism, and performance of the antioxidant and immune systems. *Small Ruminant Research* 170: 74–81. <https://doi.org/10.1016/j.smallrumres.2018.11.014>
- Arboud MM, Waheeb RS, El-Sheshtawy RI and El-Amrawi GA, 2020. Assessment of cattle bull semen preservability using tris extender enriched with turmeric extract. *Egyptian Journal of Veterinary Sciences* 51: 357-362. <https://doi.org/10.21608/ejvs.2020.27051.1168>
- Noori MS and Sultan KH, 2020. Effect of wool shearing and diet supplementing with curcumin on some productive and blood biochemical traits of Awassi lambs. *Mesopotamia Journal of Agriculture* 48: 24-32. <https://doi.org/10.33899/magrj.2020.126964.1041>
- Odhaib KJ, Ali NM, Abdulameer HA and Khudhair NA, 2021. Influence of graded levels of turmeric (*Curcuma longa*) as feed additives alternatives to promote growth and enhance health status in lambs. *Biochemical and Cellular Archives* 21: 3025-3032.
- Petruska P, Capcarova M and Sutovsky P, 2014. Antioxidant supplementation and purification of semen for improved artificial insemination in livestock species. *The Turkish Journal of Veterinary and Animal Sciences* 38: 643-652. <https://doi.org/10.3906/vet-1404-61>
- Sahoo DK, Roy A and Chainy GB, 2008. Protective effects of vitamin E and curcumin on L-thyroxine-induced rat testicular oxidative stress. *Chemico-Biological Interactions* 176: 121-128. <https://doi.org/10.1016/j.cbi.2008.07.009>
- Salahshoor MR, Jalili C, Khazaei M and Khani F, 2012. Effects of curcumin on reproductive parameters in male mice. *Research in Paramedical Sciences* 1: 31-38.
- SAS Institute, SAS/STAT user's guide for personal computers. Release V. 9.1. SAS, 2003. Institute Inc., Cary, NC (USA).
- Schwarz S, Kehrenberg C and Walsh TR, 2001. Use of antimicrobial agents in veterinary medicine and food animal production. *International Journal of Antimicrobial Agents* 17: 431-437. [https://doi.org/10.1016/s0924-8579\(01\)00297-7](https://doi.org/10.1016/s0924-8579(01)00297-7)
- Soleimanzadeh A and Saberivand A, 2013. Effect of curcumin on rat sperm morphology after the freeze-thawing process. *Veterinary Research Forum* 4: 185–189.
- Steel RGD and Torrie JH, 1984. Principles procedures of statistics. McGraw Hill Book Company, Inc., New York (USA), pp :481.
- Tvrđá E, Lukáč N, Jambor T, Lukáčová J and Massányi P, 2015. Curcumin in male fertility, effects on spermatozoa vitality and oxidative balance. *Journal of Microbiology, Biotechnology and Food Science* 4: 120-124. <https://doi.org/10.15414/jmbfs.2015.4.special2.120-124>
- Wei SM, Yan ZZ and Zhou J, 2009. Curcumin attenuates ischemia–reperfusion injury in rat testis. *Fertility and Sterility* 91: 271-277. <https://doi.org/10.1016/j.fertnstert.2007.10.082>
- Wencelová M, Váradyová Z, Mihaliková K, Čobanová K, Plachá I, Pristaš P, Jalč D and Kišidayová S, 2015. Rumen fermentation pattern, lipid metabolism and the microbial community of sheep fed a high-concentrate diet supplemented with a mix of medicinal plants. *Small Ruminant Research* 125: 64-72. <https://doi.org/10.1016/j.smallrumres.2015.01.028>
- Zhang L, Diao RY, Duan YG, Yi TH and Cai ZM, 2017. In vitro antioxidant effect of curcumin on human sperm quality in leucocytospermia. *Andrologia* 49: e12760. <https://doi.org/10.1111/and.12760>