



Improving Feed Quality by Adding Coconut Pulp in the Concentrate Formation to Enhance Production Performance of Kacang Goats

Padang^{1*}, Sri Wulan¹, Nirwana¹, Fatmawati¹, Zainal¹, Harmoko² and Naharuddin³

¹Department of Animal Husbandry, Faculty of Animal and fishery-Tadulako University, Palu 94148, Central Sulawesi, Indonesia

²Department of Animal Husbandry, Study Program Outside the Main Campus-Pattimura University, Ambon 97233, Maluku, Indonesia

³Department of Forestry, Faculty of Forestry-Tadulako University, Palu 94118, Central Sulawesi, Indonesia

*Corresponding author: padanghamid2608@gmail.com

Article History: 24-627 Received: 21-Sep-24 Revised: 25-Oct-24 Accepted: 31-Oct-24 Online First: 10-Nov-24

ABSTRACT

Efforts to increase the physiological productivity of livestock are one of the problems in responding to the increasing demand for livestock meat in Indonesia. A study was done to investigate the effects of adding coconut pulp to a supplemental diet on the productivity and physiological responses of female Kacang goats. A 2 x 3 factorial design was used in which the first factor was two levels of coconut pulp treatments (A1=non-fermented, A2=fermented), and the second factor was three levels of coconut pulp in the supplement (L0=0%, L1=10%, L2=20%). Five animals were used for each treatment. An analysis of variance showed that adding fermented or non-fermented coconut pulp at various levels did not affect weight gain significantly ($P>0.05$), dry matter intake, feed efficiency, physiological measures (body temperature, respiratory and pulse rates), or hematological parameters (white/red blood cell counts, hemoglobin, hematocrit). Additionally, no significant effects ($P>0.05$) were observed on slaughter weight, carcass metrics, or non-carcass components. However, the amount of coconut pulp significantly affected ($P<0.05$) certain parameters: white blood cell count, carcass weight, carcass percentage, and specific non-carcass weight measures, including internal, external, and edible components. Overall, coconut pulp inclusion in the supplemental diet, regardless of fermentation, did not notably influence most productivity and physiological measures, though the level of inclusion impacted specific non-carcass and blood parameters.

Key words: Kacang Goat, Coconut Pulp, Fermentation, Productivity, Physiological Condition and Carcass.

INTRODUCTION

Feed is the most significant factor in livestock farming, accounting for 60-80% of production costs and is often affected by seasonal fluctuations. In addition, the primary feed given to livestock is grass, which cannot meet their needs both in terms of quantity and quality. To overcome the limitation, farms are frequently supplement with high-quality concrete. However, the high costs of the concentrates as well as competition from humans and other livestock demands presents a significant challenge. This indicates that there is a need to explore alternative to traditional feed ingredients by using agricultural waste that does not contribute to environmental pollution. One such effort comprises repurposing agricultural and industrial by-products, such as coconut pulp, for producing animal feed.

According to previous studies, coconut pulp can be

obtained from VCO industries and household-scale oil production (Suryani et al. 2020). This pulp can be used directly as animal feed but can be processed first to extend the shelf life and improve its original quality (Khaidir et al. 2015; Singh and Krishnaswamy 2022). Coconut pulp that is not utilized by the community can still be used as an alternative feed for livestock because it contains nutrients needed for growth (Emilia et al. 2021; Radhiah et al. 2022).

In line with these findings, the process of making coconut generates solid waste in the form of coconut pulp, which accounts for approximately 30% of the raw materials used during the process (Ouma 2019; Kristianto 2023). According to Sutanto et al. (2021); Elyana et al. (2018), coconut pulp is an industrial or household waste that has great potential to be used as feed ingredients, because it is still easy to obtain from the remains of traditional oil production.

Cite This Article as: Padang, Wulan S, Nirwana, Fatmawati, Zainal, Harmoko and Naharuddin, 2025. Improving feed quality by adding coconut pulp in the concentrate formation to enhance production performance of Kacang goats. International Journal of Veterinary Science 14(2): 416-426. <https://doi.org/10.47278/journal.ijvs/2024.255>

Despite the potential, the use of coconut pulp as a substitute for animal feed is not yet widely recognized (Marhamah et al. 2019; Kholishah et al. 2021; Oktaviana et al. 2022). Although it is a by-product of coconut milk production, it has a fairly high crude fiber content (Panjaitan 2021; Londok et al. 2022). In Central Sulawesi, Indonesia, research on the use of plants is still focused, for example the effect of castor leaves on the productivity of Kacang goats (Naser et al. 2023), the effect of moringa powder and fruit on the growth performance of Kacang goats (Sagaf et al. 2024). The effect of giving *Lannea coromandelica* leaf flour on the production performance and hematological value of Kacang goats (Abdullah et al. 2023).

Adequate crude fiber in feed is beneficial for promoting smooth digestion in the intestines. Dietary fiber has been reported to play a crucial role in disease prevention and is an essential component of nutritional therapy (Yulvianti et al. 2015; Alissandra et al. 2019; Laksono et al. 2023). However, despite its high fiber content, coconut pulp has low palatability, lacks several essential amino acids, possesses anti-nutritional properties, and has low digestibility. These factors have limited its use as animal feed, and the full extent of these limitations has yet to be fully explored. To overcome these challenges, improving the quality of coconut pulp through fermentation and processing it into coconut pulp flour is essential. Pre-treating the waste with microorganisms before feeding it to livestock is one approach to enhancing the efficiency of rumen microorganisms, making the utilization of feed more effective in meeting the animals' basic needs for maintenance, production, and reproduction. Therefore, this study aims to determine the effect of adding coconut pulp to the concentrates composition on the productivity and physiological condition of female Kacang goats.

MATERIALS AND METHODS

Experimental Livestock

This study involved 30 female Kacang goats, approximately 12 months old, with body weights ranging from 10 to 20kg. The experiment utilized a Randomized

Block Design (RBD) with a 2 x 3 factorial arrangement based on initial body weight. The first factor was the type of coconut pulp: A1=unfermented coconut pulp, and A2=fermented coconut pulp. The second factor was the level of coconut pulp in the concentrate: L0=no coconut pulp, L1=10% coconut pulp, and L2=20% coconut pulp. Each treatment was replicated five times. The goats were housed in tiered pens with a zinc roof, plank flooring, and plank walls, measuring 6 x 10 meters. The pens were divided into 30 sections, each 1.0 x 1.0m, with one goat per section. Each section was equipped with a wooden feed trough and a drinking basin. The goats were acclimated to the new feed over a 14-day period, followed by a 56-day data collection phase. Feed was provided daily to monitor consumption. Body weight was recorded weekly in the morning before feeding. At the conclusion of the study, the goats were slaughtered for assessment of carcass and non-carcass components. This study received approval from the animal ethics committee of the Faculty of Animal Husbandry and Fisheries, Tadulako University, Palu, Indonesia.

Animal Feed

During the study, the feed consisted of concentrate and tubers. The concentrate was composed of 18% ground soybeans, 48% rice bran and 34% ground corn, along with varying levels of fermented and unfermented coconut pulp as treatments. The concentrate was provided at a rate of 1.0% dry matter based on the goats' body weight, while tubers were offered after the goats had consumed the concentrate and treatment as much as possible. The nutritional content of the feed ingredients is detailed in Table 1 and the nutritional content of the treatments is presented in Table 2.

Making Fermented Coconut Pulp

A total of 10 kg of coconut pulp was steamed for 20 minutes and then cooled. After cooling, the tape yeast was sprayed evenly with a dose of 1 l/1 ton and placed in plastic to facilitate fermentation for 5 days. Subsequently, the sample was dried for 24 hours and ready to be mixed with other feed ingredients.

Table 1: Nutritional Content of Feed Ingredients Used

| Feed Ingredients | Dry Ingredients (%) | Crude protein (%) | Coarse Fiber (%) | Crude Fat (%) | TDN (%) |
|------------------------------|---------------------|-------------------|------------------|---------------|---------|
| Ground Soybeans | 91.97 | 31.35 | 9.73 | 11.65 | 61.00 |
| Ground Corn | 86.82 | 9.54 | 9.92 | 8.30 | 80.87 |
| Rice Bran | 89.92 | 10.67 | 18.39 | 4.62 | 61.21 |
| Coconut Pulp (AK) | 90.98 | 7.47 | 9.01 | 31.76 | 66.77 |
| Fermented Coconut Pulp (AKF) | 93.12 | 10.50 | 6.90 | 27.34 | 59.43 |
| <i>Panicum sharmentosum</i> | 26.29 | 11.51 | 30.20 | 1.90 | 59.54 |

Description results of the analysis of the Feed Nutrition Laboratory of the Faculty of Animal Husbandry and Fisheries, Tadulako University

Table 2: Nutritional Content of Treatment during the Study

| Treatment | Dry Ingredients (%) | Crude protein (%) | Coarse Fiber (%) | Crude Fat (%) | TDN (%) |
|-------------------------------|---------------------|-------------------|------------------|---------------|---------|
| A ₁ L ₀ | 89.24 | 14.01 | 13.95 | 7.14 | 67.86 |
| A ₁ L ₁ | 89.41 | 13.35 | 13.46 | 9.60 | 67.75 |
| A ₁ L ₂ | 89.58 | 12.70 | 12.96 | 12.06 | 67.64 |
| A ₂ L ₀ | 89.24 | 14.01 | 13.95 | 7.14 | 67.86 |
| A ₂ L ₁ | 89.62 | 13.66 | 13.25 | 9.16 | 67.01 |
| A ₂ L ₂ | 90.01 | 13.31 | 12.54 | 11.18 | 66.17 |

Description A₁L₀ = 0.0% unfermented coconut pulp; A₁L₁ = 10.0% unfermented coconut pulp; A₁L₂ = 20.0% unfermented coconut pulp; A₂L₀ = 0.0% fermented coconut pulp; A₂L₁ = 10.0% fermented coconut pulp; A₂L₂ = 20.0% fermented coconut pulp

Variables and their Measurements

Some of the dependent variables observed in this study included production performance, rectal temperature, respiratory rate, and pulse rate.

Production performance

Weight gain was calculated as the difference between the final weight and the initial weight during the observation period. Weighing was carried out at the beginning and end of the reported week by referring to the formula:

$$PBBH(\text{g/head/day}) = \frac{W_2 - W_1}{T_2 - T_1}$$

Description: PBBH is daily weight gain (grams), W1 is initial weight, W2 is final weight, T1 is time of first weighing, and T2 is time of final weighing. Ration consumption based on dry matter (grams) was obtained by calculating the difference between the amount of feed given and the amount of feed remaining, then multiplied by the resulting dry matter content, expressed in grams/head/day, while the efficiency of ration use was obtained by dividing PBBH by daily dry matter consumption.

Rectal temperature, respiratory rate, pulse rate (pulse frequency)

Rectal temperature (SR), respiratory rate (FR), and pulse rate (DN) were measured daily at noon (12.30 UTC+8) and morning (3.30 UTC+8). SR was measured for 1 minute using a clinical thermometer inserted into the rectum. FR was calculated by counting the diaphragm movements for 1 minute, and DN was measured by hand on the femoral vein for 1 minute.

Blood collection

Blood was taken from the jugular vein of the experimental animals at the beginning of the study and the end of the experiment, before feeding at 07.30am, in a tube containing ethylene diamine tetra acetate (EDTA). With a disposable needle, 5mL of whole blood was taken aseptically from the jugular vein. The rest was placed in a simple tube with 2mL of blood in a vacutainer tube containing EDTA for serum examination. The EDTA tube was immediately closed and the contents were stirred slowly for one minute by inverting or shaking. Furthermore, the hematocrit value (Packed cell volume=PCV) and hemoglobin (Hb) concentration of the blood samples were determined immediately after blood sampling. After dilution, the number of red blood cells and white blood cells was determined using a Neubauer hemocytometer.

Carcass observation

After the livestock were fasted overnight, all goats were slaughtered for carcass examination. This involved severing the jugular vein and carotid artery and allowing the goats to bleed for several minutes. The goats were then skinned and eviscerated immediately after bleeding. Each goat's body parts were separated, weighed, and recorded. The carcass percentage was calculated by dividing the hot carcass weight by the slaughter weight or empty body weight. After removing all digestive tract contents, which were included as part of the slaughter weight, the empty

body weight was determined. The hot carcass weight was calculated after removing the head, as well as the contents of the chest, abdomen, and flanks, along with the skin and legs. In this report, total edible non-carcass components (edible offal) were defined as the sum of the kidneys, heart, liver, tongue, small and large intestines, reticulorumen, omasum, abomasum, tail, and total fat. Total inedible non-carcass components (non-edible offal) were calculated by summing the skin, digestive tract contents, sexual organs, bladder, spleen, lungs with trachea, diaphragm, and esophagus.

Data analysis

This study was processed statistically using RBD. Observation data obtained were analyzed by analysis of variance (F test), when there was a significant effect continued with the Least Significant Difference (LSD) Test to determine the average difference in the effect of treatment.

RESULTS AND DISCUSSION

Production performance

The results of observations on the effect of adding coconut pulp to the concentrate composition on the production performance of female Kacang goats are presented in Table 3.

The analysis of variance indicated that the addition of fermented and unfermented coconut pulp at various levels in the concentrate composition did not significantly affect ($P>0.05$) body weight gain, dry matter intake, or feed utilization efficiency in female Kacang goats.

The observed weight gain, dry matter intake, and feed utilization efficiency were attributable to the nutrient contributions from the fermented and unfermented coconut pulp at different levels, which did not alter the overall nutritional composition of the feed. Initially, the concentrate's nutritional content was 14.01% protein and 67.86% Total Digestible Nutrients (TDN) before the addition of coconut pulp. After incorporating 20% coconut pulp, the nutrient content was 12.70% protein and 67.64% TDN for unfermented coconut pulp, and 13.31% protein and 66.71% TDN for fermented coconut pulp (Table 2). This is consistent with the recommended protein requirements for goats, which range from 12-14% (Nsahlai et al. 2004; Souza et al. 2021; Fu et al. 2023). Consequently, the lack of significant impact on body weight, dry matter intake, and feed efficiency when coconut pulp was added at various levels can be attributed to the fact that these levels remained within the recommended protein range for goats.

Physiological conditions

The results of observations on the effect of adding coconut pulp to the concentrate composition on the physiological conditions of local goats are presented in Table 4.

Analysis of variance showed no interaction ($P>0.05$) between the addition of fermented and unfermented coconut pulp with different levels in the concentrate composition on body temperature, respiration frequency, pulse frequency, number of white blood cells, number of red blood cells, hemoglobin levels, and hematocrit values

Table 3: Average weight gain, feed consumption, and feed utilization efficiency in Kacang goats fed with additional coconut pulp in the concentrate composition

| Parameter | Coconut Pulp | Level (%) | | | Average |
|-----------------------------------|----------------------|-----------|--------|--------|---------|
| | | 0.0 | 10.0 | 20.0 | |
| Weight Gain (g/head/day) | Without Fermentation | 39.29 | 35.54 | 33.68 | 36.17 |
| | Fermentation | 37.32 | 38.07 | 36.07 | 37.15 |
| | Average | 38.30 | 36.80 | 34.88 | |
| Dry Feed Consumption (g/head/day) | Without Fermentation | 495.89 | 463.06 | 469.32 | 476.09 |
| | Fermentation | 489.24 | 481.70 | 484.37 | 485.11 |
| | Average | 492.57 | 472.38 | 476.84 | |
| Efficiency of Dry Feed Use | Without Fermentation | 0.080 | 0.076 | 0.072 | 0.076 |
| | Fermentation | 0.077 | 0.080 | 0.075 | 0.077 |
| | Average | 0.078 | 0.078 | 0.073 | |

Table 4: Average Physiological Condition of Kacang Goats Given Additional Coconut Pulp in the Concentrate Arrangement

| Parameter | Coconut Pulp | Level (%) | | | Average |
|--|----------------------|-----------|--------|--------|---------|
| | | 0.0 | 10.0 | 20.0 | |
| Rectal Temperature(°C) | Without Fermentation | 38.93 | 38.68 | 38.64 | 38.75 |
| | Fermentation | 38.87 | 38.87 | 38.75 | 38.83 |
| | Average | 38.90 | 38.78 | 38.69 | |
| Respiratory Rate (times/minute) | Without Fermentation | 51.30 | 46.11 | 46.02 | 47.81 |
| | Fermentation | 52.88 | 51.21 | 48.19 | 50.76 |
| | Average | 52.09 | 48.66 | 47.11 | |
| Pulsus Rate (times/minute) | Without Fermentation | 80.58 | 75.83 | 75.59 | 77.33 |
| | Fermentation | 79.60 | 78.78 | 77.68 | 78.69 |
| | Average | 80.09 | 77.30 | 76.64 | |
| White Blood Cells (thousands/mm ³) | Without Fermentation | 15.23 | 20.44 | 21.62 | 19.10 |
| | Fermentation | 16.10 | 19.78 | 20.34 | 18.74 |
| | Average | 15.66a | 20.11b | 20.98b | |
| Red Blood Cells (millions/mm ³) | Without Fermentation | 9.84 | 9.63 | 9.44 | 9.64 |
| | Fermentation | 9.83 | 9.83 | 9.82 | 9.83 |
| | Average | 9.84 | 9.73 | 9.63 | |
| Hemoglobin (g/dL) | Without Fermentation | 9.83 | 8.74 | 8.66 | 9.08 |
| | Fermentation | 9.30 | 9.24 | 9.22 | 9.25 |
| | Average | 9.56 | 8.99 | 8.94 | |
| Hematocrit (%) | Without Fermentation | 24.85 | 24.36 | 24.30 | 24.50 |
| | Fermentation | 24.75 | 24.72 | 24.54 | 24.67 |
| | Average | 24.80 | 24.54 | 24.42 | |

Description - Different letters towards the row indicate significant differences (P<0.05).

of female Kacang goats. Additionally, the provision of fermented and unfermented coconut pulp in the concentrate composition did not provide a significant effect (P>0.05) on body temperature, respiration frequency, pulse frequency, number of white blood cells, number of red blood cells, hemoglobin levels, and hematocrit values of female Kacang goats. The level of pulp provision in the concentrate composition did not provide a significant effect (P>0.05) on body temperature, respiration frequency, pulse frequency, number of red blood cells, hemoglobin levels, and hematocrit values, except for the number of white blood cells of female Kacang goats.

The results of this study indicated that the body temperature of the goats obtained was higher than the results of the report by Hairmoko and Padang (2019) namely 38.23-38.48°C. The body temperature of the goats was within the normal range according to the statement of Rajion et al. (2001) that the normal body temperature in goats ranged from 38.40-39.40°C and the SR of goats in normal conditions was 39.20-40.20°C (Rahardja et al. 2011).

This was related to metabolic activity, where the heat production produced was the same because the nutritional content was within the range of goat needs. The feed

consumed by livestock could affect the rate of heat production in the body called the calorogenic effect of feed and maintain homeostasis conditions (Härter et al. 2017; Costa et al. 2019; Fu et al. 2023).

The respiratory frequency that did not differ between treatments was caused by relatively similar feed consumption, resulting in no different results for the respiration frequency of Kacang goats. The level of feed consumption affected the rate of respiration frequency in ruminant livestock and could increase the body's metabolic process, therefore, the body heat produced also increased (Navarro et al. 2019; Palulungan et al. 2022; Loučka et al. 2023).

The frequency of pulses that did not differ between treatments was caused by the amount of feed consumption for each treatment not differing, both unfermented and fermented treatments with different levels in the concentrate composition, therefore affecting the frequency of pulses in livestock. This was as stated by Gonzalez-Rivas et al. (2020); Nirwana et al. (2021) that the frequency of pulses was closely related to the metabolic rate of livestock, where factors that influenced the frequency of pulses in livestock included gender, muscle activity, environmental temperature, and feed consumption levels.

The results of the LSD Test showed that the number of white blood cells in female Kacang goats given 10-20% coconut pulp in the concentrate composition was significantly different ($P < 0.05$) from female Kacang goats not given coconut pulp in the concentrate composition.

The number of white blood cells that did not differ between the treatments of fermented and unfermented coconut pulp feed ingredients was probably due to the possibility that fermented and unfermented coconut pulp feed ingredients were not foreign objects or feed ingredients containing toxins. Therefore, livestock responded by forming white blood cells that were not different, but after adding coconut pulp levels to the concentrate composition, the body responded to it as a foreign object or toxic substance. Elnaggar et al. (2016); Agustina et al. (2023) explained that leukocytes were mobile/active units of the body's defense system. Leukocytes had several basic properties, namely chemotaxis (attracted to certain chemicals), diapedesis (able to pass through the capillary membrane of blood vessels), amoeboid movement (moving like an amoeba), and phagocytosis (eating foreign objects) (Sipos 2019; Bradford and Contreras 2024). This could carry out their functions defensively (preventing foreign objects from entering the body) and reparatively (repairing damaged body parts) (Ruiz-Campillo et al. 2017; Hussien and Al-Sukruwah 2022; Scatà et al. 2023).

The average number of erythrocytes from this study was lower than the results conducted by Sarmin (2020); Soeyono et al. (2020). Reports by Widyono et al. (2014), on female Kacang goats that were intensively raised, obtained an average number of erythrocytes of $13.23 \pm 1.74 \times 10^6/\mu\text{L}$. Meanwhile, from the study by Bijanti et al. (2011), on female Kacang goats in Mojosari Village, Gresik, an average number of erythrocytes was obtained of $14.57 \pm 2.3 \times 10^6/\mu\text{L}$.

The decrease in the number of erythrocytes in female Kacang goats related to feed could be caused by a lack of protein and several minerals from feed that were needed in the process of erythrocyte formation such as Iron (Fe), Cobalt (Co), and Cuprum (Cu) (Iftitah et al. 2022; Shevchenko et al. 2023). The lack of minerals was caused by goats' insufficient nutritional intake from feed consumption.

The results of the examination of hemoglobin levels in female Kacang goats could be seen in Table 4, and this was 8.66-9.30g/dL. The hemoglobin levels obtained tended to be lower compared to the study results reported by Crilly and Plate (2022) 6.80-10.13g/dL and by Tresia et al. (2023) 9.16-10.84g/dL.

The hemoglobin levels obtained from the study results showed values that tended to be higher compared to the existing literature. This was due to the influence of the quality and quantity of feed given on the formation of hemoglobin however hemoglobin levels tended to be high. Singh et al. (2018), Sudarman et al. (2019) and Roza et al. (2024) stated that buffaloes given high-concentrate feed showed higher hemoglobin levels. These were also related to the Iron content in the feed (Zerfu et al. 2023). Iron was mainly needed in the process of erythrocyte formation, namely in hemoglobin synthesis (Lambrecht et al. 2021; Nohara et al. 2022). The element Iron was the main

component of hemoglobin, however, Iron deficiency affected hemoglobin formation (Choi et al. 2021; Iftitah et al. 2022; Anggita et al. 2023). Reduced Iron absorption caused the amount of ferritin (Iron stored in the body) to also decrease which had an impact on the amount of Iron that was used for hemoglobin synthesis and could cause anemia (Antoniani et al. 2018). Tana et al. (2018) and Soul et al. (2019) stated that hemoglobin levels were also influenced by the season, body activity, the presence or absence of erythrocyte damage, handling direction during examination, and nutrition in feed.

The results of the hematocrit value examination in female Kacang goats could be seen in Table 4. The average hematocrit value in female Kacang goats was 24.30-24.85%. The hematocrit value obtained was lower than the results of the reported study (Widyono et al. 2014) and higher than the report of (Bijanti et al. 2011). The previous report (Widyono et al. 2014), showed a hematocrit value of 28.58%, and another study (Bijanti et al. 2011), obtained a hematocrit value of 15.32%. Differences in hematocrit values were influenced by several factors such as age, livestock activity, water consumption, environmental temperature, and nutritional content in the feed, especially protein, minerals, and vitamins needed to maintain normality and hematocrit values (Corrales-Hernández et al. 2018; Kertawirawan et al. 2022).

Hematocrit values had a very close relationship with the number of erythrocytes. A decrease in the number of erythrocytes was generally followed by a decrease in hematocrit values. The pattern of increasing hematocrit values in this study showed appropriate values in goats. An increase in hematocrit values accompanied by an increase in hemoglobin levels and the number of red blood cells were thought to be caused by a lack of Iron and vitamin B6 in animals (Corrales-Hernández et al. 2018; Kertawirawan et al. 2022; Francisco et al. 2023)

Carcass and non-carcass

The results of observations on the effect of adding coconut pulp in the concentrate composition on the weight and percentage of carcasses and non-carcasses of female Kacang goats are presented in Table 5.

Analysis of variance showed that there was no interaction ($P > 0.05$) between the addition of fermented and unfermented coconut pulp with different levels in the concentrate composition on the slaughter weight, non-carcass weight (internal, external, and edible) and non-carcass percentage (internal, external and edible) of female Kacang goats, but had a significant effect ($P < 0.05$) on carcass weight and percentage. The provision of fermented and unfermented coconut pulp in the concentrate composition had no significant effect ($P > 0.05$) on the slaughter weight, carcass weight, carcass percentage, non-carcass weight (internal, external, and edible), and non-carcass percentage (internal, external, and edible) of female Kacang goats. The level of coconut pulp provision in the concentrate composition had no significant effect ($P > 0.05$) on the slaughter weight and edible non-carcass percentage and had an effect ($P < 0.05$) on the carcass weight, carcass percentage, non-carcass weight (internal, external and edible) and non-carcass percentage (internal and external) of female Kacang goats.

Table 5: Average Weight and Percentage of Carcass and Non-Carcass of Kacang Goats Fed with Additional Coconut Pulp in the Concentrate Arrangement

| Parameter | Coconut Pulp | Level (%) | | | Average |
|--|----------------------|-----------|---------|--------|---------|
| | | 0.0 | 10.0 | 20.0 | |
| Slaughter weight (kg/head) | Without Fermentation | 14.78 | 13.24 | 13.72 | 13.92 |
| | Fermentation | 14.13 | 14.02 | 13.71 | 13.95 |
| | Average | 14.45 | 13.63 | 13.72 | |
| Carcass weight (kg/head) | Without Fermentation | 6.91a | 5.78c | 5.77c | 6.15 |
| | Fermentation | 6.33ab | 6.24ab | 6.02bc | 6.20 |
| | Average | 6.62a | 6.01b | 5.90b | |
| Carcass Percentage (%/head) | Without Fermentation | 46.82a | 43.61bc | 42.06c | 44.16 |
| | Fermentation | 44.72b | 44.65b | 43.90b | 44.42 |
| | Average | 45.77a | 44.13b | 42.98b | |
| Non-carcass weight (kg/head) | Without Fermentation | 3.17 | 2.52 | 2.50 | 2.73 |
| | Fermentation | 2.91 | 2.74 | 2.64 | 2.76 |
| | Average | 3.04a | 2.63b | 2.57b | |
| Non-Carcass Percentage (%/head) | Without Fermentation | 21.46 | 19.20 | 18.28 | 19.65 |
| | Fermentation | 20.59 | 19.48 | 19.31 | 19.79 |
| | Average | 21.03a | 19.34b | 18.79b | |
| Internal non-carcass weight (kg/head) | Without Fermentation | 0.828 | 0.584 | 0.689 | 0.700 |
| | Fermentation | 0.718 | 0.654 | 0.641 | 0.671 |
| | Average | 0.773a | 0.619b | 0.665b | |
| Percentage of Internal Non-Carcass (%/head) | Without Fermentation | 5.61 | 4.44 | 5.07 | 5.04 |
| | Fermentation | 5.07 | 4.66 | 4.70 | 4.81 |
| | Average | 5.34a | 4.55b | 4.88b | |
| External non-carcass weight (kg/head) | Without Fermentation | 2.342 | 1.934 | 1.810 | 2.029 |
| | Fermentation | 2.197 | 2.088 | 1.998 | 2.094 |
| | Average | 2.269a | 2.011b | 1.904b | |
| Percentage of non-External Carcass (%/head). | Without Fermentation | 15.85 | 14.76 | 13.21 | 14.61 |
| | Fermentation | 15.52 | 14.82 | 14.61 | 14.98 |
| | Average | 15.69a | 14.79b | 13.91b | |
| Edible non-carcass weight (kg/head). | Without Fermentation | 1.983 | 1.645 | 1.712 | 1.780 |
| | Fermentation | 1.795 | 1.763 | 1.703 | 1.754 |
| | Average | 1.889a | 1.704b | 1.707b | |
| Percentage of non-edible carcasses (%/head). | Without Fermentation | 13.43 | 12.45 | 12.48 | 12.79 |
| | Fermentation | 12.67 | 12.65 | 12.41 | 12.58 |
| | Average | 13.05 | 12.55 | 12.45 | |

Description - Different letters towards the row indicate significant differences ($P < 0.05$).

The results of this study showed that the slaughter weight of each treatment did not show significant differences, but the carcass weight, percentage, non-carcass weight, and percentage (internal non-carcass, external non-carcass, and non-carcass fit for consumption) differed along with the increasing level of provision of fermented coconut pulp and non-fermented coconut pulp. This was in accordance with Chanjula and Cherdong (2018) and Obeidat et al. (2020) that nutrition is a factor that could affect livestock carcass production.

The results of the LSD Test showed that the carcass weight of goats that did not receive the addition of fermented coconut pulp and without fermentation in the concentrate composition was significantly higher ($P < 0.05$) compared to 10% unfermented coconut pulp, 20% fermented coconut pulp, and unfermented coconut pulp, but not significantly different ($P > 0.05$) with 10% fermented coconut pulp. A total of 10% unfermented coconut pulp was also not significantly different ($P > 0.05$) with 20% fermented coconut pulp and 20% unfermented coconut pulp in the concentrate.

This was caused by various factors and thought to affect the proportion of carcass weight, especially the nutritional content of feed consumed by livestock. The nutritional content of the concentrate given was 14.01% and decreased with the addition of fermented coconut pulp and unfermented coconut pulp as a substitute for

concentrate. Livestock carcass growth was influenced by many factors, namely genetics, gender, castration, physiology, age, body weight, and feed given both in quantity and quality (Fthenakis and Papadopoulos 2018; Purnami et al. 2021; Spandan et al. 2022). However, gender, breed of livestock, and nutrition were factors that greatly influenced the growth of carcass components (Bambou et al. 2021; Ale et al. 2022; Campelo-Lima et al. 2022; Farid et al. 2023).

The results of the LSD Test showed that the percentage of goat carcasses that did not receive the addition of raw coconut pulp in the concentrate composition was significantly higher ($P < 0.05$) compared to other treatments, while LSD that did not receive fermented coconut pulp was not significantly different ($P > 0.05$) from those that received 10% raw and fermented coconut pulp, and 20% fermented coconut pulp. The reports that received 10% unfermented coconut pulp were not significantly different from those given fermented coconut pulp and 20% unfermented coconut pulp.

The decreasing carcass percentage with increasing coconut pulp as a substitute for concentrate, both unfermented and fermented, was due to the nutritional quality obtained by goats during the study as shown in Table 2. Therefore, the carcass percentage was closely related to the nutritional quality consumed by livestock (Njoga et al. 2021; Ncube et al. 2022). Nutrition was one

of the important factors affecting body weight (Kahi and Wasike 2019; Berthel et al. 2022; Charis et al. 2022; Nelson et al. 2023). When the feed quality was good and the amount of consumption was sufficient, it accelerated growth and eventually increased body weight gain which caused high slaughter weight (Coloma-García et al. 2020; Tahuk and Bira 2023). Faster-growing livestock was more efficient in converting food into body weight gain, therefore increasing carcass weight and subsequently affecting carcass percentage (Reynolds et al. 2020; Dahmer et al. 2022; Wang et al. 2023; Yang et al. 2023).

The LSD Test results showed that the non-carcass weight and percentage (internal, external and edible) of goats that did not receive additional coconut pulp in the concentrate composition were significantly ($P < 0.05$) higher than the treatments that received 10% and 20% coconut pulp. However, those received 10% and 20% coconut pulp did not show any significant difference ($P > 0.05$) in the non-carcass weight and percentage (internal, external, and edible) of female Kacang goats.

Growth and development could result in changes in body weight and measurements, including live, carcass, and non-carcass weight. When there was an increase in live weight, there was an increase in carcass and non-carcass weight (García-Muñiz et al. 2019; Vorlaphim et al. 2021; Zhan et al. 2023; Cai et al. 2023). This report showed that the slaughter weight was not significantly different between treatments however the weight of carcass components and non-carcass weight also did not show significant differences, but tended to decrease with increasing provision of unfermented and fermented coconut pulp. This was supported by Shen et al. (2021); Keçici et al. (2022) that changes in carcass components with increasing carcass weight were due to differential growth of carcass tissue, however changes in carcass components were proportional to the increase in carcass weight.

The relative growth of some non-carcass components was the same as the relative growth rate of the livestock body (Mohammed et al. 2012). Salim and Abdulkareem (2019); Hundal et al. (2020); Alkass et al. (2023) stated that the rumen, reticulum, and omasum increased rapidly in weight in early postnatal life, but the percentage of the digestive tract decreased when it reached maturity. Furthermore, the small intestine developed faster than the large intestine and abomasum. The weight of the digestive and metabolic organs varied greatly depending on the physiological status and the food given (Al-Owaimer et al. 2013; Khatun et al. 2019).

Conclusion

Adding coconut pulp into the feed composition for Kacang goats is feasible. This is because the addition of coconut pulp both fermented and non-fermented, at various levels did not significantly affect body weight gain, dry matter consumption, dry matter use efficiency, body weight gain, respiration rate, pulse rate, or other parameters such as white blood cell count, red blood cell count, haemoglobin level, haematocrit value, slaughter weight, carcass weight, carcass weight, non-carcass weight, and non-carcass weight (internal, external, and edible). In addition, feeding fermented and unfermented coconut pulp in the ration had no significant effect on body weight,

respiration rate, and pulse rate. The only result obtained was that the provision of coconut pulp in the ration significantly affected leucocyte count, slaughter weight, carcass weight, carcass percentage, and non-carcass percentage in Kacang goats.

Conflict of interest: There is no conflict of interest with any organization or party about the anything discussed in this manuscript.

Acknowledgment: The experiment was funded through the Publication Riset Grant provided by Tadulako University for 2022, as per the "Decision of the Research and Community Service Institute with contract No: 769.ak/UN.28.2/PI/2022". The study was conducted with the support of the staff technical assistance at the Animal Production Laboratory, Faculty of Animal and fishery, Tadulako University Palu.

Author's Contribution: Padang, Sri Wulan, and Nirwana designed the research concept and sought funding. Fatmawati and Zainal carried out field and laboratory practices. Harmoko and Naharuddin drafted the script. All authors have read and approved the final manuscript.

REFERENCES

- Agustina DK, Suyadi S, Nugartiningih VMA and Kuswati K, 2023. Hematological profile and leukocyte differentiation of madura cattle and their crosses at different sexes and ages. *Advances in Animal and Veterinary Sciences* 11(7): 1193-1199. <https://doi.org/10.17582/journal.aavs/2023/11.7.1193.1199>
- Abdullah S, Padang P, Sagaf S, Naharuddin N, Cakrawati SW and Harmoko H, 2023. The impact of *Lannea coromandelica* leaf flour supplementation on the production performance and hematological values of local goats. *International Journal of Design & Nature and Ecodynamics* 18(4): 989-994. <https://doi.org/10.18280/ijdne.180427>
- Al-Owaimer A, Suliman G, El-Waziry A, Metwally H and Abouheif M, 2013. Allometric growth patterns of body and carcass components in ardhii Goat. *International Journal of Animal and Veterinary Advances* 5(5): 183-189. <https://doi.org/10.19026/ijava.5.5595>
- Ale KB, Scott J, Okere C, Abrahamsen FW, Gurung R and Gurung NK, 2022. Effects of low-fat distillers dried grains with solubles supplementation on growth performance, rumen fermentation, blood metabolites, and carcass characteristics of kiko crossbred wether goats. *Animals* 12(23): 1-13. <https://doi.org/10.3390/ani12233318>
- Alissandra PBM, Yuwalee U and Rameshprabu R, 2019. Development of a novel sequential pretreatment strategy for the production of bioethanol from coconut pulp residue. *Maejo International Journal of Energy and Environmental Communication* 1(3): 20-28. <https://doi.org/10.54279/mijeec.v1i3.244920>
- Alkass JE, Yateem CAM, Al-Sherwany DAO and Mustafa KNS, 2023. Allometry growth coefficients of carcass and non-carcass components in small ruminants: A review. *Mesopotamia Journal of Agriculture* 51(2): 25-35. <https://doi.org/10.33899/magrj.2023.137250.1227>
- Anggita A, Mudawamah M and Sumartono S, 2023. Analisis komponen darah dari berbagai fase reproduksi induk domba (*Ovis aries*) Sapudi. *Bioscientist: Jurnal Ilmiah Biologi* 11(2): 1311-1319. <https://doi.org/10.33394/bioscientist.v11i2.9107>
- Antoniani C, Meneghini V, Lattanzi A, Felix T, Romano O,

- Magrin E, Weber L, Pavani, G., Hoss S El, Kurita R, Nakamura Y, Cradick TJ, Lundberg AS, Porteus M, Amendola M, Nemer wel, Cavazzana M, Mavilio F and Miccio A, 2018. Induction of fetal hemoglobin synthesis by CRISPR/Cas9-mediated editing of the human β -globin locus. *Blood* 131(17): 1960-1973. <https://doi.org/10.1182/blood-2017-10-811505>
- Bambou JC, Cériac S, Liméa L, Arquet R, Bocage B and Alexandre G, 2021. Impact of diet supplementation and age at slaughter on carcass characteristics of creole goats. *Frontiers in Veterinary Science* 8: 671948. <https://doi.org/10.3389/fvets.2021.671948>
- Berthel R, Simmler M, Dohme-Meier F and Keil N, 2022. Dairy sheep and goats prefer the single components over the mixed ration. *Frontiers in Veterinary Science* 9: 1017669. <https://doi.org/10.3389/fvets.2022.1017669>
- Bijanti R, Eliyani H and Soeharsono, 2011. Parameter hematologi kambing Kacang desa Mojasarirejo Driyorejo Gresik 4(3): 187–190.
- Bradford BJ and Contreras GA, 2024. Adipose Tissue inflammation: linking physiological stressors to disease susceptibility. *Annual Review of Animal Biosciences* 12(1): 261-281. <https://doi.org/10.1146/annurev-animal-021122-113212>
- Cai L, Li M, Zhou S and Xu Q, 2023. The mixture of *saccharomyces cerevisiae* and *clostridium butyricum* could promote rumen fermentation and improve the growth performance of goats in hot summer. *Metabolites* 13(1): 140. <https://doi.org/10.3390/metabo13010104>
- Campelo-Lima VH, Antas-Urbano S, Andrade-Ferreira M, Cavalcante-Ribeiro PH, Gomes-Bezerra J, Cariri-Chagas JC, Moreira-aguiar E and Nascimento-Rangel AH, 2022. Sugarcane bagasse or elephant grass hay in diets for goats: Performance, feeding behavior and carcass characteristics. *Revista Colombiana de Ciencias Pecuarias* 35(1): 49-59. <https://doi.org/10.17533/udea.rccp.v35n1a05>
- Chanjula P and Cherdthong A, 2018. Effects of crude glycerin from waste vegetable oil in diets on performance and carcass characteristics of feedlot goats. *Asian-australasian Journal of Animal Sciences* 31(4): 514-525. <https://doi.org/10.5713/ajas.17.0194>
- Charis BZ, Pagalay U, Hariyadi MA and Wajdi MF, 2022. Optimalisasi formula kandungan zat bahan pakan domba dan kambing dengan multivariate linear regression. *Jurnal Ilmiah Biosaintropis (Bioscience-Tropic)* 8(1): 46-55. <https://doi.org/10.33474/e-jbst.v8i1.492>
- Choi JH, Jung TH, Seo HJ and Han KS, 2021. Development Of Livestock Feed Additives Using Porcine Blood Cells. *Journal of animal Physiology and animal Nutrition* 105 (3): 478-484. <https://doi.org/10.1111/jpn.13498>
- Coloma-García W, Mehaba N, Such X, Caja G and Salama AAK, 2020. Effects of cold exposure on some physiological, productive, and metabolic variables in lactating dairy goats. *Animals* 10(12): 2383. <https://doi.org/10.3390/ani10122383>
- Corrales-Hernández A, Mota-Rojas D, Guerrero-Legarreta I, Roldan-Santiago P, Rodríguez-Salinas S, Yáñez-Pizaña A, de la Cruz L, González-Lozano M and Mora-Medina P, 2018. Physiological responses in horses, donkeys and mules sold at livestock markets. *International Journal of Veterinary Science and Medicine* 6(1): 97-102. <https://doi.org/10.1016/j.jivsm.2018.03.002>
- Costa TC, Moura FH, Souza RO, Lopes MM, Fontes MMS, Serão NVL, Sanglard L P, Du M, Gionbelli MP and Duarte MS, 2019. Effect of maternal feed restriction in dairy goats at different stages of gestation on skeletal muscle development and energy metabolism of kids at the time of births. *Animal Reproduction Science* 206: 46-59. <https://doi.org/10.1016/j.anireprosci.2019.05.006>
- Crilly JP and Plate P, 2022. Anaemia in lambs and kids reared indoors on maternal milk and the impact of iron supplementation on haemoglobin levels and growth rates. *Animals* 12(14): 1863. <https://doi.org/10.3390/ani12141863>
- Dahmer PL, McDonald FB, Chun CKY, Zumbaugh CA, Jones CK, Crane AR, Kott T, Lattimer JM and Chao MD, 2022. Evaluating the impact of feeding dried distillers grains with solubles on Boer goat growth performance, meat color stability and antioxidant capacity. *Translational Animal Science* (2): 1-9. <https://doi.org/10.1093/tas/txac060>
- Elnaggar MM, Abdellrazeq GS, Mack V, Fry LM, Davis WC and Park KT, 2016. Characterization and use of new monoclonal antibodies to CD11c, CD14, and CD163 to analyze the phenotypic complexity of ruminant monocyte subsets. *Veterinary Immunology and Immunopathology* 178: 57-63. <https://doi.org/10.1016/j.vetimm.2016.06.010>
- Elyana P, Pangastuti A and Nugraheni ER, 2018. The effect of adding coconut pulp resulting from aspergillus oryzae fermentation in commercial feed on the growth of Nile tilapia (*Oreochromis niloticus*). *Cell Biology and Development* 2(1): 33-42. <https://doi.org/10.13057/cellbioldev/v020105>
- Emilia I, Putri YP, Novianti D and Niarti M, 2021. Pembuatan virgin coconut oil (VCO) dengan cara fermentasi di desa gunung megang kecamatan gunung megang muara enim. *Sainmatika: Jurnal Ilmiah Matematika Dan Ilmu Pengetahuan alam* 18(1): 88-92. <https://doi.org/10.31851/sainmatika.v17i3.5679>
- Farid MU, Jaspal MH, Luqman N, asghar B, Ijaz M, Yar MK, Ali Taseer MS, Iqbal Z, Naem MA and Nasir J, 2023. Age, sex, and chilling effects on goat meat. *Italian Journal of Food Science* 35(4): 12-20. <https://doi.org/10.15586/ijfs.v35i4.2195>
- Francisco PP, Evelin TT, Victoria CB and Daniela RF, 2023. Hematological alterations in farmers exposed to organophosphate pesticides. *Salud, Ciencia y Tecnología* 3: 568. <https://doi.org/10.56294/SALUDCYT2023568>
- Fthenakis GC and Papadopoulos E, 2018. Impact of parasitism in goat production. *Small Ruminant Research* 163: 21-25. <https://doi.org/10.1016/j.smallrumres.2017.04.001>
- Fu B, Gao H, Yu Y, Zhao Q, Wu H, Wang D, Li Z, Zhang Y, Chen L and Leng J, 2023. Crude protein requirement of yunshang black goats during growth period. *Chinese Journal of Animal Nutrition* 35(3): 1749-1757. <https://doi.org/10.12418/CJAN2023.165>
- García-Muñiz JG, Ramírez-Valverde R, Núñez-Domínguez R and Hidalgo-Moreno JA, 2019. Dataset on growth curves of Boer goats fitted by ten non-linear functions. *Data in Brief* 23: 103672. <https://doi.org/10.1016/j.dib.2019.01.020>
- Gonzalez-Rivas PA, Chauhan SS, Ha M, Fegan N, Dunshea FR and Warner RD, 2020. Effects of heat stress on animal physiology, metabolism, and meat quality: A review. *Meat Science* 162: 108025. <https://doi.org/10.1016/j.meatsci.2019.108025>
- Salim AH and Abdulkareem AA, 2019. CAST/MspI gene polymorphism and its impact on growth performance and carcass traits of Shami goats breed in Iraq. In *Journal of Physics: Conference Series* 1294(9), p. 092015). IOP Publishing. <https://doi.org/10.1088/1742-6596/1294/9/092015>
- Harmoko H and Padang P, 2019. Kondisi performa dan status fisiologis kambing kacang dengan pemberian pakan tepung daun jarak (*Jatropha gossypifolia*) fermentasi. *Indonesian Journal of Animal Science* 21(3): 183-191. <https://doi.org/10.25077/jpi.21.3.183-191.2019>
- Härter CJ, Lima LD, Castagnino DS, Rivera AR, Nunes AM, Sousa SF, Liesegang A, Resende KT and Teixeira IAMA, 2017. Mineral metabolism of pregnant goats under feed restriction. *Animal Production Science* 57(2): 290-300. <https://doi.org/10.1071/AN14471>
- Hundal JS, Wadhwa M and Bakshi MPS, 2020. Effect of supplementing herbal feed additive anethum sowa on

- nutrient utilization, productive performance and carcass characteristics of male beetal kids. *Animal Nutrition and Feed Technology* 20(1): 25-38. <https://doi.org/10.5958/0974-181X.2020.00003.7>
- Hussen J and Al-Sukruwah MA, 2022. The impact of the animal housing system on immune cell composition and function in the blood of dromedary camels. *Animals* 12(3): 317-324. <https://doi.org/10.3390/ani12030317>
- Iftitah D, arisandi B, Widyani RR and Juniah J, 2022. Physiological conditions of broiler chickens during transportation with vitamin treatment and distance difference. *Jurnal Ilmu-Ilmu Peternakan* 32(3): 313-327. <https://doi.org/10.21776/ub.jiip.2022.032.03.02>
- Kahi AK and Wasike CB, 2019. Dairy goat production in sub-Saharan africa: Current status, constraints and prospects for research and development. *Asian-australasian Journal of Animal Sciences* 32(8): 1266-1273. <https://doi.org/10.5713/ajas.19.0377>
- Keçici PD, Yalçintan H and ekiz B, 2022. Allometric growth of non-carcass components and hind limb tissue composition in Saanen goat kids reared under natural and artificial systems. *Journal of Research in Veterinary Medicine* 41(1): 20-26. <https://doi.org/10.30782/jrv.m.1036642>
- Kertawirawan IPA, Budiari NLG, Adijaya IN and Kusumadewi MR, 2022. Identification of concentrate quality from local raw materials through hematology profile on bali cattle. *Buletin Veteriner Udayana* 14(4): 443-449. <https://doi.org/10.24843/bulvet.2022.v14.i04.p17>
- Khaidir, Nasruddin and Syahputra D, 2015. Pengolahan ampas kelapa dalam menjadi biodiesel pada beberapa variasi konsentrasi katalis kalium hidroksida (KOH). *Jurnal Samudera* 9 (2): 77-92 <https://repository.unimal.ac.id/2026/>
- Khatun A, Sarker MB, Islam MA, Bari MS, Amin MR and Moniruzzaman M, 2019. Growth, carcass characteristics and organ development of goat kids fed different levels of cow milk. *Journal of the Bangladesh Agricultural University* 17(4): 551-559. <https://doi.org/10.3329/jbau.v17i4.44625>
- Kholishah N, Mulyaningsih S and Firyanto R, 2021. Pembuatan biodiesel dari ampas kelapa dengan metode transesterifikasi in-situ dan katalis kalium hidroksida (KOH). *CHEMTAG Journal of Chemical Engineering* 2(2): 44-49. <https://doi.org/10.56444/cjce.v2i2.1842>
- Kristianto LK, 2023. Potensi ampas kelapa sebagai bahan pakan ternak alternatif di kalimantan timur. *Warta BSIP Perkebunan* 1(1): 17-21. <https://epublikasi.pertanian.go.id/berkala/wartabun/article/view/1504>
- Laksono J, Karyono T and Haniati H, 2023. Nutritional value of coconut pulp (*cocos nuciferal*) fermented using aspergillus niger at different times as a poultry feed. *Jurnal Ilmu Dan Teknologi Peternakan Indonesia (JITPI) Indonesian Journal of Animal Science and Technology* 9(1): 41-48. <https://doi.org/10.29303/jitpi.v9i1.183>
- Lambrech NJ, Wilson ML, Baylin A, Folsom G, Naabah S, Eisenberg JNS, Adu B and Jones AD, 2021. Associations between livestock ownership and lower odds of anaemia among children 6-59 months old are not mediated by animal-source food consumption in Ghana. *Maternal and Child Nutrition* 17(3): 1-15. <https://doi.org/10.1111/mcn.13163>
- Londok J, Laihad JT and Kowel YHS, 2022. Prebiotik manan oligosakarida dari ampas kelapa dan perannya dalam nutrisi ayam pedaging-review. *Prosiding Seminar* 9: 144-152
- Loučka R, Jančík F, Kumprechtová D, Koukolová V, Kubelková P, Tyrolová Y, Výborná A, Joch M, Jambor V, Synková H, Malá S, Nedělník J, Lang J and Homolka P, 2023. Using precision livestock farming for dairy herd management. *Czech Journal of Animal Science* 68(3): 111-121. <https://doi.org/10.17221/180/2022-CJAS>
- Marhamah SU, Akbarillah T and Hidayat H, 2019. Kualitas nutrisi pakan konsentrat fermentasi berbasis bahan limbah ampas tahu dan ampas kelapa dengan komposisi yang berbeda serta tingkat akseptabilitas pada ternak kambing. *Jurnal Sain Peternakan Indonesia* 14(2): 145-153. <https://doi.org/10.31186/jspi.id.14.2.145-153>
- Mohammed S, Urge M, Animut G, Awigeche K, Abebe G and Goetsch AL, 2012. Effects of level of concentrate supplementation on growth performance of arsi-Bale and Boer × arsi-Bale male goats consuming low-quality grass hay. *Tropical Animal Health and Production* 44(6): 1181-1189. <https://doi.org/10.1007/s11250-011-0056-2>
- Naser A, Padang P, Sagaf S, Abdullah S, Naharuddin N, Wulan S, Nirwana N, Zainal Z and Mustafa M, 2023. The impact of processing methods and levels of *Jatropha gossypifolia* L. leaves on productivity of goats, considering distance. *International Journal of Design & Nature and Ecodynamics*, 18(3): 653-659. <https://doi.org/10.18280/ijdne.180317>
- Navarro G, Bravo V, Gallo C and Phillips CJC, 2019. Physiological and behavioural responses of cattle to high and low space, feed and water allowances during long distance transport in the south of Chile. *Animals* 9(5): 229. <https://doi.org/10.3390/ani9050229>
- Ncube KT, Dzomba EF, Rosen BD, Schroeder SG, Van Tassel CP and Muchadeyi FC, 2022. Differential gene expression and identification of growth-related genes in the pituitary gland of South african goats. *Frontiers in Genetics* 13: 811193. <https://doi.org/10.3389/fgene.2022.811193>
- Nelson NK, Sivasabari K, Ankitha Indu J, Krishnan D, Anjali MR, Akhil PR, Pran M, Nainu F, Praveen SV, Singh P, Chopra H, Chakraborty S, Dey A, Dhama K and Chandran D, 2023. Beneficial impacts of goat milk on the nutritional status and general well-being of human beings: Anecdotal evidence. In *Journal of Experimental Biology and Agricultural Sciences* 11(1): 1-15. [https://doi.org/10.18006/2023.11\(1\).1.15](https://doi.org/10.18006/2023.11(1).1.15)
- Nirwana Wulan S and Zainal, 2021. Complete silage of forage corn and king grass to weight gain and physiological status of goat and sheep. *Jurnal Ilmiah Agrisains* 22(2): 41-46. <https://doi.org/10.22487/jiagrisains.v22i2.2021.74-80>
- Njoga EO, Ezenduka EV, Ogbodo CG, Ogbonna CU, Jaja IF, Ofomatah AC and Okpala COR, 2021. Detection, distribution and health risk assessment of toxic heavy metals/metalloids, arsenic, cadmium, and lead in goat carcasses processed for human consumption in south-eastern Nigeria. *Foods* 10(4): 798. <https://doi.org/10.3390/foods10040798>
- Nohara M, Hisaeda K, Ono T, Inoue Y, Ogawa K, Hata A, Sibano K, Nagahata H and Fujitani N, 2022. The relationships between environmental parameters in livestock pen and physiological parameters of Holstein dairy cows. *Journal of Veterinary Medical Science* 84(7): 964-977. <https://doi.org/10.1292/jvms.22-0187>
- Nsahlai IV, Goetsch AL, Luo J, Johnson ZB, Moore JE, Sahlu T, Ferrell CL, Galyean ML and Owens FN, 2004. Metabolizable protein requirements of lactating goats. *Small Ruminant Research* 53(3): 327-337. <https://doi.org/10.1016/j.smallrumres.2004.04.004>
- Obeidat MD, Obeidat BS, Nusairat B and Al-Shareef R, 2020. Fattening performance and carcass traits of Baladi and Shami-Baladi kids. *Veterinary World* 13(1): 21-25. <https://doi.org/10.14202/vetworld.2020.21-25>
- Oktaviana A, Verdian AH and Aziz R, 2022. Supplementary of coconut pulp extract to feed to increase growth of saline tilapia. *Jurnal Perikanan Tropis* 9(1): 47-55. <https://doi.org/10.35308/jpt.v9i1.4128>
- Ouma GO, 2019. Growth performance, metabolic efficiency and nutrient utilization of BALB/C mice fed with diet made from leftovers from *cocos nucifera* pulp. *American Journal of Bioscience* 7(1): 1-6. <https://doi.org/10.11648/j.ajbio.20190701.11>
- Palulungan JA, Saragih EW, Purwaningsih and Noviyanti N, 2022. Dampak penambahan lamtoro (*Leucaena*

- leucocephala*) pada pakan terhadap status fisiologis ternak kambing kacang (*capra aegragus hircus*). jurnal ilmu peternakan dan veteriner tropis. Journal of Tropical animal and Veterinary Science 12(1): 9-15. <https://doi.org/10.46549/jipvet.v12i1.281>
- Panjaitan D, 2021. Potensi pemanfaatan limbah ampas kelapa sebagai sumber pangan atau bahan substitusi makanan kesehatan. Jurnal Riset Teknologi Pangan Dan Hasil Pertanian (RETIPA) 1(2): 63-68. <https://doi.org/10.54367/retipa.v1i2.1209>
- Purnami NA, Purbowati E and Rianto E, 2021. Meat production of young and adult kejobong goats fed diets with different protein concentration. Jurnal Ilmu-Ilmu Peternakan 31(1): 52-59. <https://doi.org/10.21776/ub.jiip.2021.031.01.07>
- Radhiah A, Martunis M and Erika C, 2022. Karakteristik fisikokimia dari *virgin coconut oil* (VCO) yang diproduksi dengan metode penggaraman dan fermentasi menggunakan ragi roti. Jurnal Ilmiah Mahasiswa Pertanian 7(2): 431-436. <https://doi.org/10.17969/jimfp.v7i2.20160>
- Rahardja DP, Toleng AL and Lestari VS, 2011. Thermoregulation and water balance in fat-tailed sheep and kacang goat under sunlight exposure and water restriction in a hot and dry area. Animal 5(10): 1587-1593. <https://doi.org/10.1017/S175173111000577>
- Rajion MA, Mohamed Saat I, Zulkifli I and Goh YM, 2001. The effects of road transportation on some physiological stress measures in goats. Asian-australasian Journal of Animal Sciences 14(9): 1250-1252. <https://doi.org/10.5713/ajas.2001.1250>
- Reynolds D, Min BR, Gurung N, mcelhenney W, Lee JH, Solaiman S and Bolden-Tiller O, 2020. Influence of tannin-rich pine bark supplementation in the grain mixes for meat goats: Growth performance, blood metabolites, and carcass characteristics. Animal Nutrition 6(1): 85-91. <https://doi.org/10.1016/j.aninu.2019.09.003>
- Roza E, Aritonang SN, Yellita Y, Susanty H and Rizqan, 2024. Hematology and milk production of murrah buffaloes with local forage-based feeding in kapau village, agam regency, West Sumatra, Indonesia. Advances in Animal and Veterinary Sciences 12(3): 451-456. <https://doi.org/10.17582/journal.aavs/2024/12.3.451.456>
- Ruiz-Campillo MT, Molina-Hernandez V, Escamilla A, Stevenson M, Perez J, Martinez-Moreno A, Donnelly S, Dalton JP and Cwiklinski K, 2017. Immune signatures of pathogenesis in the peritoneal compartment during early infection of sheep with *Fasciola hepatica*. Scientific Reports 7(1): 1-14. <https://doi.org/10.1038/s41598-017-03094-0>
- Sagaf S, Naser A, Nirwana N, Cakrawati SW, Zainal Z, Mustafa M and Naharuddin N, 2024. Effects of moringa leaf and fruit powder supplementation on hematological parameters and growth performance in Kacang goats. International Journal of Design & Nature and Ecodynamics 19(3): 963-969. <https://doi.org/10.18280/ijdne.190326>
- Sarmin SS, 2020. Pengaruh bunting dan laktasi terhadap hematologi dan mineral kambing peranakan etawa di Kulonprogo, Yogyakarta, Indonesia. Jurnal Sain Veteriner 38(3): 260-268. <https://doi.org/10.22146/jsv.54190>
- Scatà MC, Alhussien MN, Grandoni F, Reale A, Zampieri M, Hussen J and De Matteis G, 2023. Hyperthermia-induced changes in leukocyte survival and phagocytosis: a comparative study in bovine and buffalo leukocytes. Frontiers in Veterinary Science 10: 1-14. <https://doi.org/10.3389/fvets.2023.1327148>
- Shen J, Hao Z, Wang J, Hu J, Liu X, Li S, Ke N, Song Y, Lu Y, Hu L, Qiao L, Wu X and Luo Y, 2021. Comparative transcriptome profile analysis of longissimus dorsi muscle tissues from two goat breeds with different meat production performance using RNA-Seq. Frontiers in Genetics 11: 1-13. <https://doi.org/10.3389/fgene.2020.619399>
- Shevchenko L, Mitsevsky M, Shulyak S, Mykhalska V, Poliakovskiy V, Boiarchuk S, Anastasiia I, Vadym K, Dmytro N and Gruntovskiy M, 2023. The effect of colloidal silver on clinical, morphological parameters and mineral composition of blood of calves. Acta Fytotechnica et Zootechnica 26(1): 1336-9245. <https://doi.org/10.15414/afz.2023.26.01.8-14>
- Singh P and Krishnaswamy K, 2022. Sustainable zero-waste processing system for soybeans and soy by-product valorization. Trends in Food Science and Technology 128: 331-344. <https://doi.org/10.1016/j.tifs.2022.08.015>
- Singh SK, Singh VK, Kumari P, Nakade UP and Garg SK, 2018. Trypanosoma evansi induces detrimental immuno-catabolic alterations and condition like type-2 diabetes in buffaloes. Parasitology International 67(2): 140-146. <https://doi.org/10.1016/j.parint.2017.10.015>
- Sipos W, 2019. Shifts in porcine PBMC populations from adolescence to adulthood. Veterinary Immunology and Immunopathology 211: 35-41. <https://doi.org/10.1016/j.vetimm.2019.04.002>
- Soeyono G, Widhyari SD, afifah SLN, Angesti SI, Khoirunnisa I, Khaeriyah I, Maulani RN and Ayu BL, 2020. Mastektomi pada kambing peranakan etawa (*Capra aegragus hircus*). A-RSHI Veterinary Letters 4(2): 29-36. <https://doi.org/10.29244/avl.4.2.29-30>
- Soul W, Mupangwa J, Muchenje V and Mpendulo TC, 2019. Biochemical indices and hematological parameters of goats fed lablab purpureus and vigna unguiculata as supplements to a chloris gayana basal diet. Veterinary and Animal Science 8: 1-9. <https://doi.org/10.1016/j.vas.2019.100073>
- Souza AP, Vargas JAC, Fernandes MHMR, Almeida AK, Resende KT and Teixeira IA MA, 2021. Metabolizable protein: 2. requirements for maintenance in growing saanen goats. Frontiers in Veterinary Science 8: 1-8. <https://doi.org/10.3389/fvets.2021.650203>
- Sutanto TDD, Ratnawati D and HP AM, 2021. Pembuatan *virgin coconut oil* (VCO) dengan metode enzimatis dan fermentasi. Indonesian Journal of community Empowerment and Service (ICOMES) 1(1): 6-14. <https://doi.org/10.33369/icom.es.v1i1.18978>
- Spandan PV, Ruban W, Sejian V, Devaraj C, Silpa MV, Awachat V B, Manjunathareddy GB and Bhatta R, 2022. Heat stress induced changes in the major carcass traits and quantitative expression patterns of selective meat quality determining genes in kanni aadu goats. Food Chemistry 1: 1-10. <https://doi.org/10.1016/j.focha.2022.100053>
- Sudarman A, Hidayati N and Suharti S, 2019. Status nutrisi kerbau betina di peternakan rakyat cibungbulang: pengaruh suplementasi *Indigofera* sp dan galepek terhadap perubahan profil darah. Jurnal Ilmu Nutrisi Dan Teknologi Pakan 17(2): 32-37. <https://doi.org/10.29244/jintp.17.2.32-37>
- Suryani S, Sariyani S, Earnestly F, Marganof M, Rahmawati R, Sevindraja S, Indra MTM and Fudholi A, 2020. A comparative study of virgin coconut oil, coconut oil and palm oil in terms of their active ingredients. Processes 8(4): 1-11. <https://doi.org/10.3390/PR8040402>
- Tahuk PK and Bira GF, 2023. The effect of different feed restriction levels on the performance of young male kacang goats. Advances in Animal and Veterinary Sciences 11(1): 141-149. <https://doi.org/10.17582/journal.aavs/2023/11.1.141.149>
- Tana S, Saraswati TR and Yuniwati EYW, 2018. Hematology and blood chemistry status of most frequently consumed ruminants in community. Biosaintifika 10(2): 341-347. <https://doi.org/10.15294/biosaintifika.v10i2.12714>
- Tresia GE, Trisiana AF and Tiesnamurti B, 2023. Effects of road transportation on some physiological stress measures in anperaan boerka goats. Buletin Peternakan 47(3): 142-151. <https://doi.org/10.21059/buletinpeternak.v47i3.83317>
- Vorlaphim T, Paengkoum P, Purba RAP, Yuangklang C, Paengkoum S and Schonewille JT, 2021. Treatment of rice

- stubble with pleurotus ostreatus and urea improves the growth performance in slow-growing goats. *Animals* 11(4): 1-10. <https://doi.org/10.3390/ani11041053>
- Wang D, Tang G, Zhao L, Wang M, Chen L, Zhao C, Liang Z, Chen J, Cao Y and Yao J, 2023. Potential roles of the rectum keystone microbiota in modulating the microbial community and growth performance in goat model. *Journal of Animal Science and Biotechnology* 14(1): 1-15. <https://doi.org/10.1186/s40104-023-00850-3>
- Widyono I, Sarmin T, Susmiyati B and Suwignyo, 2014. Studi Nilai Hematologik Kambing Kacang. Prosiding KIVNAS Ke-13 PDHI. Palembang
- Yang Y, Hui Y, Guo Z, Song X, Zhu H, Pan C and Lan X, 2023. Investigation of the association between goat DNMT3B gene polymorphism and growth traits. *Animal Biotechnology* 34(7): 2492-2498. <https://doi.org/10.1080/10495398.2022.2101115>
- Yulvianti M, Ernayati W and Tarsono, 2015. Pemanfaatan ampas kelapa sebagai bahan baku tepung kelapa tinggi serat dengan metode *freeze drying*. *Jurnal Integrasi Proses* 5(2): 101-107. <https://dx.doi.org/10.36055/jip.v5i2.246>
- Zerfu TA, Abera A, Duncan A, Baltenweck I and mcneill G, 2023. Ownership of small livestock species, but not aggregate livestock, is associated with an increased risk of anemia among children in ethiopia: a propensity score matching analysis. *Food Science and Nutrition* 11(9): 5157-5165. <https://doi.org/10.1002/fsn3.3474>
- Zhan J, Gu Z, Wang H, Liu Y, Wang L, Huang L, Huo J and Wu Y, 2023. Effects of rutin supplementation on growth performance, slaughter performance, serum parameters, and meat quality of nubian goats. *Animal Science Journal* 94(1): 1-12. <https://doi.org/10.1111/asj.13819>